

INTRODUCTION	
CATALOG	4
STERLING	
QUALITY AND PERFORMANCE ASSURANCE	5
STERLING QUALITY	5
STERLING PERFORMANCE	5
APPLIED BIOSYSTEMS INSTRUMENTS	
STERLING CE PHOSPHORAMIDITES	6
STERLING SOLVENTS/REAGENTS	6
STERLING SUPPORTS	7
AB 3900 POLYSTYRENE MODIFIER COLUMNS	9
EXPEDITE™ INSTRUMENTS	
STERLING CE PHOSPHORAMIDITES	10
STERLING SOLVENTS/REAGENTS	10
STERLING SUPPORTS	11
HIGH THROUGHPUT SYNTHESIS	
HT DNA PHOSPHORAMIDITES	13
MERMADE INSTRUMENTS	
STERLING CE PHOSPHORAMIDITES	14
STERLING SOLVENTS/REAGENTS	14
STERLING SUPPORTS	15
GE HEALTHCARE LIFE SCIENCES INSTRUMENTS	
STERLING CE PHOSPHORAMIDITES	16
STERLING SOLVENTS/REAGENTS	17
DR. OLIGO INSTRUMENTS	
STERLING CE PHOSPHORAMIDITES	18
STERLING SOLVENTS/REAGENTS	18
STERLING SUPPORTS	19
OLIGONUCLEOTIDE PURIFICATION	19
ULTRAMILD DNA SYNTHESIS	
ULTRAMILD CE PHOSPHORAMIDITES	20
ULTRAMILD SUPPORTS	21
ULTRAMILD SOLVENTS/REAGENTS	21
SUPPORTS	
GLEN UNYSUPPORT	22
GLEN UNYSUPPORT FC	23
UNIVERSAL SUPPORT III	24
Q-SUPPORTS	25
UNIVERSAL HybridCPG™ SOLID SUPPORTS	25
HIGH LOAD CPG	27
REAGENTS	
ALTERNATIVE SOLVENTS/REAGENTS	28
5' → 3' SYNTHESIS	
5'-CE PHOSPHORAMIDITES	30
5'-SUPPORTS	31
BACKBONE MODIFICATION	
METHYL PHOSPHONAMIDITES	32
PACE PHOSPHORAMIDITES	33
METHYL PHOSPHORAMIDITES	34
ULTRAMILD SOLVENTS/REAGENTS	34
H-PHOSPHONATE MONOMERS	35
H-PHOSPHONATE REAGENTS	35
THIOPHOSPHORAMIDITES	36
SULFURIZING REAGENTS	37
OLIGONUCLEOTIDE-DIRECTED MUTAGENESIS	
TRIMER PHOSPHORAMIDITES	38

STERLING

BACKBONE MODIFIERS

MINOR BASES

MODIFICATION/LABELLING

RNA AND 2'-OME-RNA

MISCELLANEOUS

TABLE OF CONTENTS

DUPLEX STABILIZATION

C-5 PROPYNE DERIVATIVES AND G-CLAMP	42
BASES AFFECTING DUPLEX STABILITY	42
ZIP NUCLEIC ACIDS (ZNA®)	43
CAPS FOR INCREASED DUPLEX STABILITY AND BASE-PAIRING FIDELITY AT TERMINI	44

EPIGENETICS

DNA METHYLATION	45
-----------------	----

PCR/SEQUENCING UTILITIES

DUPLEX EFFECTS	46
CLEANAMP™ MONOMERS	49
CHAIN TERMINATORS	50

STRUCTURAL STUDIES

STRUCTURE/ACTIVITY RELATIONSHIP	52
HALOGENATED NUCLEOSIDES	55
DEUTERATED NUCLEOSIDES	56
DNA DAMAGE/REPAIR	57
CLICK DNA AND RNA LIGATION	60
5'-LABELLING OF MicroRNAs	60
2'-5' LINKED OLIGONUCLEOTIDES	61
MUTAGENESIS	62
CONVERTIBLE NUCLEOSIDES	63
FLUORESCENT NUCLEOSIDES	64
CAGED NUCLEOSIDES	66
THERAPEUTIC NUCLEOSIDES	67
LARGE SCALE SYNTHESIS	67

ULTRAMILD DNA SYNTHESIS

ULTRAMILD CE PHOSPHoramidites	68
ULTRAMILD SUPPORTS	69
ULTRAMILD SOLVENTS/REAGENTS	69

MODIFIERS

TERMINUS MODIFIERS	70
SEQUENCE MODIFIERS	73
3'-MODIFIERS	75
CHEMICAL PHOSPHORYLATION	78
ALDEHYDE MODIFICATION	79
SPACER MODIFIERS	80
DENDRIMERS	81
BRANCHING PHOSPHoramidite	81
PHOTOCLEAVABLE MONOMERS	82
CONJUGATION USING CLICK CHEMISTRY	83
OLIGO-CLICK KITS	84
COPPER-FREE CLICK CHEMISTRY	86

LABELLING

SERINOL REAGENTS FOR MODIFICATION AND LABELLING	89
DABCYL LABELLING	91
BIOTIN LABELLING	92
FLUORESCCEIN LABELLING	95
FLUORESCCEIN LABELLING (SIMA)	98
RHODAMINE (TAMRA) LABELLING	99
CYANINE LABELLING	100
DYLIGHT™ DYES	101
EPOCH DYES AND QUENCHER	102
BLACK HOLE QUENCHER DYES	104
BLACKBERRY® QUENCHER (BBQ-650®)	106
ACRIDINE LABELLING	107
DNP LABELLING	107
CHOLESTEROL LABELLING	108
TOCOPHEROL LABELLING	108
STEARYL LABELLING	108
PSORALEN LABELLING	109
EDTA LABELLING	110
FERROCENE LABELLING	110
METHYLENE BLUE LABELLING	110

LABELLING (CONT)	
LABELLING WITH METAL CHELATES	111
LABELLING WITH POLYAROMATIC HYDROCARBONS	111
PUROMYCIN CPG	112
QUENCHED AUTOLIGATION (QUAL) PROBES	112
LABELLING FOR PHOTO-REGULATION OF OLIGONUCLEOTIDES	113
RNA SUPPORTS	
RNA SUPPORTS FOR 3' MODIFICATION	114
RNA SYNTHESIS	
TOM-PROTECTED RNA PHOSPHoramidites	115
RNA SUPPORTS FOR TOM RNA SYNTHESIS	115
TBDMS-PROTECTED RNA PHOSPHoramidites	117
HT RNA PHOSPHoramidites	117
LC RNA PHOSPHoramidites	117
ULTRAMILD TBDMS RNA PHOSPHoramidites	118
TBDMS RNA SUPPORTS	118
ULTRAMILD SOLVENTS/REAGENTS	119
MINOR RNA BASES	
MINOR RNA PHOSPHoramidites (TOM PROTECTED)	120
RNA SEQUENCE MODIFIER (TOM PROTECTED)	121
MINOR RNA PHOSPHoramidites (TBDMS PROTECTED)	122
MINOR RNA TRIPHOSPHATES	124
2'-OME-RNA SYNTHESIS	
2'-OME-RNA PHOSPHoramidites	125
HT 2'-OME-RNA PHOSPHoramidites	125
ULTRAMILD 2'-OME-RNA	126
ULTRAMILD SOLVENTS/REAGENTS	126
MINOR 2'-OME-RNA PHOSPHoramidites	128
2'-F RNA SYNTHESIS	
2'-F-RNA MONOMERS	130
2'-F ANA SYNTHESIS	
2'-F-ARABINONUCLEIC ACID (2'-F-ANA)	131
NAIM	
INTERFERENCE MAPPING	132
A ANALOGS	133
NUCLEOSIDE α -THIOTRIPHOSPHATES	135
INTERNALLY QUENCHED NUCLEOTIDE FLUORESCENT REPORTERS	137
PURIFICATION	
GLEN-PAK™ PURIFICATION	138
POLY-PAK™ PURIFICATION	139
FLUORO-PAK™ PURIFICATION	140
GLEN GEL-PAK™ DESALTING	141
OLIGO-AFFINITY SUPPORT	142
PHYSICAL DATA	
INDEX	
GENERAL INFORMATION	
ORDERING	163
DISCOUNTS	163
TERMS AND CONDITIONS OF SALE	163
PATENTS	163

CATALOG

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

Welcome to the Glen Research Catalog containing the most complete selection of products for DNA and RNA research. This catalog format has proved to be popular with our customers and allows relatively simple navigation through our wide array of product offerings. The Table of Contents at the beginning and the Index at the end of the Catalog are the most comprehensive we have produced. The Physical Data section covers all products, including supports. This section is also useful for resolving a product name from a known catalog number. There are always limitations to printed catalogs in a fast-moving technology sector and a complete and up-to-date catalog is also maintained on our web site.

All minor bases, modifiers and RNA products are packaged for Applied Biosystems instruments. We can provide vials and columns for a wide variety of other instruments. As shown in the table to the left, we can accommodate catalog numbers for unusual products to fit all popular instruments. For example, if you wish to order 0.25g of dI-CE Phosphoramidite for an Applied Biosystems instrument, the catalog number is 10-1040-02. If you wish the same product for an Expedite instrument, the catalog number, derived from the table to the left, would be 10-1040-02E. For columns, a pack of four dl columns for an Expedite would be 20-2190-42E. The table to the left is reproduced on all relevant spreads of this catalog.

We are unique in conducting a QC test for supports to show the length of oligo that can be prepared before a drop-off in coupling due to steric effects begins to occur. The drop-off point is recorded in the Certificate of Analysis or Analytical Report. The vast majority of our minor base and modification supports are 1000Å CPG, which results in improved performance and the ability to make much longer oligos. Supports which will continue with a specific pore size are shown as such in the catalog. Polystyrene supports are also available for some of our most popular items.

For reasons of quality assurance, we do not transfer powders or oils from stock Applied Biosystems vials to vials for other instruments. Powders may be hygroscopic and electrostatic, making transfer difficult, and oils have to be dissolved and the solvent evaporated. For best performance, it is preferable for the customer to dissolve the product and immediately transfer the solution to the correct instrument vial. Consequently, the product will be delivered in an industry-standard septum-capped vial along with a clean dry vial for the appropriate instrument. We monitor product usage for each instrument type and will attempt to have stock prepackaged for some of the most popular products and configurations.

Glen Research's distributors cover a very significant percentage of countries where oligonucleotide synthesis is commonly practiced. Our vast selection of unusual products is really only comprehensively stocked here in Virginia and some of our web viewers have asked us to set up a direct shipping channel. For them, we offer the eGlen program which is described in the following web link: <http://www.glenresearch.com/Reference/eGlen.html>.

Authorized distributors for Glen Research products are listed below. Other countries not listed are covered by direct sales from our Sterling, USA office.

UK and Ireland

Cambio Ltd
Telephone Number: +44 (0) 1954 210200
Fax Number: +44 (0) 1954 210300
E-mail addresses: support@cambio.co.uk and
orders@cambio.co.uk
Website: <http://www.cambio.co.uk/>

Nordic and Baltic Countries

BioNordikaBergman AS
Telephone Number: +47 23035800
Fax Number: +47 23035801
e-mail address: info@bionordikabergman.no
Website: <http://www.bionordika.no/>

Japan

Nihon Techno Service Co., Ltd.
Telephone Number: +81 29 886 6811
Fax Number: +81 29 870 0210
e-mail address: info@ntsbio.com
Website: <http://www.ntsbio.com/>

Italy

Primm srl
Telephone Number: +39 02 2104031
Fax Number: +39 02 2640355
email address: lucia.ricchiuti@primm.it
Website: <http://www.primm.it/>

Belgium

Eurogentec S.A.
Telephone Number: +32 4 372 74 00
Fax Number: +32 4 372 75 00
e-mail address: info@eurogentec.com
Website: <http://www.eurogentec.com/>

Israel

Eisenberg Bros. Ltd.
Telephone Number: 972-3-9777000
Fax Number: 972-3-9777001
e-mail address: nicoles@eb1.co.il
Website: <http://www.eisenbros.co.il/>

Netherlands

Eurogentec b.v.
Telephone Number: +31 43 352 06 98
Fax Number: +31 43 354 19 65
e-mail address: info@eurogentec.com

Germany

Eurogentec GmbH
Telephone Number: +49 221 258 94 55
Fax Number: +49 221 258 94 54
e-mail address: info@eurogentec.com

France

Eurogentec s.a.
Telephone Number: +33 2 41 73 33 73
Fax Number: +33 2 41 73 10 26
e-mail address: info@eurogentec.com

QUALITY AND PERFORMANCE ASSURANCE

Glen Research has developed and implemented a Quality Management System (QMS) designed to enhance customer satisfaction by focusing on processes for continual improvement and on assurance of conformity to customer needs, with full consideration of applicable regulatory requirements.

STERLING QUALITY


The benchmark for excellence in DNA and RNA synthesis. All Sterling materials must pass stringent purity and identity tests prior to acceptance. Sterling products are formulated, filtered, and packaged in optimal environments using specially cleaned and dried glassware and columns. Color-coded labelling and post-packaging analysis guarantee accuracy and Sterling Quality.




STERLING is a trademark of Glen Research Corporation.

Glen Research offers the highest level of Quality Assurance for reagents for DNA and RNA synthesis - Sterling Quality and Performance. We now apply the Sterling criteria of quality and performance to all of Glen Research's established products.


The common monomers and supports, whose structures are illustrated below, are available for the variety of synthesizers listed on the following pages.




dA-CE Phosphoramidite




dC-CE Phosphoramidite




Ac-dC-CE Phosphoramidite




dG-CE Phosphoramidite




dT-CE Phosphoramidite




dA-Icaa-CPG




dC-Icaa-CPG



Ac-dC-Icaa-CPG



dG-Icaa-CPG



dT-Icaa-CPG




STERLING

QUALITY ASSURANCE

Every batch of these CE Phosphoramidites is tested as follows:

1. **HPLC**
a) Identity is confirmed by comparison with a reference sample.
b) Purity is determined by HPLC to be $\geq 98.0\%$.
2. **TLC**
Purity is verified by TLC.
3. **^{31}P NMR**
Purity is determined by ^{31}P NMR to be $\geq 98\%$.
4. **Coupling Test**
Coupling efficiency is determined to be $\geq 99\%$.
5. **Solution Test**
A 0.1M solution is determined to be clear and free of particulate contamination.
6. **Loss on Drying**
Volatile contaminants are determined to be $\leq 2\%$.



dmF-dG-CE Phosphoramidite

ABI INSTRUMENTS

1. 60mL septum-capped vials used on oldest ABI 380, 381 and 391 instruments. 200mL oxidizer and 450mL deblock screw-capped bottles also used on ABI 380, 381 and 391 instruments.
2. Small screw-capped vials used on ABI 392 and 394 instruments.
3. Larger screw-capped vials used on ABI 392, 394 and 3400 instruments.
4. Large bottles used on ABI 3900 instruments.

STERLING CE PHOSPHORAMIDITES

Glen Research CE (β -cyanoethyl) Phosphoramidites are produced and packaged to ensure the highest performance on DNA synthesizers. Every Glen Research product is accompanied by a Certificate of Analysis and HPLC trace, showing the results of our QC testing. Every Glen Research monomer vial is specially cleaned to eliminate particulate contamination and tested to ensure a tight fit on synthesizers.

Item	Catalog No.	Pack	Price (\$)
dA-CE Phosphoramidite	10-1000-02	0.25g	12.50
	10-1000-05	0.5g	25.00
	10-1000-10	1.0g	50.00
	10-1000-20	2.0g	100.00
	10-1000-40	4.0g	200.00
dC-CE Phosphoramidite	10-1010-02	0.25g	12.50
	10-1010-05	0.5g	25.00
	10-1010-10	1.0g	50.00
	10-1010-20	2.0g	100.00
	10-1010-40	4.0g	200.00
Ac-dC-CE Phosphoramidite	10-1015-02	0.25g	12.50
	10-1015-05	0.5g	25.00
	10-1015-10	1.0g	50.00
	10-1015-20	2.0g	100.00
	10-1015-40	4.0g	200.00
dG-CE Phosphoramidite	10-1020-02	0.25g	12.50
	10-1020-05	0.5g	25.00
	10-1020-10	1.0g	50.00
	10-1020-20	2.0g	100.00
	10-1020-40	4.0g	200.00
dmf-dG-CE Phosphoramidite	10-1029-02	0.25g	12.50
	10-1029-05	0.5g	25.00
	10-1029-10	1.0g	50.00
	10-1029-20	2.0g	100.00
	10-1029-40	4.0g	200.00
dT-CE Phosphoramidite	10-1030-02	0.25g	12.50
	10-1030-05	0.5g	25.00
	10-1030-10	1.0g	50.00
	10-1030-20	2.0g	100.00
	10-1030-40	4.0g	200.00

STERLING SOLVENTS/REAGENTS

All solvents and reagents are prepared to our exacting specifications to ensure the highest synthesis efficiency and are passed through a 0.2 micron filter during packaging to eliminate particulate contamination. Glen Research uses freshly sublimed 1H-tetrazole for premium performance on Applied Biosystems synthesizers.

Item	Catalog No.	Pack	Price (\$)
<i>Activator</i>			
0.45M Tetrazole in Acetonitrile	30-3100-45 ¹	45mL	40.00
	30-3100-52 ²	200mL	100.00
	30-3100-57 ³	450mL	200.00
	30-3100-62 ⁴	2L	760.00
<i>Diluent</i>			
Acetonitrile, anhydrous	40-4050-45	60mL	12.00
	40-4050-50	100mL	16.00

STERLING SOLVENTS/REAGENTS (CONT.)

Item	Catalog No.	Pack	Price (\$)
Cap Mix A			
THF/Pyridine/Ac ₂ O	40-4110-45 ¹	45mL	16.00
	40-4110-52 ²	200mL	30.00
	40-4110-57 ³	450mL	72.00
	40-4110-62 ⁴	2L	325.00
Cap Mix B			
16% MeIm in THF <i>(This Cap B solution is identical to the formulation produced by Applied Biosystems.)</i>	40-4220-45 ¹	45mL	20.00
	40-4220-52 ²	200mL	40.00
	40-4220-57 ³	450mL	96.00
	40-4220-62 ⁴	2L	425.00
Oxidizing Solution			
0.02M I ₂ in THF/Pyridine/H ₂ O	40-4330-52 ^{1,2}	200mL	30.00
	40-4330-57 ³	450mL	72.00
	40-4330-62 ⁴	2L	325.00
Deblocking Mix			
3% TCA/DCM	40-4140-57 ^{1,2}	450mL	36.00
	40-4140-62 ^{3,4}	2L	144.00

STERLING SUPPORTS

All Glen Research CPG supports use the standard long chain alkylamino (lcaa) linker but differ in the glass pore size, 500Å, 1000Å or 2000Å. The 500Å support is appropriate for shorter sequences, while the 1000Å supports perform better in the synthesis of longer (>30-mer) DNA sequences. The 2000Å support is best for very long (>150-mer) oligonucleotides. We have instituted an additional QC test for supports to show the length of oligo that can be prepared before a drop-off in coupling due to steric effects begins to occur. The drop-off point is recorded in the Certificate of Analysis. All Glen Research supports are fully end-capped to ensure that the CPG surface is totally inert, thereby avoiding the introduction of impurity sequences containing deletions at the 3'-terminus.

Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Pack	Price(\$)
dA	dC	dG	dT	dA,dC,dG,dT (1 column of each base)	Ac-dC	dmf-dG		

500Å Columns

20-2100-42	20-2110-42	20-2120-42	20-2130-42	20-2140-42	20-2113-42	4X0.2µm	40.00
20-2100-41	20-2110-41	20-2120-41	20-2130-41	20-2140-41	20-2113-41	4X1µm	60.00
20-2100-13	20-2110-13	20-2120-13	20-2130-13		20-2113-13	1X10µm	100.00

1000Å Columns

20-2101-45	20-2111-45	20-2121-45	20-2131-45	20-2141-45	20-2115-45	20-2129-45	4X40nm	40.00
20-2101-42	20-2111-42	20-2121-42	20-2131-42	20-2141-42	20-2115-42	20-2129-42	4X0.2µm	40.00
20-2101-41	20-2111-41	20-2121-41	20-2131-41	20-2141-41	20-2115-41	20-2129-41	4X1µm	60.00
20-2101-13	20-2111-13	20-2121-13	20-2131-13		20-2115-13	20-2129-13	1X10µm	100.00


ABBREVIATIONS

Ac₂O = Acetic Anhydride
 CE = Cyanoethyl
 CPG = Controlled Pore Glass
 DCM = Dichloromethane
 dmf = dimethylformamide
 I₂ = Iodine
 lcaa = long chain alkylamino
 MeIm = 1-Methylimidazole
 µm = micromole(s)
 nm = nanomole(s)
 TCA = Trichloroacetic Acid
 THF = Tetrahydrofuran

SEE ALSO

Alternative Solvents

p28



STERLING SUPPORTS (CONT.)

Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Pack	Price(\$)
dA	dC	dG	dT	dA,dC,dG,dT (1 column of each base)	Ac-dC	dmf-dG		

2000Å Columns

20-2102-42	20-2112-42	20-2122-42	20-2132-42	20-2142-42			4X0.2µm	40.00
------------	------------	------------	------------	------------	--	--	---------	-------

Low Volume (LV) Polystyrene Columns

26-2100-45	26-2110-45	26-2120-45	26-2130-45	26-2140-45			4X40nm	48.00
26-2100-42	26-2110-42	26-2120-42	26-2130-42	26-2140-42			4X0.2µm	48.00

AB 3900 Polystyrene Columns

26-2600-65	26-2610-65		26-2630-65		26-2629-65	200X40nm		825.00
26-2600-62	26-2610-62		26-2630-62		26-2629-62	200X200nm		825.00

AB 3900 1000Å CPG Columns

20-2101-65			20-2131-65		20-2115-65	20-2129-65	200X40nm	600.00
20-2101-62			20-2131-62		20-2115-62	20-2129-62	200X200nm	650.00
20-2101-61			20-2131-61		20-2115-61	20-2129-61	200X1µm	875.00

500Å Bulk CPG

20-2000-01	20-2010-01	20-2020-01	20-2030-01		20-2013-01		0.1g	9.00
20-2000-02	20-2010-02	20-2020-02	20-2030-02		20-2013-02		0.25g	20.00
20-2000-10	20-2010-10	20-2020-10	20-2030-10		20-2013-10		1.0g	75.00

1000Å Bulk CPG

20-2001-01	20-2011-01	20-2021-01	20-2031-01		20-2015-01	20-2029-01	0.1g	9.00
20-2001-02	20-2011-02	20-2021-02	20-2031-02		20-2015-02	20-2029-02	0.25g	20.00
20-2001-10	20-2011-10	20-2021-10	20-2031-10		20-2015-10	20-2029-10	1.0g	75.00

2000Å Bulk CPG

20-2002-01	20-2012-01	20-2022-01	20-2032-01				0.1g	15.00
20-2002-02	20-2012-02	20-2022-02	20-2032-02				0.25g	30.00
20-2002-10	20-2012-10	20-2022-10	20-2032-10				1.0g	105.00

Item	Catalog No.	Pack	Price(\$)
Empty Synthesis Columns (40nm, 0.2µm, or 1µm) (TWIST™ Style)	20-0030-00	10	60.00
Empty Synthesis Columns (10µm) (TWIST™ Style)	20-0040-00	10	300.00
Replacement Frits (10µm)	20-0040-0F	20	30.00

Product structures are shown on Page 5. TWIST is a trademark of Glen Research Corporation.

AB 3900 1000Å CPG COLUMNS

Glen Research's AB 3900 1000Å CPG columns bring the lower cost of CPG to this platform while maintaining the high synthesis efficiency of 1000Å CPG. Our columns offer the following key attributes:

- No need to change instrument settings
- No need to change software parameters
- Easier handling post-synthesis compared to PS
- High quality 1000Å CPG for optimal synthesis results

BULK CPG LOADING

500Å supports	35-50µmoles/g
1000Å supports	25-40µmoles/g

SEE ALSO

Universal Supports	p22
Q-Supports	p25
High Load Supports	p27

AB 3900 POLYSTYRENE MODIFIER COLUMNS

Some of our more popular minor base and modifier supports are available on polystyrene in columns fully compatible with the Applied Biosystems 3900 synthesizer. These include our popular Universal Support III, which will allow DNA, RNA or LNA oligos to be produced on the 3900 with ANY base at the 3' terminus. Structures and more complete descriptions are found in the relevant catalog sections for each item. AB 3900 columns can be prepared with virtually any of the CPG supports in this catalog. It is no longer necessary to adjust the flow using our AB 3900 CPG columns, as noted in the box to the right. Modified CPG columns are only available in 200 nmole size - simply add 'A' to the regular catalog number to order.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Universal Support III PS			
200 nmole columns	26-5110-52	Pack of 10	100.00
40 nmole columns	26-5110-55	Pack of 10	100.00
Glen UnySupport PS			
200 nmole columns	26-5140-52	Pack of 10	100.00
40 nmole columns	26-5140-55	Pack of 10	100.00
3'-Phosphate PS			
200 nmole columns	26-2900-52	Pack of 10	150.00
40 nmole columns	26-2900-55	Pack of 10	150.00
3'-PT-Amino-Modifier C6 PS			
200 nmole columns	26-2956-52	Pack of 10	220.00
40 nmole columns	26-2956-55	Pack of 10	220.00
3'-(6-FAM) PS			
200 nmole columns	26-2961-52	Pack of 10	300.00
40 nmole columns	26-2961-55	Pack of 10	300.00
3'-DabcyI PS			
200 nmole columns	26-5912-52	Pack of 10	300.00
40 nmole columns	26-5912-55	Pack of 10	300.00
3'-TAMRA PS			
200 nmole columns	26-5910-52	Pack of 10	300.00
40 nmole columns	26-5910-55	Pack of 10	300.00
3'-BiotinTEG PS			
200 nmole columns	26-2955-52	Pack of 10	300.00
40 nmole columns	26-2955-55	Pack of 10	300.00

SEE ALSO

Universal Support III p24
Glen UnySupport p22

AB 3900 1000Å CPG COLUMNS

Glen Research's AB 3900 1000Å CPG columns bring the lower cost of CPG to this platform while maintaining the high synthesis efficiency of 1000Å CPG. Our columns offer the following key attributes:

- No need to change instrument settings
- No need to change software parameters
- Easier handling post-synthesis compared to PS
- High quality 1000Å CPG for optimal synthesis results

STERLING CE PHOSPHORAMIDITES

Glen Research CE (β-cyanoethyl) Phosphoramidites are produced and packaged to ensure the highest performance on DNA synthesizers. Every Glen Research product is accompanied by a Certificate of Analysis and HPLC trace, showing the results of our QC testing. Every Glen Research monomer vial is specially cleaned to eliminate particulate contamination.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
dA-CE Phosphoramidite	10-1000-C2	0.25g	12.50
	10-1000-C5	0.5g	25.00
	10-1000-1C	1.0g	50.00
	10-1000-2C	2.0g	100.00
dC-CE Phosphoramidite	10-1010-C2	0.25g	12.50
	10-1010-C5	0.5g	25.00
	10-1010-1C	1.0g	50.00
	10-1010-2C	2.0g	100.00
Ac-dC-CE Phosphoramidite	10-1015-C2	0.25g	12.50
	10-1015-C5	0.5g	25.00
	10-1015-1C	1.0g	50.00
	10-1015-2C	2.0g	100.00
dG-CE Phosphoramidite	10-1020-C2	0.25g	12.50
	10-1020-C5	0.5g	25.00
	10-1020-1C	1.0g	50.00
	10-1020-2C	2.0g	100.00
dmf-dG-CE Phosphoramidite	10-1029-C2	0.25g	12.50
	10-1029-C5	0.5g	25.00
	10-1029-1C	1.0g	50.00
	10-1029-2C	2.0g	100.00
dT-CE Phosphoramidite	10-1030-C2	0.25g	12.50
	10-1030-C5	0.5g	25.00
	10-1030-1C	1.0g	50.00
	10-1030-2C	2.0g	100.00

STERLING SOLVENTS/REAGENTS

All solvents and reagents are prepared to our exacting specifications to ensure the highest synthesis efficiency and are passed through a 0.2 micron filter during packaging to eliminate particulate contamination. Glen Research uses freshly sublimed 1H-tetrazole for premium performance on Expedite synthesizers. Crystalline tetrazole solutions have been discontinued.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Activator</i>			
0.45M Tetrazole (sublimed) in Acetonitrile	30-3102-66 ¹	60mL	50.00
	30-3102-52 ²	200mL	100.00
	30-3100-57 ²	450mL	200.00
<i>Diluent</i>			
Acetonitrile, anhydrous	40-4050-45	60mL	12.00
	40-4050-50	100mL	16.00

QUALITY ASSURANCE

Every batch of these CE Phosphoramidites is tested as follows:

- HPLC**
a) Identity is confirmed by comparison with a reference sample.
b) Purity is determined by HPLC to be $\geq 98.0\%$.
- TLC**
Purity is verified by TLC.
- ³¹P NMR**
Purity is determined by ³¹P NMR to be $\geq 98\%$.
- Coupling Test**
Coupling efficiency is determined to be $\geq 99\%$.
- Solution Test**
A 0.1M solution is determined to be clear and free of particulate contamination.
- Loss on Drying**
Volatile contaminants are determined to be $\leq 2\%$.

EXPEDITE INSTRUMENTS

- For use on Expedite 8905 instruments.
- For use on Expedite 8909 instruments.

STERLING SUPPORTS (CONT.)

<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
dA	dC	dG	dT	dA,dC,dG,dT (1 column of each base)	Ac-dC	dmf-dG		

2000Å Columns

20-2202-42	20-2212-42	20-2222-42	20-2232-42	20-2242-42			4X0.2µm	40.00
------------	------------	------------	------------	------------	--	--	---------	-------

500Å Bulk CPG

20-2000-01	20-2010-01	20-2020-01	20-2030-01		20-2013-01		0.1g	9.00
20-2000-02	20-2010-02	20-2020-02	20-2030-02		20-2013-02		0.25g	20.00
20-2000-10	20-2010-10	20-2020-10	20-2030-10		20-2013-10		1.0g	75.00

1000Å Bulk CPG

20-2001-01	20-2011-01	20-2021-01	20-2031-01		20-2015-01	20-2029-01	0.1g	9.00
20-2001-02	20-2011-02	20-2021-02	20-2031-02		20-2015-02	20-2029-02	0.25g	20.00
20-2001-10	20-2011-10	20-2021-10	20-2031-10		20-2015-10	20-2029-10	1.0g	75.00

2000Å Bulk CPG

20-2002-01	20-2012-01	20-2022-01	20-2032-01				0.1g	15.00
20-2002-02	20-2012-02	20-2022-02	20-2032-02				0.25g	30.00
20-2002-10	20-2012-10	20-2022-10	20-2032-10				1.0g	105.00

SEE ALSO

Universal Supports
Q-Supports
High Load Supports

p22
p25
p27

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
Empty Synthesis Columns (40nm, 0.2µm)	20-0021-02	10	48.00
Empty Synthesis Columns (1µm)	20-0021-01	10	48.00
Replacement Filters (40nm, 0.2 or 1µm)	20-0021-0F	20	20.00
Empty Synthesis Columns (15µm) (TWIST™ Style)	20-0040-00	10	300.00
Replacement Frits (15µm)	20-0040-0F	20	30.00

Product structures are shown on Page 5. TWIST is a trademark of Glen Research Corporation.
Expedite is a trademark of Applied Biosystems.

HT DNA PHOSPHORAMIDITES

We offer a high quality designation of DNA phosphoramidites destined for high throughput and large-scale synthesis customers. These customers normally require high quality materials produced under the guidelines of a validated quality management system while still being priced aggressively. These products include the usual Glen Research certification and guarantees but they are only available in larger packs or in bulk. As the market evolves, we would expect to expand this line to encompass other bases, modifiers and supports. HT monomers are not subject to regular discounts and a separate HT discount schedule will be set up for our customers. For these products, please request a quote.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>HT DNA Phosphoramidites</i>			
dA-CE Phosphoramidite	10-1000-5HT	5.0g	
dC-CE Phosphoramidite	10-1010-5HT	5.0g	
Ac-dC-CE Phosphoramidite	10-1015-5HT	5.0g	
dG-CE Phosphoramidite	10-1020-5HT	5.0g	
dmf-dG-CE Phosphoramidite	10-1029-5HT	5.0g	
dT-CE Phosphoramidite	10-1030-5HT	5.0g	

HT DNA

Minimum order: 400g total
 Packaged as 5g bottles (or greater)
 or in bulk.
 Products may be combined to 400g
 total.
 Please request a quote.

STERLING CE PHOSPHORAMIDITES

MerMade synthesizers belong to a family of synthesizers, including the column-based MerMade 4, MerMade 6 and 12 instruments and the parallel array synthesizers, MerMade 192 and MerMade 192E, manufactured by BioAutomation Corporation in Plano, TX. Their web site can be found at: <http://www.BioAutomation.com>. Phosphoramidite monomers are packaged in 30mL and 240mL amber bottles for dissolving at a concentration of 1g/20mL and are connected directly to the instrument. Some instruments may also be configured to accept Applied Biosystems serum vials, as shown on page 6.

QUALITY ASSURANCE

Every batch of these CE Phosphoramidites is tested as follows:

- 1. HPLC**
a) Identity is confirmed by comparison with a reference sample.
b) Purity is determined by HPLC to be $\geq 98.0\%$.
- 2. TLC**
Purity is verified by TLC.
- 3. ³¹P NMR**
Purity is determined by ³¹P NMR to be $\geq 98\%$.
- 4. Coupling Test**
Coupling efficiency is determined to be $\geq 99\%$.
- 5. Solution Test**
A 0.1M solution is determined to be clear and free of particulate contamination.
- 6. Loss on Drying**
Volatile contaminants are determined to be $\leq 2\%$.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
dA-CE Phosphoramidite	10-1000-02M	0.25g	12.50
	10-1000-05M	0.5g	25.00
	10-1000-10M	1.0g	50.00
	10-1000-5S	5.0g	250.00
	10-1000-1S	10.0g	500.00
dC-CE Phosphoramidite	10-1010-02M	0.25g	12.50
	10-1010-05M	0.5g	25.00
	10-1010-10M	1.0g	50.00
	10-1010-5S	5.0g	250.00
	10-1010-1S	10.0g	500.00
Ac-dC-CE Phosphoramidite	10-1015-02M	0.25g	12.50
	10-1015-05M	0.5g	25.00
	10-1015-10M	1.0g	50.00
	10-1015-5S	5.0g	250.00
	10-1015-1S	10.0g	500.00
dG-CE Phosphoramidite	10-1020-02M	0.25g	12.50
	10-1020-05M	0.5g	25.00
	10-1020-10M	1.0g	50.00
	10-1020-5S	5.0g	250.00
	10-1020-1S	10.0g	500.00
dmf-dG-CE Phosphoramidite	10-1029-02M	0.25g	12.50
	10-1029-05M	0.5g	25.00
	10-1029-10M	1.0g	50.00
	10-1029-5S	5.0g	250.00
	10-1029-1S	10.0g	500.00
dT-CE Phosphoramidite	10-1030-02M	0.25g	12.50
	10-1030-05M	0.5g	25.00
	10-1030-10M	1.0g	50.00
	10-1030-5S	5.0g	250.00
	10-1030-1S	10.0g	500.00

STERLING SOLVENTS/REAGENTS

All solvents and reagents are prepared to our exacting specifications to ensure the highest synthesis efficiency and are passed through a 0.2 micron filter during packaging to eliminate particulate contamination. Parallel synthesizers typically use 5-ethylthio-1H-tetrazole (ETT) as activator to minimize the chance of crystallization. ETT is used at a concentration of 0.25M in acetonitrile, which is far below the level at which crystallization may occur.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Activator</i> 0.25M ETT in Acetonitrile	30-3140-57	450mL	200.00
	30-3140-61	960mL	365.00
	30-3140-62	2L	760.00

SEE ALSO

Alternative Activators

p28

STERLING SOLVENTS/REAGENTS (CONT.)

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Diluent</i>			
Acetonitrile, anhydrous	40-4050-50	100mL	16.00
<i>Cap Mix A</i>			
THF/Lutidine/Ac ₂ O	40-4010-57	450mL	72.00
	40-4010-61	960mL	154.00
	40-4010-62	2L	325.00
<i>Cap Mix B</i>			
16% MeIm in THF	40-4220-57	450mL	96.00
	40-4220-61	960mL	204.00
	40-4220-62	2L	425.00
<i>Oxidizing Solution</i>			
0.02M I ₂ in THF/Pyridine/H ₂ O	40-4330-57	450mL	72.00
	40-4330-61	960mL	154.00
	40-4330-62	2L	325.00
<i>Deblocking Mix</i>			
3% DCA/DCM	40-4040-57	450mL	36.00
	40-4040-61	960mL	75.00
	40-4040-62	2L	144.00
3% TCA/DCM	40-4140-57	450mL	36.00
	40-4140-61	960mL	75.00
	40-4140-62	2L	144.00

STERLING SUPPORTS

Columns containing 1000Å CPG are available in packs of 200 to fit MerMade plates. Regular 500Å or 1000Å supports, listed on page 8, may also be used to fill the wells of regular 96 well plates. However, this requires each plate to be prepared with each nucleoside accurately in all wells. A universal support clearly removes the need for four specific supports and makes preparing plates straightforward. Glen UnySupport™ 40 nmole frits, as described on page 22, can also be used.

<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
dA	dC	dG	dT	Ac-dC	dmf-dG		
<i>MerMade 1000Å CPG Columns</i>							
20-2001-65	20-2021-65	20-2031-65	20-2015-65	20-2029-65		200X50nm	750.00
20-2001-62	20-2021-62	20-2031-62	20-2015-62	20-2029-62		200X200nm	750.00
20-2001-61	20-2021-61	20-2031-61	20-2015-61	20-2029-61		48X1µm	300.00
<i>Universal Support III</i>							
				20-5110-91		Pack of 96	375.00
				20-5110-92		Pack of 96	250.00
				20-5110-95		Pack of 96	250.00
<i>Glen UnySupport</i>							
				20-5141-91		Pack of 96	375.00
				20-5141-92		Pack of 96	250.00
				20-5141-95		Pack of 96	250.00
<i>Empty MerMade Columns</i>							
				20-0050-05		Pack of 48	200.00
				20-0050-02		Pack of 48	200.00

ABBREVIATIONS

Ac₂O = Acetic Anhydride
 CE = Cyanoethyl
 CPG = Controlled Pore Glass
 DCM = Dichloromethane
 dmf = dimethylformamide
 I₂ = Iodine
 MeIm = 1-Methylimidazole
 TCA = Trichloroacetic Acid
 THF = Tetrahydrofuran

SEE ALSO

Universal Supports p22
 Q-Supports p25
 High Load Supports p27

STERLING CE PHOSPHORAMIDITES

Glen Research CE (β-cyanoethyl) Phosphoramidites are produced and packaged to ensure the highest performance on DNA synthesizers. Every Glen Research product is accompanied by a Certificate of Analysis and HPLC trace, showing the results of our QC testing. Every Glen Research monomer vial is specially cleaned to eliminate particulate contamination.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Åkta oligopilot</i>			
dA-CE Phosphoramidite	10-1000-20	2.0g	100.00
	10-1000-50	5.0g	250.00
dC-CE Phosphoramidite	10-1010-20	2.0g	100.00
	10-1010-50	5.0g	250.00
Ac-dC-CE Phosphoramidite	10-1015-20	2.0g	100.00
	10-1015-50	5.0g	250.00
dG-CE Phosphoramidite	10-1020-20	2.0g	100.00
	10-1020-50	5.0g	250.00
dmf-dG-CE Phosphoramidite	10-1029-20	2.0g	100.00
	10-1029-50	5.0g	250.00
dT-CE Phosphoramidite	10-1030-20	2.0g	100.00
	10-1030-50	5.0g	250.00

QUALITY ASSURANCE

Every batch of these CE Phosphoramidites is tested as follows:

- 1. HPLC**
a) Identity is confirmed by comparison with a reference sample.
b) Purity is determined by HPLC to be ≥98.0%.
- 2. TLC**
Purity is verified by TLC.
- 3. ³¹P NMR**
Purity is determined by ³¹P NMR to be ≥98%.
- 4. Coupling Test**
Coupling efficiency is determined to be ≥99%.
- 5. Solution Test**
A 0.1M solution is determined to be clear and free of particulate contamination.
- 6. Loss on Drying**
Volatile contaminants are determined to be ≤2%.

SEE ALSO

HT DNA Phosphoramidites p13

STERLING SOLVENTS/REAGENTS

All solvents and reagents are prepared to our exacting specifications to ensure the highest synthesis efficiency and are passed through a 0.2 micron filter during packaging to eliminate particulate contamination.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Diluent</i>			
Acetonitrile, anhydrous	40-4050-45	60mL	12.00
	40-4050-50	100mL	16.00
<i>Äkta oligopilot</i>			
<i>Activator</i>			
0.40M Tetrazole in Acetonitrile	30-3105-71	1L	380.00
<i>Cap Mix A</i>			
Acetonitrile/Melm	40-4015-71	1L	145.00
<i>Cap Mix B*</i>			
Acetonitrile/Ac ₂ O/Lutidine	40-4028-71*	1L	160.00
<i>Oxidizing Solution</i>			
0.05M I ₂ in Pyridine/H ₂ O	40-4035-71	1L	225.00
<i>Deblocking Mix</i>			
3% DCA/DCM	40-4040-71	1L	80.00
3% TCA/DCM	40-4140-71	1L	80.00
3% DCA/toluene	40-4240-71	1L	145.00

ABBREVIATIONS

Ac₂O = Acetic Anhydride
 CE = Cyanoethyl
 CPG = Controlled Pore Glass
 DCA = Dichloroacetic Acid
 DCM = Dichloromethane
 I₂ = Iodine
 Melm = 1-Methylimidazole
 µm = micromole(s)

SEE ALSO

Alternative Activators p28

* Cap Mix B is a two part formulation that is combined immediately before shipment.

STERLING CE PHOSPHORAMIDITES

Dr. Oligo synthesizers belong to a family of synthesizers, including the parallel array synthesizers, Dr. Oligo 96, Dr. Oligo 192, Dr. Oligo 384 and Dr. Oligo 768, manufactured by Biolytic® Lab Performance, Inc. in Fremont, CA. Their web site can be found at: <http://www.biolytic.com>. Phosphoramidite monomers are packaged in 30mL and 240mL amber bottles for dissolving at a concentration of 1g/20mL and are connected directly to the instrument. Some instruments may also be configured to accept Applied Biosystems serum vials, as shown on page 6.

QUALITY ASSURANCE

Every batch of these CE Phosphoramidites is tested as follows:

- 1. HPLC**
a) Identity is confirmed by comparison with a reference sample.
b) Purity is determined by HPLC to be $\geq 98.0\%$.
- 2. TLC**
Purity is verified by TLC.
- 3. ^{31}P NMR**
Purity is determined by ^{31}P NMR to be $\geq 98\%$.
- 4. Coupling Test**
Coupling efficiency is determined to be $\geq 99\%$.
- 5. Solution Test**
A 0.1M solution is determined to be clear and free of particulate contamination.
- 6. Loss on Drying**
Volatile contaminants are determined to be $\leq 2\%$.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
dA-CE Phosphoramidite	10-1000-02M	0.25g	12.50
	10-1000-05M	0.5g	25.00
	10-1000-10M	1.0g	50.00
	10-1000-5S	5.0g	250.00
	10-1000-1S	10.0g	500.00
dC-CE Phosphoramidite	10-1010-02M	0.25g	12.50
	10-1010-05M	0.5g	25.00
	10-1010-10M	1.0g	50.00
	10-1010-5S	5.0g	250.00
	10-1010-1S	10.0g	500.00
Ac-dC-CE Phosphoramidite	10-1015-02M	0.25g	12.50
	10-1015-05M	0.5g	25.00
	10-1015-10M	1.0g	50.00
	10-1015-5S	5.0g	250.00
	10-1015-1S	10.0g	500.00
dG-CE Phosphoramidite	10-1020-02M	0.25g	12.50
	10-1020-05M	0.5g	25.00
	10-1020-10M	1.0g	50.00
	10-1020-5S	5.0g	250.00
	10-1020-1S	10.0g	500.00
dmf-dG-CE Phosphoramidite	10-1029-02M	0.25g	12.50
	10-1029-05M	0.5g	25.00
	10-1029-10M	1.0g	50.00
	10-1029-5S	5.0g	250.00
	10-1029-1S	10.0g	500.00
dT-CE Phosphoramidite	10-1030-02M	0.25g	12.50
	10-1030-05M	0.5g	25.00
	10-1030-10M	1.0g	50.00
	10-1030-5S	5.0g	250.00
	10-1030-1S	10.0g	500.00

STERLING SOLVENTS/REAGENTS

All solvents and reagents are prepared to our exacting specifications to ensure the highest synthesis efficiency and are passed through a 0.2 micron filter during packaging to eliminate particulate contamination. Parallel synthesizers typically use 5-ethylthio-1H-tetrazole (ETT) as activator to minimize the chance of crystallization. ETT is used at a concentration of 0.25M in acetonitrile, which is far below the level at which crystallization may occur.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Activator</i>			
0.25M ETT in Acetonitrile	30-3140-57	450mL	200.00
	30-3140-62	2L	760.00

SEE ALSO

Alternative Activators

p28

STERLING SOLVENTS/REAGENTS (CONT.)

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Diluent</i>			
Acetonitrile, anhydrous	40-4050-50	100mL	16.00
<i>Cap Mix A</i>			
THF/Lutidine/Ac ₂ O	40-4010-57	450mL	72.00
	40-4010-62	2L	325.00
<i>Cap Mix B</i>			
16% Melm in THF	40-4220-57	450mL	96.00
	40-4220-62	2L	425.00
<i>Oxidizing Solution</i>			
0.02M I ₂ in THF/Pyridine/H ₂ O	40-4330-57	450mL	72.00
	40-4330-62	2L	325.00
<i>Deblocking Mix</i>			
3% DCA/DCM	40-4040-57	450mL	36.00
	40-4040-62	2L	144.00
3% TCA/DCM	40-4140-57	450mL	36.00
	40-4140-62	2L	144.00

STERLING SUPPORTS

Dr. Oligo instruments are designed for flexibility in the use of supports and columns. They can use fritted plates with loose CPG (page 8) and AB 3900 style polystyrene and CPG columns. Glen UnySupport™ 40 nmole frits can also be used.

<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
dA	dC	dG	dT	Ac-dC	dmf-dG		
<i>AB 3900 Polystyrene Columns</i>							
26-2600-65	26-2610-65		26-2630-65		26-2629-65	200X40nm	825.00
26-2600-62	26-2610-62		26-2630-62		26-2629-62	200X200nm	825.00
<i>AB 3900 1000Å CPG Columns</i>							
20-2101-65			20-2131-65	20-2115-65	20-2129-65	200X40nm	600.00
20-2101-62			20-2131-62	20-2115-62	20-2129-62	200X200nm	650.00
20-2101-61			20-2131-61	20-2115-61	20-2129-61	200X1µm	875.00

OLIGONUCLEOTIDE PURIFICATION

Biolytic Labs. also offers the innovative Dr. Oligo Processor for high throughput purification of oligonucleotides using Glen-Pak™ DNA Purification Cartridges: <https://www.biolytic.com/p-6814-dr-oligo-processor-fully-automated.aspx>.

ABBREVIATIONS

Ac₂O = Acetic Anhydride
 CE = Cyanoethyl
 CPG = Controlled Pore Glass
 DCM = Dichloromethane
 dmf = dimethylformamide
 I₂ = Iodine
 Melm = 1-Methylimidazole
 TCA = Trichloroacetic Acid
 THF = Tetrahydrofuran

SEE ALSO


Universal Supports p22
 Q-Supports p25
 High Load Supports p27
 Glen-Pak™ DNA p138

ULTRAMILD CE PHOSPHoramidites


The synthesis of labelled oligonucleotides has become a standard procedure in many laboratories and many labelling reagents, e.g., biotin and fluorescein, are now available as β -cyanoethyl (CE) phosphoramidites. Labels which are currently available as CE phosphoramidites have one common property - they must be stable to the strongly alkaline conditions required for removal of the base protecting groups. This property is lacking in several interesting dyes and labels. We sought an alternative protecting scheme for the normal CE phosphoramidites which would allow UltraMILD deprotection and would not react with a wider variety of tags and labels. A set of monomers using phenoxyacetyl (Pac) protected dA and 4-isopropyl-phenoxyacetyl (iPr-Pac) protected dG, along with acetyl protected dC, met the desired criteria for UltraMILD deprotection.

We recommend the use of phenoxyacetic anhydride (Pac₂O) in Cap A. This modification removes the possibility of exchange of the iPr-Pac protecting group on the dG with acetate from the acetic anhydride capping mix. Cleavage and deprotection can be carried out in 2 hours at room temperature with ammonium hydroxide or 4 hours with 0.05M potassium carbonate in methanol.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Pac-dA-CE Phosphoramidite	10-1601-02	0.25g	15.00
	10-1601-05	0.5g	30.00
	10-1601-10	1.0g	60.00
Ac-dC-CE Phosphoramidite	10-1015-02	0.25g	12.50
	10-1015-05	0.5g	25.00
	10-1015-10	1.0g	50.00
iPr-Pac-dG-CE Phosphoramidite	10-1621-02	0.25g	15.00
	10-1621-05	0.5g	30.00
	10-1621-10	1.0g	60.00



Pac-dA



Ac-dC



iPr-Pac-dG

ULTRAMILD SUPPORTS

<i>Item</i>	<i>Catalog No.</i> Pac-dA	<i>Catalog No.</i> Ac-dC	<i>Catalog No.</i> iPr-Pac-dG	<i>Pack</i>	<i>Price(\$)</i>
UltraMild CPG (Bulk)	20-2601-01	Listed	20-2621-01	0.1g	18.00
	20-2601-02	on	20-2621-02	0.25g	40.00
	20-2601-10	Page 8	20-2621-10	1.0g	150.00
ABI Columns	20-2701-45	20-2115-45	20-2721-45	4X40nm	40.00
	20-2701-42	20-2115-42	20-2721-42	4X0.2µm	40.00
	20-2701-41	20-2115-41	20-2721-41	4X1µm	60.00
	20-2701-13	20-2115-13	20-2721-13	10µm	100.00
Expedite Columns	20-2801-45	20-2215-45	20-2821-45	4X40nm	40.00
	20-2801-42	20-2215-42	20-2821-42	4X0.2µm	40.00
	20-2801-41	20-2215-41	20-2821-41	4X1µm	60.00
	20-2801-14	20-2215-14	20-2821-14	15µm	150.00

ULTRAMILD SOLVENTS/REAGENTS

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Cap Mix A</i>			
THF/Pyridine/Pac ₂ O (Applied Biosystems)	40-4210-52	200mL	140.00
	40-4210-57	450mL	300.00
THF/Pac ₂ O (Expedite)	40-4212-52	200mL	140.00
	40-4212-57	450mL	300.00
<i>Deprotection Solution</i>			
0.05M Potassium Carbonate in Methanol	60-4600-30	30mL	30.00

SEE ALSO

Universal Support III p24

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers
For Instrument type Add

Expedite	E
MerMade	M

Columns
For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

REFERENCES


- (1) A.P. Guzaev, and M. Manoharan, *J Am Chem Soc*, 2003, **125**, 2380-2381.
- (2) R.K. Kumar, A.P. Guzaev, C. Rentel, and V.T. Ravikumar, *Tetrahedron*, 2006, **62**, 4528.

ELIMINATION CONDITIONS

Reagent	Conditions
Ammonium hydroxide	80°C/2h 55°C/8h
Ammonium hydroxide/ 40% Methylamine (AMA)	80°C/0.5h 65°C/1h 55°C/8h
Methylamine Gas	65°C/0.5h/30psi
Potassium Carbonate in Methanol	RT/17h
t-Butylamine/Water (1:3 v/v)	60°C/4h

INTELLECTUAL PROPERTY

This product is covered by US Patent 7,202,264 owned by Isis Pharmaceuticals, Inc..



Glen UnySupport

GLEN UNYSUPPORT

Our original Universal Support (20-5000) has been discontinued since complete dephosphorylation using ammonium hydroxide, AMA or anhydrous methylamine gas takes longer than most companies wish to allocate. A recent development has been the use of a support based on a molecule which is "conformationally preorganized" to accelerate the dephosphorylation reaction.^{1,2} By using a rigid bicyclic molecule on the support, the rate of elimination is markedly faster than the original Universal Support. The structure of Glen UnySupport™ is shown below. The N-phenyl version, developed at Isis Pharmaceuticals as UnyLinker™, is available from several companies for large scale oligo synthesis. Glen UnySupport is the N-methyl version, which is preferred for high throughput oligonucleotide synthesis since methylamine rather than aniline is formed on deprotection. We are happy to offer Glen UnySupport in a variety of popular formats under license from Isis Pharmaceuticals.

Item	Catalog No.	Pack	Price(\$)
Bulk Supports			
Glen UnySupport (500Å CPG)	20-5040-01	0.1g	11.00
	20-5040-02	0.25g	25.00
	20-5040-10	1.0g	95.00
Glen UnySupport (1000Å CPG)	20-5041-01	0.1g	11.00
	20-5041-02	0.25g	25.00
	20-5041-10	1.0g	95.00
High Load Glen UnySupport	25-5040-01	0.1g	15.00
	25-5040-02	0.25g	30.00
	25-5040-10	1.0g	115.00
Glen UnySupport PS	26-5040-01	0.1g	16.00
	26-5040-02	0.25g	35.00
	26-5040-10	1.0g	125.00
Columns			
The 1000Å columns and frits below are routinely stocked.			
ABI Format (not LV)			
1 µmole columns	20-5141-41	Pack of 4	60.00
0.2 µmole columns	20-5141-42	Pack of 4	40.00
40 nmole columns	20-5141-45	Pack of 4	40.00
10 µmole column (TWIST Format)	20-5141-13	Pack of 1	100.00
40 nmole frits	20-5441-95	Pack of 96	150.00
Female-Female Luer Adapter for 40 nmole frits	20-0060-00	Pack of 10	20.00
AB 3900 Format			
Glen UnySupport PS			
200 nmole columns	26-5140-52	Pack of 10	100.00
40 nmole columns	26-5140-55	Pack of 10	100.00
Expedite Format			
1 µmole columns	20-5241-41	Pack of 4	60.00
0.2 µmole columns	20-5241-42	Pack of 4	40.00
40 nmole columns	20-5241-45	Pack of 4	40.00
15 µmole column (TWIST Format)	20-5241-14	Pack of 1	150.00
96 Well Format (MerMade, etc.)			
1 µmole columns	20-5141-91	Pack of 96	375.00
200 nmole columns	20-5141-92	Pack of 96	250.00
40 nmole columns	20-5141-95	Pack of 96	250.00

GLEN UNYSUPPORT FC

The extended time required to cleave the succinate linkage of the original Glen UnySupport can be problematical, especially in high-throughput production of oligos, due to the outgassing of ammonia and/or methylamine. This reduction in concentration of gas can necessitate the evaporation of the cleavage solution and addition of fresh Ammonium Hydroxide:Methylamine 1:1 (AMA) or ammonium hydroxide (NH₄OH) to ensure complete deprotection and dephosphorylation of the product oligos. Using a diglycolate linkage in Glen UnySupport FC instead of the succinate in Glen UnySupport, a significant increase in the rate of cleavage has been achieved. The minimum cleavage times for both versions are as follows:

	AMA	NH ₄ OH
Glen UnySupport	10 min.	40 min.
Glen UnySupport FC	2 min.	5 min.

With the cleavage time of Glen UnySupport FC reduced to less than 5 minutes, there is minimal loss of volatile gas and, therefore, no need to evaporate the cleavage solution and replenish with fresh AMA or ammonium hydroxide solutions.

We offer Glen UnySupport FC attached to 1000Å CPG in a variety of formats suited to high throughput synthesis, as well as in bulk for more routine use.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
Bulk Support			
Glen UnySupport FC (1000Å CPG)	22-5041-01	0.1g	11.00
	22-5041-02	0.25g	25.00
	22-5041-10	1.0g	95.00
ABI Format (not LV)			
1 µmole columns	22-5141-41	Pack of 4	60.00
0.2 µmole columns	22-5141-42	Pack of 4	40.00
40 nmole columns	22-5141-45	Pack of 4	40.00
10 µmole column (TWIST Format)	22-5141-13	Pack of 1	100.00
AB 3900 Format			
Glen UnySupport CPG			
200 nmole columns	22-5141-52	Pack of 10	100.00
40 nmole columns	22-5141-55	Pack of 10	100.00
Expedite Format			
1 µmole columns	22-5241-41	Pack of 4	60.00
0.2 µmole columns	22-5241-42	Pack of 4	40.00
40 nmole columns	22-5241-45	Pack of 4	40.00
15 µmole column (TWIST Format)	22-5241-14	Pack of 1	150.00
96 Well Format (MerMade, etc.)			
1 µmole columns	22-5141-91	Pack of 96	375.00
200 nmole columns	22-5141-92	Pack of 96	250.00
40 nmole columns	22-5141-95	Pack of 96	250.00

ELIMINATION CONDITIONS

Reagent	Conditions
Ammonium hydroxide	80°C/2h 55°C/8h
Ammonium hydroxide/ 40% Methylamine (AMA)	80°C/0.5h 65°C/1h 55°C/8h
Methylamine Gas	65°C/0.5h/30psi
Potassium Carbonate in Methanol	RT/17h
t-Butylamine/Water (1:3 v/v)	60°C/4h

INTELLECTUAL PROPERTY

This product is covered by US Patent 7,202,264 owned by Isis Pharmaceuticals, Inc..



Glen UnySupport FC

REFERENCES

- (1) A.V. Azhayev, *Tetrahedron*, 1999, **55**, 787-800.
 (2) A.V. Azhayev and M. Antopolsky, *Tetrahedron*, 2001, **57**, 4977-4986.

INTELLECTUAL PROPERTY

This product is covered by
 US Patent No.: 6,770,754 and
 European Patent No.: 1404695.

CLEAVAGE AND DEPROTECTION

1. Cleavage

For standard and UltraFast deprotection protocols, cleave the oligo from the support using 2M ammonia in methanol at room temperature for 30 minutes. (Only for oligonucleotides greater than 50 nucleotides in length, rinse the support with a further volume of water. Combine the two washes and evaporate to dryness.)

2. Deprotection

Standard

Add 1 volume of 30% ammonium hydroxide, seal and deprotect using the conditions appropriate for removal of the protecting groups on the nucleobases.

UltraFast

Add 1 volume of AMA (ammonium hydroxide/40% aqueous methylamine 1:1) seal and deprotect at 65°C for 10 minutes.


UltraMild Using Ammonium Hydroxide

Add 1 volume of ammonium hydroxide, seal and leave at room temperature for 8 hours.


UltraMild Cleavage and Deprotection

Using Potassium Carbonate in Methanol

Cleave the oligo from the support using 50 mM potassium carbonate in methanol at room temperature for 30 minutes. Seal and leave overnight at room temperature.



Universal Support II



Universal Support III

UNIVERSAL SUPPORT III

The key step in the use of any universal support in oligonucleotide synthesis is the dephosphorylation of the 3'-phosphate group to form the desired 3'-hydroxyl group. Azhayev^{1,2} has excelled in the investigation of neighboring group assistance in the dephosphorylation reaction. Amide groups may be considered to be weak N-H acids and can display basic properties in ammonium hydroxide or aqueous methylamine. (±)-3-Amino-1,2-propanediol was used to form a novel universal support. In our original US II support, a succinate linker attaches the 3-amino group to the support and the 2-OH is protected with a base-labile group to set up an amide assisted elimination in mildly basic conditions. In this way, the dephosphorylation reaction would eliminate the desired 3'-OH oligonucleotide into solution and the product of any β-elimination competing side reaction would remain bound to the support. A further improvement has been achieved by using a carbamate group to connect the universal linker to the support, now called Universal Support III. The structures of the two supports are shown below right. Because the universal linker is unchanged and the succinate or carbamate groups remain attached to the support, we use the same catalog numbers for US II and III. Using Universal Support II or III, an oligo yield of > 80% can be achieved on CPG supports and > 95% on polymeric supports, with purity equivalent to the same oligo prepared normally.

Conditions for Cleavage and Deprotection are outlined in the table opposite. Universal Support II/III has been shown to generate oligonucleotides with the same efficacy in polymerase extension reactions as regular oligos. Despite the mild elimination reaction, oligonucleotides up to 75mer in length can be prepared routinely without loss of oligo during the synthesis cycles. This support is also used for the production of siRNA oligos.

Item	Catalog No.	Pack	Price(\$)
Bulk Support			
Universal Support III (1000Å CPG)	20-5010-01	0.1g	11.00
	20-5010-02	0.25g	25.00
	20-5010-10	1.0g	95.00
Universal Support III PS	26-5010-01	0.1g	16.00
	26-5010-02	0.25g	35.00
	26-5010-10	1.0g	125.00
ABI Format (not LV)			
Universal Support III CPG 1 μmole columns	20-5110-41	Pack of 4	60.00
	20-5110-42	Pack of 4	40.00
	20-5110-45	Pack of 4	40.00
40 nmole columns			
	20-5110-13	Pack of 1	100.00
Expedite Format			
1 μmole columns	20-5210-41	Pack of 4	60.00
	20-5210-42	Pack of 4	40.00
	20-5210-45	Pack of 4	40.00
40 nmole columns			
	20-5210-14	Pack of 1	150.00
96 Well Format (MerMade, etc.)			
Universal Support III CPG 1 μmole columns	20-5110-91	Pack of 96	375.00
	20-5110-92	Pack of 96	250.00
	20-5110-95	Pack of 96	250.00
200 nmole columns			
40 nmole columns			
Universal Support III PS			
1 μmole columns	26-5110-91	Pack of 96	375.00
	26-5110-92	Pack of 96	250.00
	26-5110-95	Pack of 96	250.00
200 nmole columns			
40 nmole columns			
AB 3900 Format			
Universal Support III PS 200 nmole columns	26-5110-52	Pack of 10	100.00
	26-5110-55	Pack of 10	100.00

UNIVERSAL HybridCPG™ SOLID SUPPORTS

HybridCPG™ consists of CPG particles conformally coated with a very thin crosslinked polymer film based on polystyrene. The molecular structure of the polymer coating is designed to have a very high density of evenly distributed attachment points for optimum oligo synthesis. In this way, the pore size to loading density trade-off is minimized and the chemical resistance to glass-aggressive reagents is greatly improved. Although the nano-scale coating is subject to swelling, the rigid pore structure of the CPG substrate accommodates this and, at the same time, maintains a uniform pore space for the oligo synthesis. Thus, HybridCPG exhibits no bulk swelling in the synthesis solvents and allows much higher ligand loadings for a given pore size compared to conventional CPG. Glen Research is partnering with Prime Synthesis, developers of HybridCPG, to introduce our two popular Universal Supports, Glen UnySupport™ and Universal Support III, on HybridCPG.

Item	Catalog No.	Pack	Price(\$)
<i>Bulk Support</i>			
US III HybridCPG	28-5010-01	0.1g	16.00
	28-5010-02	0.25g	35.00
	28-5010-10	1.0g	125.00
Glen UnySupport HybridCPG	28-5040-01	0.1g	16.00
	28-5040-02	0.25g	35.00
	28-5040-10	1.0g	125.00

Q-SUPPORTS

Oligonucleotides are routinely prepared on supports to which the first nucleoside is attached via a succinate linkage. Over the years, the succinate linkage has demonstrated stability during the synthesis process but has sufficient lability to be cleaved quickly in the deprotection step. However, if the cleavage step is carried out with ammonium hydroxide manually or on the synthesizer, it consumes one hour of precious time while releasing only about 80% of the oligonucleotide. This step is, therefore, a bottleneck in the productivity of many synthesis groups.

Is it possible to find a replacement to the succinate group which offers good stability to the synthesis reagents while offering a much faster cleavage step? The oxalate group has been shown to be very labile during cleavage but its stability to the normal synthesis reagents is not good, requiring changes for successful use. In a practical but elegant study¹ of various bifunctional carboxylic acids, Richard Pon's group identified hydroquinone-*O,O'*-diacetic acid as the most satisfactory alternative to the succinate group. Nucleosides with this linker arm (Q-linker) are attached to supports with the same ease as the succinyl linker arm.

The cleavage time in ammonium hydroxide at room temperature was found to be 2 minutes, but what about the stability during synthesis? How significant was premature cleavage of oligonucleotide on the synthesizer because of the basic reagents in the capping mixes and oxidizer? Pon showed that the Q-linker is stable to the capping reagents but very slightly labile to the oxidizer (8% cleavage in overnight exposure which would correspond to about 2,000 normal synthesis cycles).

We tested the significance of premature cleavage by preparing sixteen 20mer oligonucleotides on a 0.2 μmole scale, eight with succinate and eight with Q-linkers. The succinate supported oligos were cleaved for 1 hour at room temperature, while those on the Q-support were cleaved for 2 minutes. Both sets were then deprotected normally with ammonium hydroxide. The Q-supports actually gave 5% better yields of product than the succinate supports. Oligo purities were equivalent in both sets.

The Q-linker is absolutely compatible with all hydrolytic cleavage procedures, but especially mild procedures like potassium carbonate in methanol. Pon also showed that it is preferable for RNA supports, improving the cleavage time for 2'-silyl protected nucleoside supports from 2 hours (60-65% cleavage) to 5 minutes (95% cleavage).

We are offering Q-linkers of the four regular nucleosides on 500Å CPG in 0.2 and 1 μmole scales.

TRADEMARKS

HybridCPG™ is a trademark of Prime Synthesis, Inc. Glen UnySupport™ is a trademark of Glen Research Corporation.

REFERENCE

- (1) R.T. Pon and S.Y. Yu, *Tetrahedron Lett*, 1997, **38**, 3327-3330.

INTELLECTUAL PROPERTY

Q-Supports are supplied under license from University Technologies International LP (UTI LP).

Q/SUCCINATE COMPARISON

Q-Support (2 minutes cleavage)	Succinate (60 minutes cleavage)
132 ODU*	125 ODU*

*Average crude yield from eight 1 μmole columns deprotected normally.

Q-SUPPORTS (CONT.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M






Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)


Catalog No.	Catalog No.	Catalog No.	Catalog No.	Catalog No.	Pack	Price(\$)
dA	dC	Ac-dC	dmf-dG	dT		
500Å Bulk Support						
21-2000-01	21-2010-01	21-2013-01	21-2029-01	21-2030-01	0.1g	11.00
21-2000-02	21-2010-02	21-2013-02	21-2029-02	21-2030-02	0.25g	25.00
21-2000-10	21-2010-10	21-2013-10	21-2029-10	21-2030-10	1.0g	95.00
ABI Format (not LV)						
21-2100-41	21-2110-41	21-2113-41	21-2129-41	21-2130-41	4X1µm	60.00
21-2100-42	21-2110-42	21-2113-42	21-2129-42	21-2130-42	4X0.2µm	40.00
Expedite Format						
21-2200-41	21-2210-41	21-2213-41	21-2229-41	21-2230-41	4X1µm	60.00
21-2200-42	21-2210-42	21-2213-42	21-2229-42	21-2230-42	4X0.2µm	40.00




HIGH LOAD CPG

Our high loading support is based on controlled pore silica and it retains the usual 500Å pores. The spacer is also conventional. The only significant difference is the loading which is in the range 80 - 130µmoles/g or about 2.5 times the loading of normal 500Å CPG. Typical loadings for our high load CPG are in the 100 - 120µmoles/g range. As a consequence of the high loading, this support should not be used for sequences longer than 40mers. This high loading support is available in columns for most synthesizers. The 2.5µmole column is identical to our standard 1µmole column (with the exception of the loading). It should be used on occasions when greater than 1µmole is desired but when a 10 or 15µmole synthesis is too high. It should be run using the 1µmole cycle. The 25µmole column is identical to the 10µmole column used on Applied Biosystems synthesizers. It is run using the 10µmole cycle. The 35µmole column is used as an alternative to the 15µmole Expedite column. Again no changes to the standard cycle are recommended. The support is of course available in bulk for use on large-scale synthesizers. A word of caution is in order. When using a column with a higher load than recommended by the instrument manufacturer, there is a much smaller margin for error. All reagents must be fresh and anhydrous diluent and activator must be used. Should you decide to prepare higher-loading columns, ensure that the molar excess of monomer to support nucleoside is at least 5X and preferably 10X.


Item	Catalog No. dA	Catalog No. dC	Catalog No. dG	Catalog No. dT	Pack	Price(\$)
Columns						
(ABI)	25-2100-46	25-2110-46	25-2120-46	25-2130-46	4X2.5µm	75.00
	25-2100-17	25-2110-17	25-2120-17	25-2130-17	1X25µm	125.00
(Expedite)	25-2200-46	25-2210-46	25-2220-46	25-2230-46	4X2.5µm	75.00
	25-2200-18	25-2210-18	25-2220-18	25-2230-18	1X35µm	185.00
Bulk						
	25-2000-02	25-2010-02	25-2020-02	25-2030-02	0.25g	25.00
	25-2000-10	25-2010-10	25-2020-10	25-2030-10	1.0g	90.00




dA-CPG



dC-CPG



dG-CPG



dT-CPG

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M

Columns

For Instrument type Add

Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)

SEE ALSO


Glen UnySupport

p22


ALTERNATIVE SOLVENTS/REAGENTS

ABBREVIATIONS


Ac₂O = Acetic Anhydride
 DCA = Dichloroacetic Acid
 DCM = Dichloromethane
 DMAP = Dimethylaminopyridine
 I₂ = Iodine
 MeIm = 1-Methylimidazole
 TCA = Trichloroacetic Acid
 THF = Tetrahydrofuran




5-Ethylthio-1H-tetrazole



DCI



5-Benzylthio-1H-tetrazole



Saccharin 1-Methylimidazole

INTELLECTUAL PROPERTY

SMI is sold under license from Avecia Biotechnology Inc.

Glen Research offers alternative solvents and reagents in suitable bottles and formulations for use on various DNA synthesizers. All solvents and reagents are prepared to our exacting specifications to ensure the highest coupling efficiencies and are passed through a 0.2 micron filter during packaging to eliminate particulate contamination. Glen Research offers the activators below in powder form for later dissolution in anhydrous acetonitrile or as a prepared solution.

Item	Catalog No.	Pack	Price (\$)
Activator			
5-Ethylthio-1H-tetrazole (ETT) <i>(Dissolve 1g in 31mL anhydrous acetonitrile for a 0.25M solution)</i>	30-3040-10	1g	35.00
	30-3040-20	2g	60.00
	30-3040-25	25g	500.00
0.25M 5-Ethylthio-1H-tetrazole in Acetonitrile <i>(Applied Biosystems)</i>	30-3140-45	45mL	40.00
	30-3140-52	200mL	100.00
	30-3140-57	450mL	200.00
	30-3140-62	2L	760.00
	30-3142-52	200mL	100.00
	30-3140-57	450mL	200.00
<i>(Expedite)</i>			
4,5-Dicyanoimidazole (DCI), crystalline <i>(Dissolve 1g in 34mL anhydrous acetonitrile for a 0.25M solution)</i>	30-3050-10	1g	35.00
	30-3050-25	25g	500.00
0.25M DCI in Acetonitrile <i>(Applied Biosystems)</i>	30-3150-45	45mL	40.00
	30-3150-52	200mL	100.00
	30-3150-57	450mL	200.00
	30-3150-62	2L	760.00
	30-3152-52	200mL	100.00
	30-3150-57	450mL	200.00
<i>(Expedite)</i>			
5-Benzylthio-1H-tetrazole (BTT) <i>(Dissolve 1g in 21.3mL anhydrous acetonitrile for a 0.25M solution)</i>	30-3070-10	1g	35.00
	30-3070-20	2g	60.00
	30-3070-25	25g	500.00
0.25M 5-Benzylthio-1H-tetrazole in Acetonitrile <i>(Applied Biosystems)</i>	30-3170-45	45mL	40.00
	30-3170-52	200mL	100.00
	30-3170-57	450mL	200.00
	30-3170-62	2L	760.00
	30-3172-52	200mL	100.00
	30-3170-57	450mL	200.00
<i>(Expedite)</i>			
Saccharin 1-Methylimidazole (SMI) <i>(Dissolve 1g in 31mL anhydrous acetonitrile for a 0.2M solution)</i>	30-3080-10	1g	35.00
	30-3080-20	2g	60.00
	30-3080-25	25g	500.00
0.2M Saccharin 1-Methylimidazole (SMI) in Acetonitrile <i>(Applied Biosystems)</i>	30-3180-45	45mL	40.00
	30-3180-52	200mL	100.00
	30-3180-57	450mL	200.00
	30-3180-62	2L	760.00
	30-3182-52	200mL	100.00
	30-3180-57	450mL	200.00
<i>(Expedite)</i>			


ALTERNATIVE SOLVENTS/REAGENTS (CONT.)

Item	Catalog No.	Pack	Price (\$)
<i>Cap Mix A</i>			
THF/Lutidine/Ac ₂ O	40-4010-52	200mL	30.00
	40-4010-57	450mL	72.00
	40-4010-62	2L	325.00
THF/Ac ₂ O (9:1)	40-4012-62	2L	275.00
<i>Cap Mix B</i>			
6.5% DMAP in THF (Cap B solutions containing DMAP are preferred by some researchers for preparing long oligos.)	40-4020-52	200mL	42.00
10% Melm in THF	40-4120-52	200mL	30.00
	40-4120-57	450mL	72.00
	40-4120-62	2L	325.00
10% Melm in THF/Pyridine (8:1)	40-4122-62	2L	325.00
<i>Oxidizing Solution</i>			
0.02M I ₂ in THF/Pyridine/H ₂ O	40-4132-62	2L	325.00
0.5M CSO in Anhydrous Acetonitrile (ABI)	40-4632-52	200mL	250.00
0.5M CSO in Anhydrous Acetonitrile (Expedite) (CSO is an alternative anhydrous Oxidizer. A minimum oxidation time of 3 minutes is required on small scales.)	40-4632-52E	200mL	250.00
<i>Deblocking Mix</i>			
3% DCA/DCM (DCA solutions are more mildly acidic than the TCA equivalents, possibly causing less depurination of dA sites.)	40-4040-57	450mL	36.00
	40-4040-62	2L	144.00
2.5% DCA/DCM	40-4042-57	450mL	36.00
	40-4042-62	2L	144.00
DCA (Add to 4L of DCM to make a 3% solution.)	40-4044-54	120mL	60.00

Alternative Capping Reagent

The phosphoramidite of diethylene glycol monoethyl ether, UniCap, is the basis for an alternative capping reagent. To use UniCap as a capping amidite on the Expedite 8909 or AB synthesizers, dilute it to the standard amidite concentration and place the vial in position 5 on the instrument. Cycles can be modified by adding coupling steps for amidite reservoir 5 after the last column coupling step. The standard capping steps can be left out of the cycle. UniCap Phosphoramidite was originally developed for oligo synthesis on the surface of chips and is the capping reagent of choice for this application.

Item	Catalog No.	Pack	Price (\$)
UniCap Phosphoramidite	10-4410-02	0.25g	50.00
	10-4410-05	0.5g	100.00
	10-4410-10	1.0g	200.00
	10-4410-20	2.0g	400.00




CSO

SEE ALSO

0.1M CSO in PACE Chemistry p33

INTELLECTUAL PROPERTY

This capping reagent is supplied under license.



UniCap Phosphoramidite

5'-CE PHOSPHORAMIDITES

Glen Research 5'-CE (β-cyanoethyl) Phosphoramidites are designed for the production of 5'-5' or 3'-3' linkages, useful in antisense studies, or to synthesize oligonucleotide segments in the opposite sense from normal synthesis, for structural studies. These monomers are packaged in ABI-style vials (see note box).

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
dA-5'-CE Phosphoramidite	10-0001-02	0.25g	75.00
	10-0001-05	0.5g	150.00
	10-0001-10	1.0g	300.00
dC-5'-CE Phosphoramidite	10-0101-02	0.25g	75.00
	10-0101-05	0.5g	150.00
	10-0101-10	1.0g	300.00
dmf-dG-5'-CE Phosphoramidite	10-9201-02	0.25g	75.00
	10-9201-05	0.5g	150.00
	10-9201-10	1.0g	300.00
dT-5'-CE Phosphoramidite	10-0301-02	0.25g	75.00
	10-0301-05	0.5g	150.00
	10-0301-10	1.0g	300.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add


Expedite E
MerMade M

Columns


For Instrument type Add

Expedite E
Applied Biosystems 3900 A
MerMade M


(Please inquire for availability of vials and columns for other instrument types.)




dA-5'-CE Phosphoramidite



dC-5'-CE Phosphoramidite



Dmf-dG-5'-CE Phosphoramidite







dT-5'-CE Phosphoramidite

5'-SUPPORTS

The following supports are used to produce oligonucleotides with nuclease resistant 3'-3' linkages at the 3' terminus (by attaching regular 3'-CE phosphoramidites) or to produce oligonucleotide sections in the opposite sense (by attaching 5'-CE phosphoramidites). ABI-style columns are supplied unless otherwise requested (see note box).

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
dA-5'-CPG	20-0002-01	0.1g	50.00
	20-0002-10	1.0g	375.00
1 μmole columns	20-0012-41	Pack of 4	100.00
0.2 μmole columns	20-0012-42	Pack of 4	75.00
10 μmole column (ABI)	20-0012-13	Pack of 1	225.00
15 μmole column (Expedite)	20-0012-14	Pack of 1	300.00
dC-5'-CPG	20-0102-01	0.1g	50.00
	20-0102-10	1.0g	375.00
1 μmole columns	20-0112-41	Pack of 4	100.00
0.2 μmole columns	20-0112-42	Pack of 4	75.00
10 μmole column (ABI)	20-0112-13	Pack of 1	225.00
15 μmole column (Expedite)	20-0112-14	Pack of 1	300.00
dG-5'-CPG	20-0202-01	0.1g	50.00
	20-0202-10	1.0g	375.00
1 μmole columns	20-0212-41	Pack of 4	100.00
0.2 μmole columns	20-0212-42	Pack of 4	75.00
10 μmole column (ABI)	20-0212-13	Pack of 1	225.00
15 μmole column (Expedite)	20-0212-14	Pack of 1	300.00
dT-5'-CPG	20-0302-01	0.1g	50.00
	20-0302-10	1.0g	375.00
1 μmole columns	20-0312-41	Pack of 4	100.00
0.2 μmole columns	20-0312-42	Pack of 4	75.00
10 μmole column (ABI)	20-0312-13	Pack of 1	225.00
15 μmole column (Expedite)	20-0312-14	Pack of 1	300.00



METHYL PHOSPHONAMIDITES

REFERENCE

- (1) M.P. Reddy, F. Farooqui, and N.B. Hanna, *Tetrahedron Lett.*, 1996, **37**, 8691-8694.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M

Columns


For Instrument type Add

Expedite E
Applied Biosystems 3900 A
MerMade M


(Please inquire for availability of vials and columns for other instrument types.)

Methyl Phosphonamidites may be used in DNA synthesizers following conventional CE Phosphoramidite protocols to produce oligonucleotides containing one or more methyl phosphonate linkages. However, deprotection and purification techniques differ and a description of the procedures is included in the Technical Bulletin. We also offer the dC monomer with acetyl base protection.¹ This protecting group is removed with ammonium hydroxide during the cleavage step, eliminating modification at the dC sites during the deprotection step using ethylenediamine in ethanol.


Item	Catalog No.	Pack	Price (\$)
dA-Me Phosphonamidite	10-1100-02	0.25g	50.00
	10-1100-05	0.5g	100.00
Ac-dC-Me Phosphonamidite	10-1115-02	0.25g	50.00
	10-1115-05	0.5g	100.00
dG-Me Phosphonamidite	10-1120-02	0.25g	50.00
	10-1120-05	0.5g	100.00
dT-Me Phosphonamidite	10-1130-02	0.25g	50.00
	10-1130-05	0.5g	100.00




dA-Me Phosphonamidite



Ac-dC-Me Phosphonamidite



dG-Me Phosphonamidite



dT-Me Phosphonamidite


PACE PHOSPHORAMIDITES

Phosphonoacetate (PACE) modified oligonucleotides show great potential as biological modifiers in a wide variety of research applications. PACE monomers are part of a family of Phosphonocarboxylate monomers. The monomers can be easily incorporated into complex oligonucleotides and are compatible with a wide variety of other sugar or heterobase modifications. PACE DNA can be conjugated through the carboxylic acid functional group. They have been shown to be active in siRNA duplexes and accelerate the initial rate of cleavage by RNase H-1 when incorporated with phosphorothioates. However, the most interesting observation to date is that they exhibit an unprecedented enhancement in penetration of cultured cells.


The phosphonoacetates are fully soluble in acetonitrile at a recommended concentration of 0.1M and are compatible with standard DNA synthesizers. A recommended coupling time of 33.3 minutes with 1H-Tetrazole is necessary when using the standard protocol. A modified LV cycle for AB instruments that reduces coupling time to 15 minutes with 1H-Tetrazole is available on our website. Oxidation must precede capping in the synthesis cycle. Reagents for oxidation depend on the type of synthesis. For fully modified oligos, we recommend the non-aqueous oxidizer camphorsulfonyloxaziridine (CSO) as a 0.1M solution. For mixed phosphodiester and phosphonoacetate modified oligos, a 0.5M CSO solution is recommended. Low water oxidizer, 40-4032, is an alternative oxidizing reagent although it has been reported that this can result in conversion of a small percentage of the phosphonoacetate to the phosphodiester. We also recommend the use of the Cap Mix B with DMAP (40-4020) instead of the standard Cap Mix B containing 1-Methylimidazole.

The standard protocol for cleavage and deprotection requires a two step method with pretreatment using 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU) and subsequent cleavage using methylamine. The DBU is used to deprotect the dimethylcyanoethyl (DMCE) protecting groups and to prevent alkylation of the bases during deprotection. Cleavage with 40% methylamine in water is recommended and we have also had good results when using AMA deprotection.


Item	Catalog No.	Pack	Price(\$)
dA-PACE Phosphoramidite	10-1140-02	0.25g	100.00
	10-1140-05	0.5g	200.00
	10-1140-10	1.0g	400.00
Ac-dC-PACE Phosphoramidite	10-1150-02	0.25g	100.00
	10-1150-05	0.5g	200.00
	10-1150-10	1.0g	400.00
dG-PACE Phosphoramidite	10-1160-02	0.25g	100.00
	10-1160-05	0.5g	200.00
	10-1160-10	1.0g	400.00
dT-PACE Phosphoramidite	10-1170-02	0.25g	100.00
	10-1170-05	0.5g	200.00
	10-1170-10	1.0g	400.00
0.1M CSO in Anhydrous Acetonitrile (ABI)	40-4631-52	200mL	85.00
0.1M CSO in Anhydrous Acetonitrile (Expedite)	40-4631-52E	200mL	85.00




dA-PACE Phosphoramidite



Ac-dC-PACE Phosphoramidite



dG-PACE Phosphoramidite



dT-PACE Phosphoramidite

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers For Instrument type Add

Expedite	E
MerMade	M

Columns For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

INTELLECTUAL PROPERTY

These products are covered by patents, US 6,693,187 and 7,067,641, and patents pending owned by Metasense Technologies. Purchase of all or any of these products includes a limited license to use the products solely for the manufacture of oligonucleotides for research use only. This license specifically excludes the use of the product or oligonucleotides containing the product for: (a) therapeutic or diagnostic applications (including kits, pools, libraries and other products or services that incorporate oligonucleotides containing the product), (b) any in vivo toxicity/safety study in support of an investigational new drug application (or foreign counterpart), or (c) resale (including sale of kits, pools, libraries and other products or services that incorporate the product or oligonucleotides containing the product). If such activities have commercial application, a separate license is required from Metasense Technologies. Neither the product nor any product created through its use may be used in human clinical trials.

A simple agreement must be signed before end-users and custom oligo services may purchase these products for use as defined above. <http://www.glenresearch.com/Reference/PACE.pdf>

SEE ALSO

0.5M CSO

p29


METHYL PHOSPHORAMIDITES

For many years, Glen Research has supplied methyl phosphoramidites in addition to β -cyanoethyl (CE) phosphoramidites for the few situations where the more labile cyanoethyl group is not an advantage. Some of our customers, probably remembering that the methyl group was removed specifically with thiophenol, have tried to use these monomers to prepare the interesting, uncharged, and nuclease-resistant methyl phosphotriester linkage. Unfortunately, this linkage is labile to ammonium hydroxide and the regular phosphodiester linkage is formed (along with a small amount of chain scission). We offer UltraMild methyl phosphoramidites for this application. Oligos produced from these monomers can be deprotected with potassium carbonate in methanol to produce methyl phosphotriester linkages. Since these linkages are diastereomeric and uncharged, the oligos may be hard to handle. Consequently, it is likely that chimeras will be produced using these monomers along with the regular UltraMild CE phosphoramidites. If many dG residues are included in the oligonucleotide, we recommend the use of phenoxyacetic anhydride (Pac₂O) in Cap A. This modification removes the possibility of exchange of the isopropyl-phenoxyacetate (iPr-Pac) protecting group on the dG with acetate from the acetic anhydride capping mix.


Item	Catalog No.	Pack	Price(\$)
Pac-dA-Me Phosphoramidite	10-1301-02	0.25g	25.00
	10-1301-05	0.5g	50.00
	10-1301-10	1.0g	100.00
Ac-dC-Me Phosphoramidite	10-1315-02	0.25g	25.00
	10-1315-05	0.5g	50.00
	10-1315-10	1.0g	100.00
iPr-Pac-dG-Me Phosphoramidite	10-1321-02	0.25g	25.00
	10-1321-05	0.5g	50.00
	10-1321-10	1.0g	100.00
dT-Me Phosphoramidite	10-1330-02	0.25g	25.00
	10-1330-05	0.5g	50.00
	10-1330-10	1.0g	100.00

ULTRAMILD SOLVENTS/REAGENTS


Item	Catalog No.	Pack	Price (\$)
<i>Cap Mix A</i> THF/Pyridine/Pac ₂ O (Applied Biosystems)	40-4210-52	200mL	140.00
	40-4210-57	450mL	300.00
THF/Pac ₂ O (Expedite)	40-4212-52	200mL	140.00
	40-4212-57	450mL	300.00
<i>Deprotection Solution</i> 0.05M Potassium Carbonate in Methanol	60-4600-30	30mL	30.00




Pac-dA-Me Phosphoramidite



Ac-dC-Me Phosphoramidite



iPr-Pac-dG-Me Phosphoramidite



dT-Me Phosphoramidite

H-PHOSPHONATE MONOMERS


Glen Research H-Phosphonates are analyzed by HPLC and are synthesis-tested. H-Phosphonates are especially useful for the preparation of modified internucleotide linkages which are unattainable by phosphoramidite chemistry. The most popular application is the preparation of radiolabelled phosphorothioates, since the sulfurization reaction is carried out off the synthesizer. These monomers are packaged in ABI-style vials (see note box).

Item	Catalog No.	Pack	Price(\$)
dA-H-Phosphonate, TEA Salt	10-1200-02	0.25g	40.00
	10-1200-05	0.5g	80.00
dC-H-Phosphonate, DBU Salt	10-1210-02	0.25g	40.00
	10-1210-05	0.5g	80.00
dG-H-Phosphonate, TEA Salt	10-1220-02	0.25g	40.00
	10-1220-05	0.5g	80.00
dT-H-Phosphonate, TEA Salt	10-1230-02	0.25g	40.00
	10-1230-05	0.5g	80.00


H-PHOSPHONATE REAGENTS

Our H-Phosphonate solvents and reagents have been discontinued. H-Phosphonate reagents are easily prepared using high purity products and the formulations shown below.


Item	Catalog No.	Pack	Price (\$)
1-Adamantanecarbonyl chloride is available from Aldrich, Catalog No. 117722. Dilute to 0.1M. <i>(Activator for monomers and capping reagent)</i>			
Acetonitrile/Pyridine (50:50), anhydrous <i>(Monomer Diluent)</i>			
Acetonitrile/Pyridine (95:5), anhydrous <i>(Activator Diluent)</i>			
1% Isopropyl Phosphite in Acetonitrile/Pyridine (50:50) <i>(Capping Reagent)</i>			
Acetonitrile/Pyridine (50:50) <i>(Neutralizer and Wash Solvent)</i>			
4% I ₂ in Pyridine/H ₂ O/THF (10:10:80) THF/H ₂ O/TEA (80:10:10) <i>(Both reagents are required for oxidation of H-phosphonate linkages)</i>			




dA-H-Phosphonate



dC-H-Phosphonate



dG-H-Phosphonate



dT-H-Phosphonate

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

 Monomers
 For Instrument type Add

Expedite	E
MerMade	M

 Columns
 For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

ABBREVIATIONS

I₂ = Iodine
 TEA = Triethylamine
 THF = Tetrahydrofuran

THIOPHOSPHORAMIDITES

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

REFERENCES

- (1) J. Nielsen, W.K.D. Brill, and M.H. Caruthers, *Tetrahedron Letters*, 1988, **29**, 2911-2914.
- (2) L. Cummins, D. Graff, G. Beaton, W.S. Marshall, and M.H. Caruthers, *Biochemistry*, 1996, **35**, 8734-41.
- (3) X. Yang, and D.G. Gorenstein, *Curr Drug Targets*, 2004, **5**, 705-15.
- (4) W.S. Marshall, and M.H. Caruthers, *Science*, 1993, **259**, 1564-70.
- (5) J.L. Tonkinson, et al., *Antisense Research and Development*, 1994, **4**, 269-278.
- (6) X. Yang, et al., *Bioorg Med Chem Lett*, 1999, **9**, 3357-62.
- (7) X. Yang, et al., *Ann N Y Acad Sci*, 2006, **1082**, 116-9.
- (8) X. Yang, et al., *Nucleic Acids Res*, 2002, **30**, e132.





Replacing two non-bridging oxygen atoms with sulfur atoms in a DNA phosphodiester linkage creates a phosphorodithioate (PS2) linkage.¹ Like natural DNA, the phosphorodithioate linkage is achiral at phosphorus. This analog is completely resistant to nuclease degradation and forms complexes with DNA and RNA with somewhat reduced stabilities.² Moreover, it has been found that PS2-ODNs bind proteins with a higher affinity than their phosphodiester analogues²⁻⁶ suggesting that PS2-ODNs may have additional utility in the form of sulfur-modified phosphate ester aptamers (thioaptamers)^{3,6-8} for therapeutic and diagnostic applications. Thiophosphoramidites are now commercially available after recent work at AM Biotechnologies (www.thioaptamer.com).

- 1) Thiophosphoramidites (ThioPAs) are not soluble in anhydrous acetonitrile diluent. Rather, 10% DCM (v/v) in acetonitrile is an ideal diluent for all four of the thioPAs for a final amidite concentration of 0.15 M.
- 2) ThioPAs are somewhat less stable than normal DNA phosphoramidites in anhydrous acetonitrile containing 10% DCM; however, the coupling efficiency of all four thioPAs is not reduced after two days in solution at room temperature.
- 3) After synthesis, the thioPA bottle on the synthesizer should be replaced with one containing acetonitrile diluent and the synthesizer line flushed with acetonitrile.

A typical cycle for the solid-phase synthesis of a PS2 linkage is different from a standard cycle for the synthesis of normal phosphate linkages. After coupling, the resulting thiophosphite triester is then sulfurized with DDTT. Capping is carried out AFTER sulfurization.

Upon completion of the automated synthesis, deprotection is carried out using a concentrated ammonia:ethanol (3:1, v:v) mix containing 20 mM DTT at 55 °C for 15-16 h.

Item	Catalog No.	Pack	Price(\$)
dA-Thiophosphoramidite	10-1700-90	100 µmole	150.00
	10-1700-02	0.25g	360.00
dC-Thiophosphoramidite	10-1710-90	100 µmole	150.00
	10-1710-02	0.25g	360.00
dG-Thiophosphoramidite	10-1720-90	100 µmole	150.00
	10-1720-02	0.25g	360.00
dT-Thiophosphoramidite	10-1730-90	100 µmole	150.00
	10-1730-02	0.25g	360.00



SULFURIZING REAGENTS


Glen Research's Sulfurizing Reagents are used to prepare phosphorothioate linkages using CE phosphoramidite chemistry. Each reagent exhibits the following attributes:

- 1) Reliably soluble, making them safe to use on automated synthesizers.
- 2) Reaction is fast (30 seconds), making the process convenient on small scales and readily amenable to scale-up.
- 3) Process is efficient, with better than 96% of the linkages being phosphorothioate and the remainder being phosphodiester.


When ordering Sulfurizing Reagent (Beaucage Reagent), please also order the silanized bottle appropriate for your instrument. There will be no charge for one bottle when supplied with a reagent order. The silanized bottle can be reused after rinsing with acetonitrile and drying. Additional silanized bottles are priced as shown below. Dissolve the reagent in acetonitrile at a concentration of 1g/100mL.

Sulfurizing Reagent II (3-((Dimethylamino-methylidene)amino)-3H-1,2,4-dithiazole-3-thione, DDTT) exhibits all the properties of Beaucage Reagent while adding stability in solution on the synthesizer AND offering strong ability to sulfurize RNA linkages. Sulfurizing Reagent II is available in powder form and as a stable solution.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Sulfurizing Reagent (Beaucage Reagent)	40-4036-10	1g	50.00
	40-4036-20	2g	100.00
Silanized Bottle (one bottle at no charge when ordered with Sulfurizing Reagent)			
<i>ABI (240mL capacity)</i>	40-4036-A1	each	10.00
<i>ABI (450mL capacity)</i>	40-4036-A2	each	10.00
<i>Expedite 8905 (150mL capacity)</i>	40-4036-C3	each	10.00
<i>Expedite 8909 and MerMade (240mL capacity)</i>	40-4036-C4	each	10.00
Sulfurizing Reagent II (DDTT) (Dissolve at a concentration of 1g/100mL to form an approximate 0.05M solution)	40-4037-10	1g	50.00
	40-4037-20	2g	100.00
0.05M Sulfurizing Reagent II in pyridine/acetonitrile	40-4137-51	100mL	100.00
	40-4137-52	200mL	200.00
	40-4137-57	450mL	450.00



Sulfurizing Reagent




Sulfurizing Reagent II

TRIMER PHOSPHORAMIDITES

REFERENCES

- (1) A.L. Kayushin, M.D. Korosteleva, A.I. Miroshnikov, W. Kosch, D. Zubov, and N. Piel, *Nucleic Acids Research*, 1996, **24**, 3748-3755.
- (2) A. Kayushin, et al., *Nucleos Nucleot*, 1999, **18**, 1531-1533.
- (3) A. Kayushin, M. Korosteleva, and A. Miroshnikov, *Nucleos Nucleot Nucleic Acids*, 2000, **19**, 1967-1976.
- (4) T. Mauriala, S. Auriola, A. Azhaye, A. Kayushin, M. Korosteleva, and A. Miroshnikov, *J Pharm Biomed Anal*, 2004, **34**, 199-206.
- (5) C. Neylon, *Nucleic Acids Res*, 2004, **32**, 1448-59.
- (6) L.R. Krumpke, K.M. Schumacher, J.B. McMahon, L. Makowski, and T. Mori, *BMC Biotechnol*, 2007, **7**, 65-72.
- (7) F.A. Fellouse, et al., *J Mol Biol*, 2007, **373**, 924-40.
- (8) W.P. Stemmer, A. Cramer, K.D. Ha, T.M. Brennan, and H.L. Heyneker, *Gene*, 1995, **164**, 49-53.
- (9) P.M. Sharp, and W.H. Li, *Nucleic Acids Res*, 1987, **15**, 1281-95.



General Structure of Trimer Phosphoramidites,
where B=A^{2'}, C^{2'}, G^{2'}, T

Trimer phosphoramidites¹⁻⁴ have proven to be extremely valuable because they allow codon-based mutagenesis, which circumvents the common problems of codon-bias, frame-shift mutations, and the introduction of nonsense or stop codons.⁵ This is accomplished by introducing a mixture of all 20 amino acid codons (or subset thereof) at any location within the sequenced to be mutated. This leads to the production of clonal libraries of exceptional diversity with order-of-magnitude increases in amino acid sequence variance while either maintaining a uniform amino acid distribution⁶ or one that is biased toward a desired set of amino acids.⁷

However, difficulties arise when trying to introduce mutations in multiple distal regions of a gene simultaneously. The synthesis of long oligonucleotides is required, which inevitably leads to lower sequence fidelity due to deletion mutants, depurination events and, to a lesser extent, mutations arising from deamination of cytidine, for example.

An elegant solution to this problem is the use of Antisense Trimer Phosphoramidites. These trimers are the reverse complement of the canonical 'sense' codons. When these antisense codons are put into the noncoding strand of a template DNA and amplified by PCR, they will code for the sense codon in the opposite strand of DNA. This allows the powerful technique of PCR Assembly⁸ to generate not only kilobase-sized genes from short 50mer oligonucleotides, but to simultaneously mutate multiple distal regions of that gene, as shown in Figure 1.

The sense and their corresponding antisense codons are listed in Table 1. Conveniently, many of our existing sense trimers can act as antisense codons. For example, AAC, which codes for asparagine, has the anticodon GTT, which is the sense codon for valine. However, some of the existing trimers, while they can act as an antisense codon, are not good choices for use. For example, TGG, which codes for tryptophan, could be used as an antisense codon for proline because CCA is one of proline's synonymous codons. However, CCA has a relatively low Codon Adaptation Index (CAI) value⁹ in *E. coli*, which could limit protein expression in that commonly used organism. For this reason, the anticodon CGG was chosen for optimal expression in *E. coli*, as were the other new antisense codons shown in bold in Table 1.

Included in Table 1 are the reaction factors (RFs) for each of the sense and antisense trimers. The reaction factor is critical since the trimers will likely be mixed and they exhibit different rates of reaction when coupling during oligonucleotide synthesis. An example where the RF is used to compensate for differing rates of coupling follows. The RF for AAC is 1.0 and for TAC is 1.6. Therefore, 1.6 equivalents of TAC are needed for every 1.0 equivalent of AAC for equal coupling rates. So to obtain 25 umoles of trimer mix that yields, on average, a 1:1 ratio of AAC/TAC at the mutation site, 9.6 umoles of AAC would be added to 15.4 umoles of TAC.

All of the trimers are available individually so the researchers can prepare custom trimer mixes. Two pre-made catalog trimer mixes are available: 13-1991-xx, for incorporating all 20 amino acid codons equally into a sequence and 13-1992-xx, for incorporating 19 amino acid codons (-Cys). For a custom trimer mix of a particular subset of codons or a trimer mix that represents a set of trimers that is biased toward a particular codon or codons, please contact support@glenresearch.com for a quotation and projected delivery date.

There is a concern that the sequence of the trimers has to be verified. For example, CAT coding for histidine, has to be differentiated from TAC, coding for tyrosine. These two trimers have virtually identical lipophilicity and their identity cannot be clearly confirmed by HPLC. This problem has been solved⁴ using HPLC electrospray mass spectrometric analysis of the trimers, which provides data confirming molecular weight and sequence.

FIGURE 1: SIMULTANEOUS MUTATION OF MULTIPLE DISTAL REGIONS OF GENE





TABLE 1: RF OF TRIMER PHOSPHoramidites

<i>Sense codons (5'→3')</i>	<i>Reaction Factor (RF)</i>	<i>Antisense codons (5'→3')</i>	<i>Reaction Factor (RF)</i>
AAA (Lys)	1.10	TTT	1.70
AAC (Asn)	1.00	GTT	1.90
ACT (Thr)	1.60	GGT	1.10
ATC (Ile)	1.50	GAT	1.40
ATG (Met)	1.30	CAT	1.30
CAG (Gln)	2.00	CTG	1.20
CAT (His)	1.30	ATG	1.30
CCG (Pro)	1.80	CGG	0.80
CGT (Arg)	1.40	GCG	0.60
CTG (Leu)	1.20	CAG	2.00
GAA (Glu)	1.40	TTC	1.30
GAC (Asp)	1.60	ATC	1.50
GCT (Ala)	1.50	TGC	1.50
GGT (Gly)	1.10	ACC	0.90
GTT (Val)	1.90	AAC	1.00
TAC (Tyr)	1.60	GTA	1.50
TCT (Ser)	1.30	AGA	1.40
TGC (Cys)	1.50	GCA	1.00
TGG (Trp)	1.10	CCA	1.10
TTC (Phe)	1.30	GAA	1.40



OLIGONUCLEOTIDE-DIRECTED MUTAGENESIS

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

Item	Catalog No.	Pack	Price(\$)
<i>Sense Trimers</i>			
AAA Trimer Phosphoramidite (Lys)	13-1000-95 13-1000-90	50 µm 100 µm	350.00 700.00
AAC Trimer Phosphoramidite (Asn)	13-1001-95 13-1001-90	50 µm 100 µm	350.00 700.00
ACT Trimer Phosphoramidite (Thr)	13-1013-95 13-1013-90	50 µm 100 µm	350.00 700.00
ATC Trimer Phosphoramidite (Ile)	13-1031-95 13-1031-90	50 µm 100 µm	350.00 700.00
ATG Trimer Phosphoramidite (Met)	13-1032-95 13-1032-90	50 µm 100 µm	350.00 700.00
CAG Trimer Phosphoramidite (Gln)	13-1102-95 13-1102-90	50 µm 100 µm	350.00 700.00
CAT Trimer Phosphoramidite (His)	13-1103-95 13-1103-90	50 µm 100 µm	350.00 700.00
CCG Trimer Phosphoramidite (Pro)	13-1112-95 13-1112-90	50 µm 100 µm	350.00 700.00
CGT Trimer Phosphoramidite (Arg)	13-1123-95 13-1123-90	50 µm 100 µm	350.00 700.00
CTG Trimer Phosphoramidite (Leu)	13-1132-95 13-1132-90	50 µm 100 µm	350.00 700.00
GAA Trimer Phosphoramidite (Glu)	13-1200-95 13-1200-90	50 µm 100 µm	350.00 700.00
GAC Trimer Phosphoramidite (Asp)	13-1201-95 13-1201-90	50 µm 100 µm	350.00 700.00
GCT Trimer Phosphoramidite (Ala)	13-1213-95 13-1213-90	50 µm 100 µm	350.00 700.00
GGT Trimer Phosphoramidite (Gly)	13-1223-95 13-1223-90	50 µm 100 µm	350.00 700.00
GTT Trimer Phosphoramidite (Val)	13-1233-95 13-1233-90	50 µm 100 µm	350.00 700.00
TAC Trimer Phosphoramidite (Tyr)	13-1301-95 13-1301-90	50 µm 100 µm	350.00 700.00
TCT Trimer Phosphoramidite (Ser)	13-1313-95 13-1313-90	50 µm 100 µm	350.00 700.00
TGC Trimer Phosphoramidite (Cys)	13-1321-95 13-1321-90	50 µm 100 µm	350.00 700.00
TGG Trimer Phosphoramidite (Trp)	13-1322-95 13-1322-90	50 µm 100 µm	350.00 700.00
TTC Trimer Phosphoramidite (Phe)	13-1331-95 13-1331-90	50 µm 100 µm	350.00 700.00
Trimer Phosphoramidite Mix 1 (Mix of above 20 trimers)	13-1991-95 13-1991-90	50 µm 100 µm	515.00 1030.00
Trimer Phosphoramidite Mix 2 (Mix of above 20 trimers less TGC-Cys)	13-1992-95 13-1992-90	50 µm 100 µm	515.00 1030.00

OLIGONUCLEOTIDE-DIRECTED MUTAGENESIS

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
<i>Antisense Trimers</i>			
AAC Trimer Phosphoramidite (Anti Val)	13-1001-95	50 µm	350.00
	13-1001-90	100 µm	700.00
ACC Trimer Phosphoramidite (Anti Gly)	13-1011-95	50 µm	350.00
	13-1011-90	100 µm	700.00
AGA Trimer Phosphoramidite (Anti Ser)	13-1020-95	50 µm	350.00
	13-1020-90	100 µm	700.00
ATC Trimer Phosphoramidite (Anti Asp)	13-1031-95	50 µm	350.00
	13-1031-90	100 µm	700.00
ATG Trimer Phosphoramidite (Anti His)	13-1032-95	50 µm	350.00
	13-1032-90	100 µm	700.00
CAG Trimer Phosphoramidite (Anti Leu)	13-1102-95	50 µm	350.00
	13-1102-90	100 µm	700.00
CAT Trimer Phosphoramidite (Anti Met)	13-1103-95	50 µm	350.00
	13-1103-90	100 µm	700.00
CCA Trimer Phosphoramidite (Anti Trp)	13-1110-95	50 µm	350.00
	13-1110-90	100 µm	700.00
CGG Trimer Phosphoramidite (Anti Pro)	13-1122-95	50 µm	350.00
	13-1122-90	100 µm	700.00
GAA Trimer Phosphoramidite (Anti Phe)	13-1200-95	50 µm	350.00
	13-1200-90	100 µm	700.00
GAT Trimer Phosphoramidite (Anti Ile)	13-1203-95	50 µm	350.00
	13-1203-90	100 µm	700.00
GCA Trimer Phosphoramidite (Anti Cys)	13-1210-95	50 µm	350.00
	13-1210-90	100 µm	700.00
GCG Trimer Phosphoramidite (Anti Arg)	13-1212-95	50 µm	350.00
	13-1212-90	100 µm	700.00
GGT Trimer Phosphoramidite (Anti Thr)	13-1223-95	50 µm	350.00
	13-1223-90	100 µm	700.00
GTA Trimer Phosphoramidite (Anti Tyr)	13-1230-95	50 µm	350.00
	13-1230-90	100 µm	700.00
TGC Trimer Phosphoramidite (Anti Ala)	13-1321-95	50 µm	350.00
	13-1321-90	100 µm	700.00
TTC Trimer Phosphoramidite (Anti Glu)	13-1331-95	50 µm	350.00
	13-1331-90	100 µm	700.00
TTT Trimer Phosphoramidite (Anti Lys)	13-1333-95	50 µm	350.00
	13-1333-90	100 µm	700.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

INTELLECTUAL PROPERTY

C-5 Propyne and AP-dC Phosphoramidites

This product is covered by patents or patents pending owned by Isis Pharmaceuticals, Inc. ("Isis"). Purchase of this product includes a limited license to use this product solely for internal research. This license specifically excludes (and you have no right to use this product for): (a) therapeutic or diagnostic applications (including products or services that incorporate this product), (b) any in vivo toxicity/safety study in support of an investigational new drug application (or foreign counterpart), (c) resale (including sale of any products or services that incorporate this product) or (d) gene functionalization activities (including products or services that incorporate data derived from gene functionalization activities) if such activities have commercial application, any and all of which require a separate license from Isis. Neither this product nor any product created through its use may be used in human clinical trials.

A simple agreement must be signed before end-users and custom oligo services may purchase these products for use as defined above. <http://www.glenresearch.com/Reference/PropyneFax.pdf>

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M

Columns

For Instrument type Add

Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)

C-5 PROPYNE DERIVATIVES AND G-CLAMP


Substitution of C-5 propynyl-dC (pdC) for dC and C-5 propynyl-dU (pdU) for dT are effective strategies to enhance base pairing. Using these base substitutions, duplex stability and melting temperatures are raised by the following amounts: C-5 propynyl-C 2.8° per substitution; C-5 propynyl-U 1.7° per substitution. AP-dC (G-clamp) substitutes for dC and is another very important modified nucleoside that enhances hybridization by 7-21° per substitution depending upon the sequence and location of the AP-dC. The ability of these modified bases to enhance binding while maintaining specificity has proven useful in antisense research and in the synthesis of high affinity probes. AP-dC is also a fluorescent nucleoside and should find uses in DNA structural research.

Item	Catalog No.	Pack	Price(\$)
pdC-CE Phosphoramidite	10-1014-90	100 µmole	85.00
	10-1014-02	0.25g	245.00
	10-1014-05	0.5g	490.00
pdU-CE Phosphoramidite	10-1054-90	100 µmole	65.00
	10-1054-02	0.25g	195.00
	10-1054-05	0.5g	390.00
AP-dC-CE Phosphoramidite (G-Clamp)	10-1097-95	50 µmole	230.00
	10-1097-90	100 µmole	460.00
	10-1097-02	0.25g	1175.00

BASES AFFECTING DUPLEX STABILITY

C-5 methyl pyrimidine nucleosides are known to stabilize duplexes relative to the non-methylated bases. Therefore, enhanced binding can be achieved using 5-methyl-dC in place of dC, duplex melting temperature being increased by 1.3°. Ac-5-Me-dC-CE Phosphoramidite is fully compatible with AMA deprotection and none of the N4-Me transamination mutant is observed on deprotection. 2,6-Diaminopurine 2'-deoxyribose (2-amino-dA) forms an additional hydrogen bond with Thymidine, thereby leading to duplex stabilization with a melting temperature increase of 3°. Our 2-amino-dA monomer exhibits fast and effective deprotection in ammonium hydroxide and it is stabilized to depurination during synthesis.

Item	Catalog No.	Pack	Price(\$)
5-Me-dC-CE Phosphoramidite	10-1060-90	100 µmole	50.00
	10-1060-02	0.25g	120.00
5-Me-dC-CPG	20-2060-01	0.1g	50.00
	20-2160-41	Pack of 4	200.00
	20-2160-42	Pack of 4	120.00
Ac-5-Me-dC-CE Phosphoramidite	10-1560-90	100 µmole	50.00
	10-1560-02	0.25g	120.00
2-Amino-dA-CE Phosphoramidite (2,6-diaminopurine)	10-1085-95	50 µmole	70.00
	10-1085-90	100 µmole	125.00
	10-1085-02	0.25g	250.00



BASES AFFECTING DUPLEX STABILITY (CONT.)

Sequences with high GC content may contain mismatches and still hybridize because of the high stability of the G-C base pair. The N4-ethyl analogue of dC (N4-Et-dC) hybridizes specifically to natural dG but the stability of the base pair is reduced to about the level of an AT base pair.


Coupling N6-Me-dA (10-1003) and N4-Et-dC (10-1068) with 1H-tetrazole leads to a trace of branching at the secondary amine positions, while DCI leads to around 15% branching. In collaboration with Berry and Associates, the acetyl protected monomers were prepared. Acetyl protection was chosen since it would block branching reactions. Oligonucleotides synthesized using these monomers proved to be compatible with all popular deprotection strategies from UltraMild to UltraFast. When the acetyl protected monomers were compared with the unprotected monomers using DCI as activator, branching was reduced from 15% to zero.

Item	Catalog No.	Pack	Price(\$)
N4-Et-dC-CE Phosphoramidite	10-1068-95	50 µmole	125.00
	10-1068-90	100 µmole	225.00
	10-1068-02	0.25g	675.00
N4-Ac-N4-Et-dC-CE Phosphoramidite	10-1513-95	50 µmole	125.00
	10-1513-90	100 µmole	225.00
	10-1513-02	0.25g	675.00
N6-Me-dA-CE Phosphoramidite	10-1003-90	100 µmole	162.50
	10-1003-02	0.25g	495.00
N6-Ac-N6-Me-dA-CE Phosphoramidite	10-1503-90	100 µmole	162.50
	10-1503-02	0.25g	495.00

ZIP NUCLEIC ACIDS (ZNA®)

Spermine phosphoramidite is used to produce oligospermine-oligonucleotide conjugates - Zip Nucleic Acids (ZNA®) Oligos. The name reflects the presumed mode of action. The conjugates are believed to use the oligospermine to seek out and move along (scan) oligonucleotide strands until the probe complementary sequence is located. The oligospermine then performs the function of stabilizing the formed duplex by reducing electrostatic repulsion, thereby leading to significantly increased binding affinities. ZNA® Oligos have found use in the following applications: Multiplex PCR; PCR of AT-rich Regions; RT qPCR; Detection of MicroRNA; Improved SNP Discrimination; and Antisense and Antigenic Effects. Spermine phosphoramidite is simple to use in oligonucleotide synthesis and can be added multiple times at the 3' or 5' terminus. Deprotection and isolation are also straightforward. HPLC analysis of the conjugates requires high pH to suppress the ionization of the spermine residues.

Item	Catalog No.	Pack	Price(\$)
Spermine Phosphoramidite	10-1939-95	50 µmole	145.00
	10-1939-90	100 µmole	270.00
	10-1939-02	0.25g	525.00



SEE ALSO

N6-Me-dA

p62

INTELLECTUAL PROPERTY

"Spermine phosphoramidite" synthon is the subject matter of U.S. Patent Application No. 12/086.599, European Patent Application No. EP20060847298 and foreign equivalents for which Polyplus-transfection is the co-owner. Product is sold for research purposes only. Product shall not be used to manufacture oligospermine-oligonucleotide conjugates for use in diagnostics, clinical or commercial applications including use in humans. There is no implied license to manufacture oligospermine-oligonucleotide conjugates for diagnostic, clinical or commercial applications, including but not limited to contract research. Please contact Polyplus-transfection at licensing@polyplus-transfection.com to obtain a license for such use.

Spermine phosphoramidite is covered under the intellectual property of Polyplus-transfection SA. ZNA® is a registered trademark of Polyplus-transfection SA.

CAPS FOR INCREASED DUPLEX STABILITY AND BASE-PAIRING FIDELITY AT TERMINI

New cap structures allow for the preparation of hybridization probes with increased affinity for complementary sequences. The monomers used to prepare capped oligonucleotides are phosphoramidites that can be readily introduced via automated DNA synthesis at the end of solid phase syntheses. The caps favor the formation of stable Watson-Crick duplexes by stacking on the terminal base pair (Figures 1 and 2).




FIGURE 1: STACKING OF CAP ON 5' TERMINAL BASE PAIR



FIGURE 2: STACKING OF Uaq CAP ON 3' TERMINAL BASE PAIR


Melting point increases of over 10 °C per modification can be realized for short duplexes.^{1,2} The caps fit canonical Watson-Crick base pairs and do not stack well on mismatched base pairs. This leads to increased base pairing selectivity at the terminal and the penultimate position of oligonucleotides featuring the caps. Base pairing fidelity is usually low at the termini, where fraying occurs frequently in the absence of caps. The beneficial effects of the caps are also realized when longer target strands are bound, so there is no need for blunt ends for the duplexes formed.^{1,2} The caps, when attached to the 5' terminus of an oligonucleotide, also facilitate purification as their lipophilicity leads to prolonged retention on reversed phase columns or cartridges. Finally, capping of termini may discourage the degradation of oligonucleotides by exonucleases.

3'-Uaq Cap CPG, a Uridine support modified with a 2'- anthraquinone residue, is the most effective oligonucleotide cap known to date.^{3,4} For short hybrid duplexes between DNA probes and RNA target strands, the increase in T_m is up to 18 °C and the modification is effective in increasing the T_m of DNA:DNA, RNA:RNA, and DNA:RNA hybrid duplexes. 3'-Uaq Cap also increases probe specificity by depressing the melting point of terminal mismatches.


REFERENCES

- (1) Dogan, Z.; Paulini, R.; Rojas Stütz, J. A.; Narayanan, S.; Richert, C. *J. Amer. Chem. Soc.* **2004**, *126*, 4762-4763.
- (2) Narayanan, S.; Gall, J.; Richert, C. *Nucleic Acids Res.* **2004**, *32*, 2901-2911.
- (3) A. Patra, C. Richert, *J. Amer. Chem. Soc.*, 2009, **131**, 12671-12681.
- (4) C. Ahlborn, K. Siegmund, C. Richert, *J. Amer. Chem. Soc.*, 2007, **129**, 15218-15232.


Item	Catalog No.	Pack	Price(\$)	
5'-Trimethoxystilbene Cap Phosphoramidite	10-1986-90	100 µmole	195.00	
	10-1986-02	0.25g	495.00	
5'-Pyrene Cap Phosphoramidite	10-1987-90	100 µmole	195.00	
	10-1987-02	0.25g	495.00	
3'-Uaq Cap CPG	20-2980-01	0.1g	180.00	
	20-2980-10	1.0g	1500.00	
	1 µmole columns	20-2980-41	Pack of 4	300.00
	0.2 µmole columns	20-2980-42	Pack of 4	150.00
	10 µmole column (ABI)	20-2980-13	Pack of 1	750.00
15 µmole column (Expedite)	20-2980-14	Pack of 1	1125.00	



5'-Trimethoxystilbene Cap



5'-Pyrene Cap



3'-Uaq Cap CPG


DNA METHYLATION

One of the fastest growing fields in biology and cancer research is epigenetics. While the underlying genetic code defines which proteins and gene products are synthesized, it is epigenetic control that defines when and where they are expressed. This dynamic control of gene expression is essential for X chromosome inactivation, embryogenesis, cellular differentiation and appears integral to memory formation and synaptic plasticity.


In 2009, two reports^{1,2} described the discovery of 5-hydroxymethyl-2'-deoxyCytidine (hmdC), a novel dC modification in Purkinje neurons and embryonic stem cells. Later, a third report found this modification to be strongly enriched in brain tissues associated with higher cognitive functions.³ This dC modification is generated by the action of α -ketoglutarate dependent ten eleven translocation (TET) enzymes, which oxidizes 5-Me-dC to hmdC. This finding stimulated discussion about active demethylation pathways that could occur, e.g., via base excision repair (BER), with the help of specialized DNA glycosylases. Alternatively, one could envision a process in which the hydroxymethyl group of hmdC is further oxidized to 5-formyl-dC (fdC) or 5-carboxy-dC (cadC) followed by elimination of either formic acid or carbon dioxide^{4,5}.

Glen Research has supported this research since its inception by providing the building blocks for the synthesis of oligonucleotides containing all the new dC derivatives - hmdC, fdC and cadC. The first generation hmdC phosphoramidite was fairly very well accepted but requires fairly harsh deprotection conditions. Therefore, a second generation building block (5-Hydroxymethyl-dC II) developed by Carell and co-workers that is compatible with UltraMild deprotection was introduced.⁶ 5-Formyl-dC III has been designed to meet all of the requirements to prepare an oligo containing all of the methylated variants.⁷


Item	Catalog No.	Pack	Price(\$)
5-Hydroxymethyl-dC-CE Phosphoramidite	10-1062-95	50 μ mole	335.00
	10-1062-90	100 μ mole	650.00
	10-1062-02	0.25g	1675.00
5-Carboxy-dC-CE Phosphoramidite	10-1066-95	50 μ mole	230.00
	10-1066-90	100 μ mole	450.00
	10-1066-02	0.25g	1200.00
5-Formyl-dC-CE Phosphoramidite	10-1514-95	50 μ mole	610.00
	10-1514-90	100 μ mole	1200.00
	10-1514-02	0.25g	3225.00
5-Hydroxymethyl-dC II-CE Phosphoramidite	10-1510-95	50 μ mole	345.00
	10-1510-90	100 μ mole	670.00
	10-1510-02	0.25g	2100.00
5-Formyl-dC III-CE Phosphoramidite	10-1564-95	50 μ mole	360.00
	10-1564-90	100 μ mole	700.00
	10-1564-02	0.25g	1800.00




5-Hydroxymethyl-dC




5-Hydroxymethyl-dC II



5-Carboxy-dC



5-Formyl-dC



5-Formyl-dC III

SEE ALSO

5-Me-dC p42
5-hmdU pPB

REFERENCES

- (1) S. Kriaucionis, and N. Heintz, *Science*, 2009, **324**, 929-30.
- (2) M. Tahiliani, et al., *Science*, 2009, **324**, 930-935.
- (3) M. Münzel, et al., *Angewandte Chemie-International Edition*, 2010, **49**, 5375-5377.
- (4) D. Globisch, et al., *PLoS One*, 2010, **5**, e15367.
- (5) S.C. Wu, and Y. Zhang, *Nat Rev Mol Cell Biol*, 2010, **11**, 607-20.
- (6) M. Münzel, D. Globisch, C. Trindler, and T. Carell, *Org Lett*, 2010, **12**, 5671-3.
- (7) A.S. Schroder, et al., *Angewandte Chemie-International Edition*, 2014, **53**, 315-318.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M

Columns

For Instrument type Add

Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers
For Instrument type Add

Expedite	E
MerMade	M

Columns
For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

DUPLEX EFFECTS


The design of primers is frequently complicated by the degeneracy of the genetic code. Three strategies are now available to confront this problem. In the first, a mixed base addition (N) is used to form the degenerate site. This approach is best if the number of degenerate sites is small. A second option is the use of 2'-deoxyinosine or 2'-deoxyNebularine which exhibit low, but unequal, hydrogen bonding to the other four bases. The third option is the use of a universal nucleoside. In this strategy, the base analog does not hybridize significantly to the other four bases and makes up some of the duplex destabilization by acting as an intercalating agent. 3-Nitropyrrole 2'-deoxynucleoside (M) is the first example of a set of universal bases. Subsequently, 5-nitroindole was determined to be an effective universal base and to be superior to 3-nitropyrrole, based on duplex melting experiments.

The modified bases designated P and K show considerable promise as degenerate bases. The pyrimidine derivative P, when introduced into oligonucleotides, base pairs with either A or G, while the purine derivative K base pairs with either C or T. A dP+dK mix also can serve as a mixed base with much less degeneracy than dA+dC+dG+dT (N).


Item	Catalog No.	Pack	Price(\$)
dA+dG-CE Phosphoramidites	10-1002-02	0.25g	40.00
dC+dT-CE Phosphoramidites	10-1013-02	0.25g	40.00
dA+dC+dG+dT-CE Phosphoramidites	10-1023-02	0.25g	40.00

Other pack sizes, mixed base combinations and custom doping of individual monomers are available on request. Also, mixed base columns are available in 0.2 and 1.0 μmole sizes on request.

dI-CE Phosphoramidite	10-1040-90 10-1040-02	100 μmole 0.25g	50.00 120.00
dI-CPG 500	20-2040-01	0.1g	30.00
1 μmole columns	20-2190-41	Pack of 4	120.00
0.2 μmole columns	20-2190-42	Pack of 4	72.00
dI-CPG 1000	20-2041-01	0.1g	30.00
1 μmole columns	20-2191-41	Pack of 4	120.00
0.2 μmole columns	20-2191-42	Pack of 4	72.00
dU-CE Phosphoramidite	10-1050-90 10-1050-02	100 μmole 0.25g	35.00 100.00
dU-CPG 500	20-2050-01	0.1g	30.00
1 μmole columns	20-2150-41	Pack of 4	120.00
0.2 μmole columns	20-2150-42	Pack of 4	72.00
dU-CPG 1000	20-2051-01	0.1g	50.00
1 μmole columns	20-2151-41	Pack of 4	200.00
0.2 μmole columns	20-2151-42	Pack of 4	120.00




2'-deoxyinosine




2'-deoxyUridine

DUPLEX EFFECTS (CONT.)


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
2'-DeoxyNebularine-CE Phosphoramidite (Purine)	10-1041-90	100 μ mole	105.00
	10-1041-02	0.25g	255.00
5-Nitroindole-CE Phosphoramidite	10-1044-90	100 μ mole	125.00
	10-1044-02	0.25g	325.00
dP-CE Phosphoramidite	10-1047-90	100 μ mole	195.00
	10-1047-02	0.25g	595.00
dK-CE Phosphoramidite	10-1048-90	100 μ mole	195.00
	10-1048-02	0.25g	595.00
dP+dK-CE Phosphoramidite	10-1049-90	100 μ mole	195.00
	10-1049-02	0.25g	595.00




2'-deoxyNebularine




3-Nitropyrrole



5-Nitroindole



dP



dK

DUPLEX EFFECTS (CONT.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)









INTELLECTUAL PROPERTY

Product containing Ds, s or Pa bases is covered by patents or patents pending owned by TagCyx Biotechnologies. Purchase of this product includes a limited license to use this product solely for research. This license specifically excludes: (a) therapeutic or diagnostic applications (including products or services that incorporate this product), (b) any in vivo toxicity/safety study in support of an investigational new drug application (or foreign counterpart), (c) resale, or (d) gene functionalization activities (including products or services that incorporate data derived from gene functionalization activities) if such activities have commercial application. All of the above require a separate license from TagCyx Biotechnologies. Neither this product nor any product created through its use may be used in human clinical trials.

Unnatural base pairs display unique abilities in duplex DNA and in nucleic acid and protein biosyntheses. A standard Watson and Crick base pair is formed between iso-C and iso-G, but the hydrogen bonding pattern is quite different from the natural base pairs A-T and C-G. (The 5-methyl analogue was chosen as the synthetic target due to the reported instability of 2'-deoxyisocytidine caused by deamination during oligonucleotide synthesis or deprotection.)

The unnatural base pair between 7-(2-thienyl)-imidazo[4,5-b]pyridine (Ds) and pyrrole-2-carbaldehyde (Pa) is formed by specific hydrophobic shape complementation. The shape of the Ds-Pa pair is different from those of the natural A-T and G-C pairs, but the Ds-Pa pair works together with the natural pairs in vitro replication and transcription. Pa also functions as a template base for incorporating another unnatural base, 2-amino-6-(2-thienyl)purine (s), into RNA. The s base also acts as a unique fluorescent base analog in DNA and RNA fragments. dDss is strongly fluorescent and is useful as a fluorescent tag for DNA detection. dDss also forms a base pair with dPa. Biotin PaTP can be site-specifically incorporated into RNA, opposite dDs at a desired position in DNA templates, by T7 transcription. Similarly, the fluorescent s base can be site-specifically incorporated into RNA opposite dPa in DNA templates.

Item	Catalog No.	Pack	Price(\$)
dmf-5-Me-iso-dC-CE Phosphoramidite	10-1065-90	100 μ mole	100.00
	10-1065-02	0.25g	255.00
dmf-iso-dG-CE Phosphoramidite	10-1078-90	100 μ mole	165.00
	10-1078-02	0.25g	355.00
dDs-CE Phosphoramidite	10-1521-90	100 μ mole	145.00
	10-1521-02	0.25g	420.00
Pac-ds-CE Phosphoramidite	10-1522-90	100 μ mole	170.00
	10-1522-02	0.25g	420.00
dDss-CE Phosphoramidite	10-1524-95	50 μ mole	130.00
	10-1524-90	100 μ mole	250.00
	10-1524-02	0.25g	675.00
dPa-CE Phosphoramidite	10-1523-90	100 μ mole	130.00
	10-1523-02	0.25g	420.00
sTP 10mM	81-3522-02	25 μ L	350.00
Biotin PaTP 10mM	81-3525-02	25 μ L	450.00



CLEANAMP™ MONOMERS

CleanAmp™ Primers offer an alternative to other Hot Start technologies and allow greater control of primer hybridization and extension during PCR. It has been demonstrated that CleanAmp™ Primers outperform other technologies in multiple applications. Indeed, over a broad range of applications, CleanAmp™ Primers reduce or eliminate off-target amplification. Greater amplicon yield is also achieved, due to improvement in specificity and sensitivity. By using either the slow-releasing Precision primers with two CleanAmp™ phosphotriester linkages or the faster-releasing Turbo Primers with a single CleanAmp™ phosphotriester linkage, the rate of formation of unmodified primer can be controlled to suit reaction needs. A table to aid in the selection of Turbo and Precision Primers for specific applications is shown below.

Turbo Primers


Fast cycling
 Multiplex PCR
 Improves amplicon yield
 Reduces mis-priming/ primer dimer formation

Precision Primers


Standard cycling
 One-step reverse-transcription PCR
 Improved specificity and limit of detection
 Greatest reduction in mis-priming/primer dimer formation

Synthesis of CleanAmp™ Primers requires the use of UltraMild Chemistry.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
CleanAmp™-Pac-dA-CE Phosphoramidite	10-1440-90	100 µmole	100.00
	10-1440-02	0.25g	240.00
	10-1440-05	0.5g	480.00
CleanAmp™-Ac-dC-CE Phosphoramidite	10-1450-90	100 µmole	100.00
	10-1450-02	0.25g	240.00
	10-1450-05	0.5g	480.00
CleanAmp™-Pac-dG-CE Phosphoramidite	10-1460-90	100 µmole	100.00
	10-1460-02	0.25g	240.00
	10-1460-05	0.5g	480.00
CleanAmp™-dT-CE Phosphoramidite	10-1470-90	100 µmole	100.00
	10-1470-02	0.25g	240.00
	10-1470-05	0.5g	480.00




CleanAmp™-Pac-dA



CleanAmp™-Ac-dC



CleanAmp™-Pac-dG



CleanAmp™-dT

SEE ALSO

UltraMild DNA Synthesis p20

INTELLECTUAL PROPERTY

CleanAmp™ is a trademark of TriLink BioTechnologies, Inc. CleanAmp™ products or portions thereof are covered by TriLink pending Patent Applications, US 2007281308 and WO2007139723, US Provisional Patent Application Serial # 61/056,324 and US Patent 6762298 licensed from the Department of Health and Human Services. CleanAmp™ products are sold exclusively for R & D use by the purchaser. They may not be resold, distributed or re-packaged. No license is granted or implied with the purchase of this product. Amplification methods used in connection with Polymerase Chain Reaction ("PCR") Process are covered by many patents. It may be necessary to obtain a separate license for certain patented applications in which the product is used. CleanAmp™ Licenses are available directly from TriLink BioTechnologies. www.trilinkbiotech.com

CHAIN TERMINATORS

SEE ALSO

5'-Phosphoramidites	p30
5'-Supports	p31
3'-Spacer C3 CPG	p80

REFERENCE

(1) P.Y. Chen, et al., *RNA*, 2008, **14**, 263-274..

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns


For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M


(Please inquire for availability of vials and columns for other instrument types.)

In situations where ligation must be blocked at the 5' terminus, 5'-OMe-dT may be used. 5'-OMe modification of a strand of siRNA using 5'-OMe-T can control guide strand selection and targeting specificity.¹ 5'-Amino-dT terminates an oligonucleotide with a 5'-amino group which may be used for attaching a peptide or a PNA sequence. To avoid polymerase extension at the 3' terminus, 2',3'-dideoxynucleoside and 3'-deoxynucleoside CPGs have proved to be effective. 2',3'-Phosphoramidites are designed to be used with the 5'-phosphoramidites and supports. Since these phosphoramidites have no DMT group, they are not compatible with purification by the DMT-on technique. Ion exchange HPLC or PAGE should be used to purify these dideoxy terminated oligos to ensure that shorter sequences (containing 3'-OH) groups are removed. (3'-Termination can also be effected using a 3'-3' linkage formed using 5'-supports, or 3'-spacer C3 CPG.)


Item	Catalog No.	Pack	Price(\$)
5'-OMe-dT-CE Phosphoramidite	10-1031-90	100 µmole	135.00
	10-1031-02	0.25g	355.00
5'-Amino-dT-CE Phosphoramidite	10-1932-90	100 µmole	150.00
	10-1932-02	0.25g	400.00
3'-dA-CPG	20-2004-01	0.1g	400.00
	1 µmole columns	Pack of 4	675.00
	0.2 µmole columns	Pack of 4	225.00
3'-dC-CPG	20-2064-01	0.1g	300.00
	1 µmole columns	Pack of 4	600.00
	0.2 µmole columns	Pack of 4	200.00
3'-dG-CPG	20-2074-01	0.1g	300.00
	1 µmole columns	Pack of 4	600.00
	0.2 µmole columns	Pack of 4	200.00
3'-dT-CPG	20-2084-01	0.1g	300.00
	1 µmole columns	Pack of 4	600.00
	0.2 µmole columns	Pack of 4	200.00




5'-OMe-Thymidine




5'-Amino-dT




3'-dA-CPG



3'-dC-CPG



3'-dG-CPG



3'-dT-CPG

CHAIN TERMINATORS (CONT.)

Item	Catalog No.	Pack	Price(\$)
2',3'-ddC-CPG	20-2017-01	0.1g	300.00
1 μmole columns	20-2117-41	Pack of 4	600.00
0.2 μmole columns	20-2117-42	Pack of 4	200.00
2',3'-ddA-CE Phosphoramidite	10-7001-90	100 μmole	130.00
	10-7001-02	0.25g	545.00
2',3'-ddC-CE Phosphoramidite	10-7101-90	100 μmole	130.00
	10-7101-02	0.25g	545.00
2',3'-ddG-CE Phosphoramidite	10-7201-90	100 μmole	145.00
	10-7201-02	0.25g	675.00
2',3'-ddT-CE Phosphoramidite	10-7301-90	100 μmole	130.00
	10-7301-02	0.25g	545.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add






Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)



STRUCTURE/ACTIVITY RELATIONSHIP

STABILITY NOTES

7-Deaza-dG is unstable to iodine oxidation. Add a maximum of 2 times when using iodine oxidation or use 0.5M (10-camphorsulfonyl)-oxaziridine (CSO) in anhydrous acetonitrile and 3 min. oxidation time. (See Glen Report-Vol.9, No.1, 1996,page 8.)


INTELLECTUAL PROPERTY

The use of PPG is subject to proprietary rights of Epoch Biosciences, Inc. and it is sold under license from Epoch Biosciences, Inc.


(1) I.V. Kutyavin, et al., *Nucleic Acids Res.*, 2002, **30**, 4952-4959.

The following products are used to investigate the effect on the activity of an oligonucleotide when key structural elements are changed. The 7-deaza purine monomers lack groups critical for hydrogen bonding. 7-Deaza-8-aza-A and 7-deaza-8-aza-G (PPG) monomers are isomers of A and G and have similar electron density. Their presence in oligos is slightly stabilizing relative to A and G. Unlike G, PPG does not lead to aggregation and G-rich oligos can be easily prepared and isolated. 5'-Fluorescein oligos with PPG at the 5'-terminus are much less quenched than the equivalent G oligos. As a purine analogue of Thymidine, 7-deaza-2'-deoxyXanthosine (7-deaza-dX) promises to have interesting effects on DNA structure of triplexes. 7-Deaza-dX also forms a non-standard base pair with a 2,4-diaminopyrimidine nucleoside analogue. Standard nucleobases have an unshared pair of electrons that project into the minor groove of duplex DNA. Enzymes that interact with DNA, polymerases, reverse transcriptases, restriction enzymes, etc., may use a hydrogen bond donating group to contact the hydrogen bond acceptor in the minor groove. 3-Deaza-2'-deoxyadenosine is very interesting in that it maintains the ability for regular Watson-Crick hydrogen bonding to T but is lacking the electron pair at the 3-position normally provided by N3.


Item	Catalog No.	Pack	Price(\$)
7-Deaza-dA-CE Phosphoramidite	10-1001-95	50 μ mole	177.50
	10-1001-90	100 μ mole	355.00
	10-1001-02	0.25g	975.00
7-Deaza-8-aza-dA-CE Phosphoramidite	10-1083-95	50 μ mole	177.50
	10-1083-90	100 μ mole	355.00
	10-1083-02	0.25g	975.00
7-Deaza-dG-CE Phosphoramidite	10-1021-95	50 μ mole	177.50
	10-1021-90	100 μ mole	355.00
	10-1021-02	0.25g	975.00
7-Deaza-8-aza-dG-CE Phosphoramidite (PPG)	10-1073-95	50 μ mole	207.50
	10-1073-90	100 μ mole	395.00
	10-1073-02	0.25g	1150.00
7-deaza-dX-CE Phosphoramidite	10-1076-95	50 μ mole	177.50
	10-1076-90	100 μ mole	355.00
	10-1076-02	0.25g	975.00
3-Deaza-dA-CE Phosphoramidite	10-1088-95	50 μ mole	177.50
	10-1088-90	100 μ mole	355.00
	10-1088-02	0.25g	975.00




7-Deaza-2'-deoxyAdenosine




7-Deaza-8-Aza-2'-deoxyAdenosine




7-Deaza-2'-deoxyGuanosine



7-Deaza-8-Aza-2'-deoxyGuanosine



7-deaza-dX




3-Deaza-dA

STRUCTURE/ACTIVITY RELATIONSHIP (CONT.)


The C-nucleoside 2'-deoxypseudouridine, in contrast to dU, forms stable C:pseudoU-A triplets. 2-Aminopurine lacks groups critical for hydrogen bonding and is a mildly fluorescent base.

Demand for sulfur modified bases continues to expand for investigations of oligonucleotide structure, but primarily for cross-linking purposes. 6-Thio-dG, 4-Thio-dT and 4-thio-dU are very useful modifications for photo cross-linking and photoaffinity labelling experiments. Oligos containing 2-thio-dT are useful in examining protein-DNA interaction by acting as photosensitizing probes. The thiocarbonyl group in 2-thio-dT is especially interesting in that it is available to react with compounds associating with the minor groove of DNA. 2-Amino-A forms a very stable base pair with T containing three hydrogen bonds but the stability of the base pair with 2-thio-T is greatly diminished. Due to steric interactions between the 2-thio group of thymidine and the 2-amino group of 2-amino-A, the base pair contains only a single hydrogen bond. Oligos containing 2-amino-dA and 2-thio-dT exhibit high affinity for natural oligonucleotides but show little affinity for other similar oligos even of a complementary sequence.


Item	Catalog No.	Pack	Price(\$)
2'-deoxypseudoU-CE Phosphoramidite	10-1055-95	50 μ mole	177.50
	10-1055-90	100 μ mole	355.00
	10-1055-02	0.25g	975.00
2-Aminopurine-CE Phosphoramidite	10-1046-90	100 μ mole	135.00
	10-1046-02	0.25g	355.00
6-Thio-dG-CE Phosphoramidite	10-1072-95	50 μ mole	177.50
	10-1072-90	100 μ mole	355.00
	10-1072-02	0.25g	975.00
4-Thio-dT-CE Phosphoramidite	10-1034-95	50 μ mole	165.00
	10-1034-90	100 μ mole	295.00
	10-1034-02	0.25g	675.00
4-Thio-dU-CE Phosphoramidite	10-1052-95	50 μ mole	165.00
	10-1052-90	100 μ mole	295.00
	10-1052-02	0.25g	675.00
2-Thio-dT-CE Phosphoramidite	10-1036-95	50 μ mole	165.00
	10-1036-90	100 μ mole	295.00
	10-1036-02	0.25g	675.00




2'-dpseudoU




2-Aminopurine




6-Thio-dG



4-Thio-dT



4-Thio-dU



2-Thio-dT

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

 Monomers
For Instrument type Add

Expedite	E
MerMade	M

 Columns
For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

STABILITY NOTES

6-Thio-dG, 4-Thio-dT and 4-thio-dU are protected as the S-cyanoethyl ether which is stable during synthesis and readily removed by ammonium hydroxide. It is critical to add 50mM sodium hydrosulfide (NaSH) to the ammonium hydroxide used for deprotection. Especially if room temperature deprotection is carried out, this technique radically reduces the level of ammonolysis which would lead to undesired aminated bases. Moreover, it is also desirable to remove the cyanoethyl protecting group (1M DBU in acetonitrile, 2-5 h/RT) prior to the ammonium hydroxide cleavage and deprotection.

STRUCTURE/ACTIVITY RELATIONSHIP (CONT.)



STABILITY NOTE

Synthetic oligonucleotides containing 8-amino-dG must be cleaved and deprotected with ammonium hydroxide containing 0.25M 2-mercaptoethanol to avoid oxidative degradation of 8-amino-dG sites.

8-Amino-dA and 8-amino-dG are useful in triplex formation due to the presence of the additional amino groups.

2'-DeoxyXanthosine (dX) is a naturally occurring nucleoside that may be derived from oxidative deamination of 2'-deoxyGuanosine (dG). dX has a similar bonding pattern to thymidine and it may base pair with dA, with such purine-purine interactions causing duplex distortion. dX also featured in attempts to extend the genetic alphabet with a new base pair of dX and pyrimidine-2,4-diamine nucleoside. dX has also interested researchers in the field of DNA damage and repair since it is a product of nitric oxide-induced mutagenesis.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
8-Amino-dA-CE Phosphoramidite	10-1086-95	50 μ mole	177.50
	10-1086-90	100 μ mole	355.00
	10-1086-02	0.25g	975.00
8-Amino-dG-CE Phosphoramidite	10-1079-95	50 μ mole	177.50
	10-1079-90	100 μ mole	355.00
	10-1079-02	0.25g	975.00
2'-dX-CE Phosphoramidite	10-1537-95	50 μ mole	105.00
	10-1537-90	100 μ mole	200.00
	10-1537-02	0.25g	420.00




HALOGENATED NUCLEOSIDES

Brominated and iodinated nucleosides are used in crystallography studies of oligonucleotide structure. They are also photolabile and are used for cross-linking studies to probe the structure of protein-DNA complexes. Antibodies exist to Br-dU and oligonucleotides containing Br-dU can be used as probes.


Item	Catalog No.	Pack	Price(\$)
8-Br-dA-CE Phosphoramidite	10-1007-90	100 μ mole	115.00
	10-1007-02	0.25g	295.00
8-Br-dG-CE Phosphoramidite	10-1027-90	100 μ mole	105.00
	10-1027-02	0.25g	255.00
5-Br-dC-CE Phosphoramidite	10-1080-90	100 μ mole	60.00
	10-1080-02	0.25g	160.00
5-I-dC-CE Phosphoramidite	10-1081-90	100 μ mole	135.00
	10-1081-02	0.25g	355.00
5-Br-dU-CE Phosphoramidite	10-1090-90	100 μ mole	60.00
	10-1090-02	0.25g	160.00
5-I-dU-CE Phosphoramidite	10-1091-90	100 μ mole	60.00
	10-1091-02	0.25g	160.00
5-F-dU-CE Phosphoramidite	10-1092-90	100 μ mole	135.00
	10-1092-02	0.25g	355.00
5-Br-dU-CPG	20-2090-01	0.1g	50.00
	20-2090-41	Pack of 4	200.00
	20-2090-42	Pack of 4	120.00




8-Bromo-2'-deoxyAdenosine




8-Bromo-2'-deoxyGuanosine




5-Bromo-2'-deoxyCytidine




5-Iodo-2'-deoxyCytidine



5-Bromo-2'-deoxyUridine



5-Iodo-2'-deoxyUridine



5-Fluoro-2'-deoxyUridine

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

STABILITY NOTE

Oligonucleotides containing a bromo or iodo group are prepared conventionally with the exception that deprotection is carried out in ammonium hydroxide at room temperature for 24 hours. Under these conditions, degradation of the halogen group was less than 2%.


DEUTERATED NUCLEOSIDES

Perdeuteration and selective deuteration have been useful approaches for simplification of NMR spectra and for other structural studies of large biomolecules. Driven by the progress in multinuclear multidimensional NMR spectroscopy, deuteration of nucleic acids has especially found wide applications in the NMR studies of these complex molecules in solution. 8-Deutero-2'-deoxyGuanosine phosphoramidite will be of interest to our customers involved in NMR spectroscopy.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
8-D-dG-CE Phosphoramidite	10-1520-90	100 μ mole	90.00
	10-1520-02	0.25g	240.00

STABILITY NOTES

Synthetic oligonucleotides containing 8-D-dG must be cleaved and deprotected with 25% deuterated ammonium hydroxide for 40 hours at room temperature to minimize deuterium exchange.




8-Deutero-2'-deoxyGuanosine


DNA DAMAGE/REPAIR

Cellular DNA is constantly being damaged by oxidation and alkylation, by free radicals, and by ultraviolet and ionizing radiation. The body has therefore evolved a number of repair enzyme systems to excise and repair these lesions. The 8-oxo purine monomers allow investigation of the structure and activity of oligonucleotides containing an 8-oxo mutation which is formed naturally when DNA is subjected to oxidative conditions or ionizing radiation. 5,6-Dihydro pyrimidines are naturally occurring compounds that are structural components of alanine transfer RNA. Dihydrouracil and the hydroxy pyrimidines are major base damage products formed by exposure of DNA to ionizing radiation.


Item	Catalog No.	Pack	Price(\$)
8-Oxo-dA-CE Phosphoramidite	10-1008-90	100 μmole	135.00
	10-1008-02	0.25g	355.00
8-Oxo-dG-CE Phosphoramidite	10-1028-95	50 μmole	177.50
	10-1028-90	100 μmole	355.00
	10-1028-02	0.25g	975.00
5,6-Dihydro-dT-CE Phosphoramidite	10-1530-90	100 μmole	195.00
	10-1530-02	0.25g	595.00
5,6-Dihydro-dU-CE Phosphoramidite	10-1550-90	100 μmole	195.00
	10-1550-02	0.25g	595.00
5-OH-dC-CE Phosphoramidite	10-1063-90	100 μmole	275.00
	10-1063-02	0.25g	775.00
5-OH-dU-CE Phosphoramidite	10-1053-90	100 μmole	225.00
	10-1053-02	0.25g	675.00
5-Hydroxymethyl-dU-CE Phosphoramidite	10-1093-90	100 μmole	225.00
	10-1093-02	0.25g	675.00




8-oxo-2'-deoxyAdenosine




8-oxo-2'-deoxyGuanosine




5,6-Dihydro-dT




5,6-Dihydro-dU



5-OH-dC



5-OH-dU



5-Hydroxymethyl-dU

STABILITY NOTES

Synthetic oligonucleotides containing 8-oxo-dG must be cleaved and deprotected with ammonium hydroxide containing 0.25M 2-mercaptoethanol to avoid oxidative degradation of 8-oxo-dG sites.

Oligonucleotides synthesized using 5,6-dihydro-dU or 5,6-dihydro-dT and UltraMILD monomers can be cleaved using either concentrated ammonium hydroxide or 50 mM potassium carbonate in anhydrous methanol. Complete cleavage and deprotection can be accomplished at room temperature in 2-4 hours without damaging either the dihydro-dU or dihydro-dT bases.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

SEE ALSO

5-Hydroxymethyl-dC	p45
dX	p54

STABILITY NOTES

Synthetic oligonucleotides containing 8-amino-dG must be cleaved and deprotected with ammonium hydroxide containing 0.25M 2-mercaptoethanol to avoid oxidative degradation of 8-amino-dG sites.

Oligonucleotides synthesized using Thymidine Glycol and UltraMILD monomers can be cleaved using either concentrated ammonium hydroxide or 50 mM potassium carbonate in anhydrous methanol. Complete cleavage and deprotection can be accomplished at room temperature in 2-4 hours without damaging Thymidine Glycol base. The best method to remove the TBDMS groups was achieved using TEA.3HF at 40°C overnight.

REFERENCE

(1) K. Groebke, and C.J. Leumann, *Helv Chim Acta*, 1990, **73**, 608-617.

SEE ALSO


dSpacer p80
Pyrrolidine p59

DNA DAMAGE/REPAIR (CONT.)


8-Amino-G is formed along with 8-oxo-G as the major mutagenic lesions formed in DNA damage caused by 2-nitropropane. 2-Nitropropane is an industrial solvent and a component of paints, dyes and varnishes, and is also present in cigarette smoke. Thymine glycol (5,6-dihydroxy-5,6-dihydrothymine) is formed when thymine is subjected to oxidative stress, including ionizing radiation. Oxidation of the 5,6 double bond of Thymidine generates two chiral centers at C5 and C6. The cis-5R,6S form is generated as the predominant product along with the other diastereomer, the cis-5S,6R form. The presence of thymidine glycol in DNA has significant biological consequences and many organisms possess specific repair enzymes for the excision of this lesion.

Hydrolysis of nucleoside residues in DNA occurs to generate abasic sites. Most commonly, dA sites are hydrolyzed causing depurination and leading to abasic residues. For researchers trying to determine if their source of depurination in chemical synthesis of DNA is reagent, fluidics or protocol-based, we offer a depurination-resistant dA monomer. A new chemical method allows the generation of abasic sites in double and single stranded oligonucleotides using very mild specific conditions and with very low probability of side reactions. The original Abasic Phosphoramidite (10-1924) has been discontinued since it exhibits low coupling efficiency and the post-synthesis chemistry is fairly challenging. Abasic II Phosphoramidite¹ is the replacement for the preparation of a true abasic site. This product has the advantage of simplicity in that the silyl group is removed post-synthesis using aqueous acetic acid. dSpacer has also been used successfully as a mimic of the highly base-labile abasic site.


Item	Catalog No.	Pack	Price(\$)
8-Amino-dG-CE Phosphoramidite	10-1079-95	50 µmole	177.50
	10-1079-90	100 µmole	355.00
	10-1079-02	0.25g	975.00
Thymidine Glycol CE Phosphoramidite	10-1096-95	50 µmole	180.00
	10-1096-90	100 µmole	360.00
	10-1096-02	0.25g	975.00
Abasic II Phosphoramidite (dR Precursor)	10-1927-95	50 µmole	80.00
	10-1927-90	100 µmole	150.00
	10-1927-02	0.25g	475.00
dbf-dA-CE Phosphoramidite (Discontinued)	10-1500		




8-Amino-dG



Thymidine Glycol



dbf-dA



Abasic II Phosphoramidite



DNA DAMAGE/REPAIR (CONT.)

One of the major sources of DNA damage in all organisms is the UV component of sunlight. The predominant reaction induced by UV light on DNA is dimerization of adjacent pyrimidine bases leading to cyclobutane dimers (CPDs). The dimers formed in the most significant quantity are the *cis-syn* cyclobutane dimer of two thymine bases. Although formed routinely, these dimer products are efficiently excised and repaired enzymatically by nucleotide excision repair (NER) or the dimerization is reversed by photolase enzymes. A further mode of oxidative damage is radiation-induced damage of DNA, which has been shown to lead to bridged cyclonucleosides. The purines, cyclo-dA and cyclo-dG, are predominantly formed, although the cyclo pyrimidines have also been detected. Cyclo-dA is doubly intriguing since it contains both damaged base and damaged sugar residues and, as such, should have a considerable biological impact. In a manner analogous to thymine dimer, cyclo purines cause significant distortion of the regular DNA helix and these lesions are repaired not by base excision repair (BER) but by NER.


Item	Catalog No.	Pack	Price(\$)
<i>Cis-syn</i> Thymine Dimer Phosphoramidite	11-1330-95	50 μ mole	2100.00
	11-1330-90	100 μ mole	4200.00
	11-1330-02	0.25g	10200.00
5',8-Cyclo-dA CE Phosphoramidite	10-1098-95	50 μ mole	950.00
	10-1098-90	100 μ mole	1850.00
	10-1098-02	0.25g	5350.00
5',8-Cyclo-dG CE Phosphoramidite	10-1598-95	50 μ mole	1250.00
	10-1598-90	100 μ mole	2450.00

Base excision repair (BER) is one of the most studied repair mechanisms. In this pathway, DNA glycosylases recognize the damaged bases and catalyze their excision through hydrolysis of the N-glycosidic bond. Attempts to understand the structural basis for DNA damage recognition by DNA glycosylases have been hampered by the short-lived association of these enzymes with their DNA substrates. To overcome this problem, the Verdine group at Harvard synthesized a pyrrolidine analog that mimics the charged transition state of the enzyme-substrate complex. When incorporated into double-stranded DNA, they found the pyrrolidine analog (PYR), introduced as the Pyrrolidine-CE Phosphoramidite, forms an extremely stable complex with the DNA glycosylase AlkA, exhibiting a dissociation constant in the μ M range and potently inhibited the reaction catalyzed by the enzyme.


Item	Catalog No.	Pack	Price(\$)
Pyrrolidine-CE Phosphoramidite (PYR)	10-1915-95	50 μ mole	190.00
	10-1915-90	100 μ mole	380.00
	10-1915-02	0.25g	1085.00

*Cis-syn* Thymine Dimer

5',8-Cyclo-dA



5',8-Cyclo-dG




Pyrrolidine

INSTRUMENT TYPES

For these very expensive phosphoramidites, an ABI septum vial is the standard vial. Add E to the catalog no. for an Expedite vial or V to the catalog no. for an Expedite V vial.

BIOCOMPATIBLE TRIAZOLE LINKAGE



SEE ALSO

5'-I-dT in Click Chemistry p85
Click Chemistry p84

REFERENCES - CLICK LIGATION

- (1) A.H. El-Sagheer, A.P. Sanzone, R. Gao, A. Tavassoli, and T. Brown, *Proc Natl Acad Sci U S A*, 2011, **108**, 11338-43.
- (2) A.H. el-Sagheer, and T. Brown, *Chem Commun (Camb)*, 2011, **47**, 12057-8.
- (3) A.P. Sanzone, A.H. El-Sagheer, T. Brown, and A. Tavassoli, *Nucleic Acids Res*, 2012.
- (4) A. Dallmann, et al., *Chemistry*, 2011, **17**, 14714-7.
- (5) A.H. El-Sagheer, and T. Brown, *Proc Natl Acad Sci U S A*, 2010, **107**, 15329-34.

REFERENCES - MicroRNA LABELLING

- (1) H. Vogel, and C. Richert, *ChemBioChem*, 2012, **13**, 1474-82.
- (2) R. Eisenhuth, and C. Richert, *Journal of Organic Chemistry*, 2008, **74**, 26-37.
- (3) E. Kervio, A. Hochgesand, U.E. Steiner, and C. Richert, *Proc Natl Acad Sci U S A*, 2010, **107**, 12074-9.

CLICK DNA AND RNA LIGATION


Ligation of an oligo containing a 5'-azide with an oligo containing a 3'-propargyl group using Click Chemistry leads to a triazole linkage that has been shown to have *in vivo* biocompatibility. This technique has been used to synthesize DNA constructs up to 300 bases in length. When the resultant triazole linkage was placed in a PCR template, various polymerases were able to copy the sequence correctly. The linkage has also been shown to be compatible with transcription and rolling circle amplification, as well as gene expression in *E. coli*. In the RNA world, a hammerhead ribozyme containing the triazole linkage at the substrate cleavage site has been shown to retain its activity. A large variety of applications is envisaged for this biocompatible chemical ligation. Support for this technology is offered with the help of Tom Brown's group at the University of Southampton.

Item	Catalog No.	Pack	Price(\$)
3'-Propargyl-5-Me-dC CPG	20-2982-01	0.1g	180.00
	20-2982-10	1.0g	1500.00
1 μ mole columns	20-2982-41	Pack of 4	300.00
0.2 μ mole columns	20-2982-42	Pack of 4	150.00
10 μ mole column (ABI)	20-2982-13	Pack of 1	750.00
15 μ mole column (Expedite)	20-2982-14	Pack of 1	1125.00


5'-LABELLING OF MicroRNAs

Several methods have been developed for the detection of miRNAs, however, few allow the simultaneous detection of multiple miRNAs. To overcome this analytical deficiency, the Richert group at the University of Stuttgart has recently developed an ingenious method to selectively detect miRNAs on microarrays without interference from endogenous pre-mRNAs, mRNAs and other RNA species. In this method, a short oligonucleotide containing 3'-amino-dT and a 5' reporter molecule is chemically ligated to the microRNA in a one-step procedure by *in situ* activation of the microRNA. This is specifically achieved by taking advantage of the fact that miRNAs, unlike other RNAs, are 5'-phosphorylated. The reaction is template-directed (and thus sequence specific) and can be performed together with enzymatic 3'-extension/labelling, either in solution or on a support. The short DNA labelling strand may feature one of a variety of different labels, such as a biotin group or a fluorophore.

Item	Catalog No.	Pack	Price(\$)
3'-Amino-dT CPG	20-2981-01	0.1g	120.00
	20-2981-10	1.0g	995.00
1 μ mole columns	20-2981-41	Pack of 4	200.00
0.2 μ mole columns	20-2981-42	Pack of 4	120.00
10 μ mole column (ABI)	20-2981-13	Pack of 1	500.00
15 μ mole column (Expedite)	20-2981-14	Pack of 1	750.00



3'-Propargyl-5-Me-dC CPG









3'-Amino-dT CPG

2'-5' LINKED OLIGONUCLEOTIDES

Cellular DNA and RNA are made up of ribo- and 2'-deoxyribonucleic acids linked together via 3'-5' phosphodiester linkages and by far comprise the bulk of polynucleic acids found in cells. Much less common are oligonucleotides which have 2'-5' linkages. However, a unique feature of 2'-5' linked oligonucleotides is their ability to bind selectively to complementary RNA. These features suggest a number of interesting uses for 2'-5' linked oligos such as their use as RNA specific probes or in antisense oligos. Recently, oligos have been synthesized using 3'-deoxy-2'-phosphoramidites and 2'-deoxy-3'-phosphoramidites to produce chimeras with 2'-5' linked ends and 3'-5' linked central regions. It was found that 2'-5' phosphorothioate oligos: 1) bind selectively to complementary RNA with the same affinity as phosphodiester oligos; 2) exhibit much less nonspecific binding to cellular proteins; 3) do not activate RNase H. A 3'-deoxynucleoside at the 3'-terminus of an otherwise normal oligonucleotide effectively blocks polymerase extension.

Item	Catalog No.	Pack	Price(\$)
3'-dA-CE Phosphoramidite	10-1004-95	50 μ mole	177.50
	10-1004-90	100 μ mole	355.00
	10-1004-02	0.25g	975.00
3'-dC-CE Phosphoramidite	10-1064-95	50 μ mole	177.50
	10-1064-90	100 μ mole	355.00
	10-1064-02	0.25g	975.00
3'-dG-CE Phosphoramidite	10-1074-95	50 μ mole	177.50
	10-1074-90	100 μ mole	355.00
	10-1074-02	0.25g	975.00
3'-dT-CE Phosphoramidite	10-1084-95	50 μ mole	177.50
	10-1084-90	100 μ mole	355.00
	10-1084-02	0.25g	975.00
3'-dA-CPG 1 μ mole columns 0.2 μ mole columns	20-2004-01	0.1g	300.00
	20-2104-41	Pack of 4	600.00
	20-2104-42	Pack of 4	200.00
3'-dC-CPG 1 μ mole columns 0.2 μ mole columns	20-2064-01	0.1g	300.00
	20-2164-41	Pack of 4	600.00
	20-2164-42	Pack of 4	200.00



OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers
For Instrument type Add

Expedite	E
MerMade	M

Columns
For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

SEE ALSO

3'-deoxynucleoside CPG p50

2'-5' LINKED OLIGONUCLEOTIDES (CONT.)

Item	Catalog No.	Pack	Price(\$)
3'-dG-CPG	20-2074-01	0.1g	300.00
1 μmole columns	20-2174-41	Pack of 4	600.00
0.2 μmole columns	20-2174-42	Pack of 4	200.00
3'-dT-CPG	20-2084-01	0.1g	300.00
1 μmole columns	20-2184-41	Pack of 4	600.00
0.2 μmole columns	20-2184-42	Pack of 4	200.00

MUTAGENESIS


Cellular polynucleotides are alkylated by endogenous components, such as S-adenosylmethionine, or after reacting with two general classes of environmental and laboratory chemicals. SN1 chemical agents include alkyl nitrosourea and N-alkyl-N-nitro-N-nitrosoguanidine that react with the N7 position of guanine, N3 of adenine, O6 of guanine, O2 or O4 of pyrimidines, and the non-phosphodiester oxygen atoms of the phosphate backbone. In contrast, SN2 chemical agents such as methyl methanesulfonate and dimethyl sulfate react primarily with the N1 position of adenine (1-Methyl-2'-deoxyadenosine) and N3 of cytosine. To avoid chain branching during synthesis when using DCl as activator, N6-Me-dA is offered with acetyl protection.

SEE ALSO

N6-Me-dA

p43





Item	Catalog No.	Pack	Price(\$)
06-Me-dG-CE Phosphoramidite	10-1070-90	100 μmole	105.00
	10-1070-02	0.25g	255.00
N6-Me-dA-CE Phosphoramidite	10-1003-90	100 μmole	162.50
	10-1003-02	0.25g	495.00
N6-Ac-N6-Me-dA-CE Phosphoramidite	10-1503-90	100 μmole	162.50
	10-1503-02	0.25g	495.00
04-Me-dT-CE Phosphoramidite	10-1032-90	100 μmole	135.00
	10-1032-02	0.25g	355.00
1-Me-dA-CE Phosphoramidite	10-1501-95	50 μmole	125.00
	10-1501-90	100 μmole	250.00
	10-1501-02	0.25g	750.00



CONVERTIBLE NUCLEOSIDES

The convertible nucleoside strategy is one of the most versatile methods for producing modifications in bases to examine their effects on DNA structure and activity. In some cases, with versatility comes difficulty in that the convertible base is modified after oligonucleotide synthesis. The chemistry is sometimes complex and base composition analysis of the final oligonucleotide is required to verify structure. The convertible dU monomer can be used to introduce a variety of modifications at the convertible position, including N, O and S modifications. Convertible F-dC is by far the simplest approach to the preparation of oligonucleotides containing F-dC - normal ammonium hydroxide treatment effects the conversion to F-dC. Convertible dA has been used to prepare oligonucleotides containing multiple points for attachment to solid supports. In this way, high capacity affinity supports for the purification of DNA binding proteins have been prepared. 2-F-dI is a convertible nucleoside for the preparation of 2'-dG derivatives following the displacement of the 2-fluorine by primary amines.

Item	Catalog No.	Pack	Price(\$)
TMP-F-dU-CE Phosphoramidite (Convertible F-dC)	10-1016-90	100 μ mole	195.00
	10-1016-02	0.25g	495.00
O6-Phenyl-dI-CE Phosphoramidite (Convertible dA)	10-1042-90	100 μ mole	135.00
	10-1042-02	0.25g	355.00
O4-Triazolyl-dU-CE Phosphoramidite (Convertible dU)	10-1051-90	100 μ mole	135.00
	10-1051-02	0.25g	355.00
2-F-dI-CE Phosphoramidite (Convertible dG)	10-1082-95	50 μ mole	180.00
	10-1082-90	100 μ mole	360.00
	10-1082-02	0.25g	975.00



OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers For Instrument type Add

Expedite	E
MerMade	M

Columns For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

ABBREVIATION

TMP = 2,4,6-trimethylphenyl

SEE ALSO

2-Aminopurine	p53
AP-dC (G-Clamp)	p42
UltraMild Chemistry	p68
Pyrrolo-C	p121
Pyrrolo-CTP	p124

INTELLECTUAL PROPERTY

Pyrrolo-dC is a joint development project of Berry & Associates, Inc. and Glen Research Corporation. Pyrrolo-dC is covered by US Patent No.: 7,144,995.

SPECTRAL PROPERTIES

The spectral properties of pyrrolo-dC, coupled with its unique base-pairing ability, make this fluorescent analog extremely valuable in probing DNA structure. When the pyrrolo-dC is base-paired, its fluorescence is significantly quenched through what is most likely base stacking or dG interactions. The quantum yield of fluorescence for pyrrolo-dC is quite sensitive to its hybridization state, making it ideally suited for probing the dynamic structure of DNA.

QY	λ	ϵ
		(L/mol.cm)
single-stranded	0.07	260nm 4000
		347nm 3700
double-stranded	0.02	

(QY determined relative to quinine sulfate in 0.5M H₂SO₄)

REFERENCES


1. D.A. Berry, et al., *Tetrahedron Lett*, 2004, **45**, 2457-2461.
2. The Glen Report, 2007, **19**, 8-9.
3. P. Sandin, et al., *Nucleic Acids Res.*, 2008, **36**, 157-167.
4. P. Sandin, et al., *Nucleic Acids Res.*, 2005, **33**, 5019-5025.
5. K.C. Engman, et al., *Nucleic Acids Res.*, 2004, **32**, 5087-5095.

FLUORESCENT NUCLEOSIDES


Etheno-dA is a fluorescent nucleoside which is especially useful in observing the transition between DNA structural types. It is quite base labile and should be deprotected with ammonium hydroxide at room temperature for 24 hours. Alternatively, UltraMild chemistry can be used. 2-Aminopurine and AP-dC (G-Clamp) are also useful fluorescent nucleosides.

Pyrrolo-dC is a fluorescent deoxycytidine analog that is an ideal probe of DNA structure and dynamics.^{1,2} It base-pairs as a normal dC nucleotide. An oligo fully substituted with pyrrolo-dC has the same T_m as the control dC oligo with the same specificity for dG. Its small size does not perturb the structure of the DNA helix and it is well tolerated by a number of DNA and RNA polymerases. It is highly fluorescent and its excitation and emission are well to the red of most fluorescent nucleotide analogs, which eliminates or reduces background fluorescence from proteins. Pyrrolo-dCTP has potential uses in biological assay development.


Item	Catalog No.	Pack	Price(\$)
Etheno-dA-CE Phosphoramidite	10-1006-90	100 μ mole	105.00
	10-1006-02	0.25g	255.00
Pyrrolo-dC-CE Phosphoramidite	10-1017-95	50 μ mole	110.00
	10-1017-90	100 μ mole	220.00
	10-1017-02	0.25g	675.00
Pyrrolo-dCTP (10 mM)	81-1017-01	100 μ L	150.00



Etheno-2'-deoxyAdenosine



Pyrrolo-dC






Pyrrolo-dCTP

FLUORESCENT NUCLEOSIDES (CONT.)

When a benzylcarbamoyl analogue of AP-dC (G-Clamp) was synthesized, it was found that, when incorporated into an oligo, it exhibited similar fluorescence to AP-dC. However, when base-paired against the 8-oxo-dG, its fluorescence was severely quenched. Rather remarkably, however, when base paired with dG or any of the other bases, A, C or T, there was no change in fluorescence – making it a specific probe for 8-oxo-dG.

By attaching pyrene or perylene to the 5 position of deoxyuridine through a triple bond, the fluorophore is electronically coupled to the deoxyuridine base. This electronic coupling of the base and the fluorophore makes the fluorescence sensitive to the base pairing of the dU portion of the molecule, allowing the discrimination between perfect and one base mismatched targets.

Item	Catalog No.	Pack	Price(\$)
8-Oxo-G Clamp-CE Phosphoramidite (Discontinued)	10-1515		
Pyrene-dU-CE Phosphoramidite	10-1590-95	50 μ mole	105.00
	10-1590-90	100 μ mole	210.00
	10-1590-02	0.25g	550.00
Perylene-dU-CE Phosphoramidite	10-1591-95	50 μ mole	150.00
	10-1591-90	100 μ mole	300.00
	10-1591-02	0.25g	720.00



SEE ALSO

8-Oxo-dG	p57
AP-dC (G-Clamp)	p42

SPECTRAL PROPERTIES

	Absorbance Maximum	Emission Maximum
Pyrene-dU	402nm	472nm
Perylene-dU	473nm	490nm

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

THERAPEUTIC NUCLEOSIDES

Cytosine Arabanoside (Ara-C) is an anti-viral drug which has achieved limited use. Its effect on DNA structure and activity can be investigated by incorporating it into synthetic oligonucleotides.

Zebularine (pyrimidin-2-one ribonucleoside) is a cytidine analogue that acts as a DNA demethylase inhibitor, as well as a cytidine deaminase inhibitor. This structure is very active biologically and Zebularine is now used as a potent anti-cancer drug. A 2'-deoxynucleoside analogue of Zebularine, 5-methyl-pyrimidin-2-one, 2'-deoxynucleoside, has been used to probe the initiation of the cellular DNA repair process by making use of its mildly fluorescent properties. This combination of biological activity and fluorescence properties would make 5-Me-2'-deoxyZebularine a strong addition to our array of nucleoside analogues.


Cytosine-5-methyltransferases are found in everything from archaebacteria to mammals and when the regulation of cytosine-5-methyltransferases goes awry, cancer can result. The mechanism of action for this family of enzymes involves attack of a cysteine thiol group on the C6 position of cytosine, leading to a transient dihydrocytosine intermediate, which then facilitates the nucleophilic attack by C5 on the activated methyl group of the S-adenosyl-L-methionine cofactor. As with many enzymes, the intermediate can be trapped using a suicide substrate and 5-fluoro-cytosine has been used extensively in this role. An alternate strategy is to use a transition-state mimic that binds to the active site with high affinity. An excellent candidate was found in 5-aza-5,6-dihydrocytosine. Despite not being covalently bound to the enzyme, it was found^{1,2} to be a more potent inhibitor of cytosine-5-methyltransferases than 5-fluoro-cytosine. 5-Aza-5,6-dihydro-dC is compatible with standard oligonucleotide synthesis and deprotection conditions and is an excellent tool for use in methyltransferase research.

Item	Catalog No.	Pack	Price(\$)
Ara-C-CE Phosphoramidite (Discontinued)	10-4010		
5-Me-2'-deoxyZebularine-CE Phosphoramidite	10-1061-95	50 μ mole	200.00
	10-1061-90	100 μ mole	400.00
	10-1061-02	0.25g	975.00
5-Aza-5,6-dihydro-dC-CE Phosphoramidite	10-1511-95	50 μ mole	180.00
	10-1511-90	100 μ mole	360.00
	10-1511-02	0.25g	1120.00


LARGE SCALE SYNTHESIS

The most common side reaction during deprotection of oligonucleotides on a large scale is the alkylation of dT residues by acrylonitrile, formed by β -elimination of the cyanoethyl phosphate protecting groups, to generate N3-cyanoethyl-dT.


Item	Catalog No.	Pack	Price(\$)
N3-Cyanoethyl-dT	10-1531-90	100 μ mole	200.00
	10-1531-02	0.25g	600.00




Ara-C



5-Me-2'-deoxyZebularine



5-Aza-5,6-Dihydro-dC



N3-Cyanoethyl-dT

REFERENCES

- (1) G. Sheiknejad, et al., *J Mol Biol*, 1999, **285**, 2021-2034.
- (2) V.E. Marquez, et al., *Antisense Nucleic Acid Drug D*, 1999, **9**, 415-421.

SEE ALSO

Convertible F-dC p63

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M


(Please inquire for availability of vials and columns for other instrument types.)

ULTRAMILD CE PHOSPHORAMIDITES


The synthesis of labelled oligonucleotides has become a standard procedure in many laboratories and many labelling reagents, e.g., biotin and fluorescein, are now available as β -cyanoethyl (CE) phosphoramidites. Labels which are currently available as CE phosphoramidites have one common property - they must be stable to the strongly alkaline conditions required for removal of the base protecting groups. This property is lacking in several interesting dyes and labels. We sought an alternative protecting scheme for the normal CE phosphoramidites which would allow UltraMILD deprotection and would not react with a wider variety of tags and labels. A set of monomers using phenoxyacetyl (Pac) protected dA and 4-isopropyl-phenoxyacetyl (iPr-Pac) protected dG, along with acetyl protected dC, met the desired criteria for UltraMILD deprotection.

We recommend the use of phenoxyacetic anhydride (Pac₂O) in Cap A. This modification removes the possibility of exchange of the iPr-Pac protecting group on the dG with acetate from the acetic anhydride capping mix. Cleavage and deprotection can be carried out in 2 hours at room temperature with ammonium hydroxide or 4 hours with 0.05M potassium carbonate in methanol.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Pac-dA-CE Phosphoramidite	10-1601-02	0.25g	15.00
	10-1601-05	0.5g	30.00
	10-1601-10	1.0g	60.00
Ac-dC-CE Phosphoramidite	10-1015-02	0.25g	12.50
	10-1015-05	0.5g	25.00
	10-1015-10	1.0g	50.00
iPr-Pac-dG-CE Phosphoramidite	10-1621-02	0.25g	15.00
	10-1621-05	0.5g	30.00
	10-1621-10	1.0g	60.00



Pac-dA



Ac-dC



iPr-Pac-dG

ULTRAMILD SUPPORTS

<i>Item</i>	<i>Catalog No.</i> Pac-dA	<i>Catalog No.</i> Ac-dC	<i>Catalog No.</i> iPr-Pac-dG	<i>Pack</i>	<i>Price(\$)</i>
UltraMild CPG (Bulk)	20-2601-01	Listed	20-2621-01	0.1g	18.00
	20-2601-02	on	20-2621-02	0.25g	40.00
	20-2601-10	Page 8	20-2621-10	1.0g	150.00
ABI Columns	20-2701-45	20-2115-45	20-2721-45	4X40nm	40.00
	20-2701-42	20-2115-42	20-2721-42	4X0.2µm	40.00
	20-2701-41	20-2115-41	20-2721-41	4X1µm	60.00
	20-2701-13	20-2115-13	20-2721-13	10µm	100.00
Expedite Columns	20-2801-45	20-2215-45	20-2821-45	4X40nm	40.00
	20-2801-42	20-2215-42	20-2821-42	4X0.2µm	40.00
	20-2801-41	20-2215-41	20-2821-41	4X1µm	60.00
	20-2801-14	20-2215-14	20-2821-14	15µm	150.00

ULTRAMILD SOLVENTS/REAGENTS

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Cap Mix A</i>			
THF/Pyridine/Pac ₂ O (Applied Biosystems)	40-4210-52	200mL	140.00
	40-4210-57	450mL	300.00
THF/Pac ₂ O (Expedite)	40-4212-52	200mL	140.00
	40-4212-57	450mL	300.00
<i>Deprotection Solution</i>			
0.05M Potassium Carbonate in Methanol	60-4600-30	30mL	30.00

SEE ALSO

Universal Support III p24

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

TERMINUS MODIFIERS

INTELLECTUAL PROPERTY

5'-Carboxy-Modifier C10 is offered for sale under license from TriLink BioTechnologies, Inc. It is intended for research and development purposes only, and may not be used for commercial, clinical, diagnostic or any other use. It is covered under US Patent No. 6,320,041.

SEE ALSO

PC modifiers

p82

ABBREVIATIONS


CNEt = Cyanoethyl
 CPG = Controlled Pore Glass
 DMT = 4,4'-Dimethoxytrityl
 Fmoc = Fluorenylmethoxycarbonyl
 iPr = Isopropyl
 MMT = 4-Monomethoxytrityl
 T = Trityl
 TFA = Trifluoroacetyl

Glen Research 5'-Modifiers are designed for use in DNA synthesizers to functionalize the 5'-terminus of the target oligonucleotide. The 5'-Amino-Modifiers are available with a variety of chain lengths to fit exactly the desired application.


The DMS(O)MT-protected amino group is easier to deprotect compared to the MMT-protected one. The sulfoxy derivative survives conditions of oligonucleotide synthesis and can either be cleaved with standard deblock solution, or left intact for HPLC purification. At the same time, the DMS(O)MT group is fully compatible with cartridge purification. When detritylation on a cartridge is carried out, the DMS(O)MT+, which is more stable than MMT+, does not reattach itself to an amine. We now offer 5'-DMS(O)MT-Amino-Modifier C6 utilizing this new trityl based protecting group.

5'-Amino-Modifier TEG, a hydrophilic triethylene glycol ethylamine derivative, is 12 atoms in length and fully soluble in aqueous media.


Item	Catalog No.	Pack	Price (\$)
5'-Amino-Modifier C3-TFA	10-1923-90	100 µmole	50.00
	10-1923-02	0.25g	175.00
5'-Amino-Modifier C6	10-1906-90	100 µmole	60.00
	10-1906-02	0.25g	200.00
5'-Amino-Modifier C6-TFA	10-1916-90	100 µmole	30.00
	10-1916-02	0.25g	100.00
5'-Amino-Modifier C12	10-1912-90	100 µmole	90.00
	10-1912-02	0.25g	300.00
5'-Amino-Modifier 5	10-1905-90	100 µmole	60.00
	10-1905-02	0.25g	200.00
5'-DMS(O)MT-Amino-Modifier C6	10-1907-90	100 µmole	60.00
	10-1907-02	0.25g	200.00
5'-Amino-Modifier TEG	10-1917-90	100 µmole	115.00
	10-1917-02	0.25g	500.00




5'-Amino-Modifier C3-TFA




5'-Amino-Modifier C6




5'-Amino-Modifier C6-TFA




5'-Amino-Modifier C12



5'-Amino-Modifier 5



5'-Amino-Modifier TEG



5'-DMS(O)MT-Amino-Modifier C6

TERMINUS MODIFIERS (CONT.)


Our more recent 5'-amino modifiers, protected by a novel phthalic acid diamide (PDA) protecting group, are stable solids. In contrast to the TFA protected amino modifiers, which are viscous oils, the analogous PDA protected compounds are granular powders. This important property of these compounds allows straightforward handling, storage and aliquoting and leads to a significant increase in stability.

Deprotection with methylamine in gas phase or aqueous solution or AMA leads to fast and complete removal of the PDA protecting group. However, ammonium hydroxide will not drive the equilibrium reaction to completion and only partial deprotection occurs - overnight deprotection with ammonium hydroxide will yield around 80% active amine.


We are offering three PDA Amino-Modifiers:

- 5'-Amino-Modifier C6-PDA
- Hydrophobic 5'-Amino-Modifier C12-PDA
- Hydrophilic 5'-Amino-Modifier-TEG-PDA


Item	Catalog No.	Pack	Price (\$)
5'-Amino-Modifier C6-PDA	10-1947-90	100 μ mole	30.00
	10-1947-02	0.25g	100.00
5'-Amino-Modifier C12-PDA	10-1948-90	100 μ mole	65.00
	10-1948-02	0.25g	240.00
5'-Amino-Modifier-TEG-PDA	10-1949-90	100 μ mole	105.00
	10-1949-02	0.25g	420.00



5'-Amino-Modifier C6-PDA



5'-Amino-Modifier C12-PDA



5'-Amino-Modifier-TEG-PDA

INTELLECTUAL PROPERTY

PDA amino-modifiers were developed by Stefan Pitsch and ReseaChem GmbH (S. Berger), Patent pending.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

TERMINUS MODIFIERS (CONT.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

INTELLECTUAL PROPERTY


These products or portions thereof are subject to proprietary rights of FRIZ Biochem Gesellschaft für Bioanalytik mbH. US Patent No. 7,601,848 and European Patent No. EP 1 626 952 DTPA and derived products (Chemical Products) are for research purposes only, and may not be used for commercial, clinical, diagnostic or any other use. All oligo synthesis organizations are constrained to sell oligos containing the Chemical Products for research purposes only and any other use requires a license from FRIZ Biochem (www.frizbiochem.com). From Oct. 1st 2014 even all oligo synthesis organizations require a license from FRIZ to sell oligos containing the Chemical Products.

5'-Maleimide Modifier


Phosphoramidite is protected by a patent application and is offered by Glen Research under a non-exclusive license agreement from the University of Barcelona.

The disulfide thiol modifier may be used for introducing 3'- or 5'-thiol linkages. Dithiol Phosphoramidite (DTPA) is a disulfide-containing modifier designed to functionalize synthetic DNA or RNA with multiple thiol groups and can be incorporated at any position of the oligonucleotide. Each DTPA addition leads to two thiol groups. This modifier was designed for optimal tethering of oligonucleotides to a gold surface but it can also be used for multiple reactions with maleimides and other thiol-specific derivatives. 5'-Carboxy-Modifier C10 is a unique linker designed to be added at the terminus of an oligonucleotide synthesis. It generates an activated carboxylic acid N-hydroxysuccinimide (NHS) ester suitable for immediate conjugation on the synthesis column with molecules containing a primary amine, resulting in a stable amide linkage. An alternative carboxylate protecting group is the 2-chlorotrityl group, which is simply removed using the standard deblock cycle to generate a free carboxyl group on an otherwise fully protected oligonucleotide. The 2-chlorotrityl group is also removed during oligo deprotection with ammonium hydroxide or AMA and is incompatible with RP purification techniques. PC Amino-Modifier is a photocleavable C6 amino-modifier, part of our line of photocleavable (PC) modifiers. 5'-AminoOxy-Modifier 11 is based on a tetraethylene glycol linkage for improved solubility and for reducing the potential negative impact on hybridization of the oligo. The oxime formed from the reaction of alkyloxyamines with aldehydes creates a stable covalent bond. In comparison, the imine formed by the conjugation of primary amines with aldehydes is not stable to acidic or basic conditions and requires subsequent reduction with borohydride to form stable amine conjugates. 5'-Maleimide Modifier Phosphoramidite, developed at the University of Barcelona, incorporates a maleimide cycloadduct that is stable to ammonium hydroxide at room temperature. This phosphoramidite can be incorporated into DNA and RNA with both phosphate and phosphorothioate linkages. A retro-Diels-Alder reaction deprotects the maleimide immediately prior to conjugation.


Item	Catalog No.	Pack	Price (\$)
5'-Thiol-Modifier C6	10-1926-90 10-1926-02	100 µmole 0.25g	60.00 200.00
Thiol-Modifier C6 S-S	10-1936-90 10-1936-02	100 µmole 0.25g	150.00 360.00
Dithiol Phosphoramidite (DTPA) (Discontinued. Contact Glen Research Technical Support for more information.)	10-1937-90		
PC Amino-Modifier Phosphoramidite	10-4906-90 10-4906-02	100 µmole 0.25g	135.00 395.00
5'-Carboxy-Modifier C10	10-1935-90 10-1935-02	100 µmole 0.25g	50.00 200.00
5'-Carboxy-Modifier C5	10-1945-90 10-1945-02	100 µmole 0.25g	95.00 330.00
5'-AminoOxy-Modifier 11	10-1919-95 10-1919-90 10-1919-02	50 µmole 100 µmole 0.25g	140.00 265.00 895.00
5'-Maleimide-Modifier Phosphoramidite	10-1938-90 10-1938-02	100 µmole 0.25g	70.00 335.00




5'-Thiol-Modifier C6




Thiol-Modifier C6 S-S




DTPA




PC Amino-Modifier




5'-Carboxy-Modifier C10



5'-Carboxy-Modifier C5



5'-AminoOxy-Modifier 11



5'-Maleimide-Modifier


SEQUENCE MODIFIERS

Sequence Modifiers are designed for use in automated synthesis. The carboxy-dT is hydrolyzed during deprotection and can be coupled directly to a molecule containing a primary amino group by a standard peptide coupling or via the intermediate N-hydroxysuccinimide (NHS) ester. Amino-Modifier dA, Amino-Modifier dC, N2-Amino-Modifier dG and both Amino-Modifier dT products can be added in place of a dA, dC, dG and dT residue, respectively, during oligonucleotide synthesis. Corresponding Amino-Modifier supports can replace their respective deoxynucleoside supports. After deprotection, the primary amine on the C6 analogues is separated from the oligonucleotide by a spacer arm with a total of 7–10 atoms and can be labelled or attached to an enzyme. The C2 analogue is more suitable for the attachment of molecules designed to react with the oligonucleotide.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Amino-Modifier C6 dA	10-1089-90	100 μ mole	205.00
	10-1089-02	0.25g	455.00
Amino-Modifier C6 dC	10-1019-90	100 μ mole	225.00
	10-1019-02	0.25g	450.00
N2-Amino-Modifier C6 dG	10-1529-95	50 μ mole	240.00
	10-1529-90	100 μ mole	480.00
	10-1529-02	0.25g	1100.00
Carboxy-dT	10-1035-90	100 μ mole	180.00
	10-1035-02	0.25g	360.00
Amino-Modifier C2 dT	10-1037-90	100 μ mole	180.00
	10-1037-02	0.25g	360.00
	10-1037-05	0.5g	720.00
Amino-Modifier C6 dT	10-1039-90	100 μ mole	180.00
	10-1039-02	0.25g	360.00
	10-1039-05	0.5g	720.00

SEE ALSO

Amino-Modifier supports p75



SEQUENCE MODIFIERS (CONT.)

SEE ALSO

Carboxy-Modifier C10 p72
Carboxy-Modifier C5 p72

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers
For Instrument type Add

Expedite E
MerMade M

Columns
For Instrument type Add


Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)


Our repertoire of NHS ester derivatives has been expanded to include the NHS-Carboxy-dT-CE Phosphoramidite. By making a dT analog of the Carboxy-Modifier C10, it is possible to label one or multiple sites within an oligonucleotide. This opens up the possibility to label any number of different dyes or molecules within an oligonucleotide when the phosphoramidite is unavailable. Doing so is straightforward and may be done manually off the synthesizer or even in a fully-automated manner on the DNA synthesizer.

We have never found conditions which allow the TFA group to be removed from an amino-modifier while the oligonucleotide remains attached to the support. We are able to solve this problem by using a 9-fluorenylmethoxycarbonyl (Fmoc) protecting group. The Fmoc group is removed using a two step procedure, the first to remove the cyanoethyl protection groups and flush the formed acrylonitrile from the synthesis column using 1% diisopropylamine in acetonitrile, and the second to remove the Fmoc group using 10% piperidine in DMF. The amino group so formed on the column can be reacted with a variety of activated esters. We offer Fmoc-Amino-Modifier C6 dT Phosphoramidite as a nucleosidic option and Amino-Modifier Serinol Phosphoramidite as a non-nucleosidic alternative. We also offer S-Bz-Thiol-Modifier C6-dT to join the ranks of thiol-modifiers for oligonucleotide synthesis. Thiol-Modifier C6-dT can be added as usual at the desired locations within a sequence.


Item	Catalog No.	Pack	Price (\$)
NHS-Carboxy-dT	10-1535-90	100 µmole	210.00
	10-1535-02	0.25g	550.00
Fmoc-Amino-Modifier C6 dT	10-1536-90	100 µmole	180.00
	10-1536-02	0.25g	360.00
S-Bz-Thiol-Modifier C6-dT	10-1538-95	50 µmole	130.00
	10-1538-90	100 µmole	245.00
	10-1538-02	0.25g	550.00
Amino-Modifier Serinol Phosphoramidite	10-1997-95	50 µmole	125.00
	10-1997-90	100 µmole	225.00
	10-1997-02	0.25g	595.00




NHS-Carboxy-dT



Fmoc-Amino-Modifier C6 dT



S-Bz-Thiol-Modifier C6-dT




Amino-Modifier Serinol Phosphoramidite


3'-MODIFIERS

3'-Amino-Modifier CPGs, containing amino groups protected with the base-labile Fmoc group, are designed to functionalize the 3'-terminus of the target oligonucleotide by the introduction of a primary amine. In an alternative approach, the nitrogen destined to become the 3'-amino group is included in a phthalimide (PT) group which is attached to the support through an amide group attached to the aromatic ring. This simple linkage is very stable to all conditions of oligonucleotide synthesis and contains no chiral center. Using an extended ammonium hydroxide treatment (55°C for 17 hours), the cleavage of the amine from the phthalimide is accomplished along with the deprotection of the oligonucleotide. ABI-style columns are supplied unless otherwise requested.


<i>Item</i>	<i>Cat. No.</i>	<i>Pack</i>	<i>Price (\$)</i>
3'-Amino-Modifier C7 CPG 500 (Discontinued. 20-2958 below is the replacement.)	20-2957-01		
3'-Amino-Modifier C7 CPG 1000	20-2958-01	0.1g	95.00
	20-2958-10	1.0g	675.00
1 μmole columns	20-2958-41	Pack of 4	140.00
0.2 μmole columns	20-2958-42	Pack of 4	85.00
10 μmole column (ABI)	20-2958-13	Pack of 1	250.00
15 μmole column (Expedite)	20-2958-14	Pack of 1	375.00
3'-Amino-Modifier Serinol CPG	20-2997-01	0.1g	95.00
	20-2997-10	1.0g	675.00
0.2 μmole columns	20-2997-42	Pack of 4	85.00
1 μmole columns	20-2997-41	Pack of 4	140.00
10 μmole column (ABI)	20-2997-13	Pack of 1	250.00
15 μmole column (Expedite)	20-2997-14	Pack of 1	375.00
3'-PT-Amino-Modifier C3 CPG	20-2954-01	0.1g	95.00
	20-2954-10	1.0g	675.00
1 μmole columns	20-2954-41	Pack of 4	140.00
0.2 μmole columns	20-2954-42	Pack of 4	85.00
10 μmole column (ABI)	20-2954-13	Pack of 1	250.00
15 μmole column (Expedite)	20-2954-14	Pack of 1	375.00
3'-PT-Amino-Modifier C6 CPG	20-2956-01	0.1g	95.00
	20-2956-10	1.0g	675.00
1 μmole columns	20-2956-41	Pack of 4	140.00
0.2 μmole columns	20-2956-42	Pack of 4	85.00
10 μmole column (ABI)	20-2956-13	Pack of 1	250.00
15 μmole column (Expedite)	20-2956-14	Pack of 1	375.00
3'-PT-Amino-Modifier C6 PS	26-2956-01	0.1g	125.00
	26-2956-10	1.0g	1025.00
200 nmole columns (AB 3900)	26-2956-52	Pack of 10	220.00
40 nmole columns (AB 3900)	26-2956-55	Pack of 10	220.00




3'-Amino-Modifier C7 CPG



3'-PT Amino-Modifier C3 CPG



3'-PT Amino-Modifier C6 CPG



3'-Amino-Modifier Serinol CPG

3'-MODIFIERS (CONT.)

The 3'-Thiol-Modifier S-S CPG supports are used to introduce 3'-thiol linkages with three and six atom spacers into oligonucleotides. DTPA CPG is used to introduce a dithiol group at the 3'-terminus. In conjunction with DTPA Phosphoramidite, it is simple to produce oligonucleotides with multiple thiol groups at the 3' terminus, which is ideal for conjugation to gold surfaces. With Glyceryl CPG the 3'-terminus of an oligonucleotide is readily oxidized by sodium periodate to form a 3'-phosphoglycylaldehyde. The aldehyde may be further oxidized to the corresponding carboxylic acid. Either the aldehyde or the carboxylate may be used for subsequent conjugation to amine-containing products.

SEE ALSO

DTPA

p72

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M


Columns

For Instrument type Add


Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)


Item	Cat. No.	Pack	Price (\$)
3'-Thiol-Modifier C3 S-S CPG	20-2933-01	0.1g	85.00
	20-2933-10	1.0g	600.00
	1 µmole columns	Pack of 4	125.00
	0.2 µmole columns	Pack of 4	75.00
	10 µmole column (ABI)	Pack of 1	225.00
15 µmole column (Expedite)	20-2933-14	Pack of 1	350.00
3'-Thiol-Modifier 6 S-S CPG	20-2938-01	0.1g	85.00
	20-2938-10	1.0g	600.00
	0.2 µmole columns	Pack of 4	75.00
	1 µmole columns	Pack of 4	125.00
	10 µmole column (ABI)	20-2938-13	Pack of 1
15 µmole column (Expedite)	20-2938-14	Pack of 1	350.00
3'-DTPA CPG (Discontinued. Contact Glen Research Technical Support for more information.)	20-2937-01		
3'-Glyceryl CPG	20-2902-01	0.1g	85.00
	20-2902-10	1.0g	600.00
	1 µmole columns	Pack of 4	125.00
	0.2 µmole columns	Pack of 4	75.00
	10 µmole column (ABI)	20-2902-13	Pack of 1
15 µmole column (Expedite)	20-2902-14	Pack of 1	350.00




3'-Thiol-Modifier C3 S-S CPG




3'-Thiol-Modifier C6 S-S CPG



3'-Thiol-Modifier 6 S-S CPG



3'-DTPA CPG




3'-Glyceryl CPG


3'-MODIFIERS (CONT.)

3'-Amino-Modifier C6 dC CPG and 3'-Amino-Modifier C6 dT CPG replace a dC and T, respectively, at the 3'-terminus. These products allow convenient labelling at the 3' without blocking the terminus from desired enzymatic activity.

<i>Item</i>	<i>Cat. No.</i>	<i>Pack</i>	<i>Price (\$)</i>	
3'-Amino-Modifier C6 dC CPG	20-2019-01	0.1g	120.00	
	20-2019-10	1.0g	995.00	
	1 μ mole columns	20-2019-41	Pack of 4	200.00
	0.2 μ mole columns	20-2019-42	Pack of 4	120.00
	10 μ mole column (ABI)	20-2019-13	Pack of 1	300.00
	15 μ mole column (Expedite)	20-2019-14	Pack of 1	450.00
3'-Amino-Modifier C6 dT CPG	20-2039-01	0.1g	96.00	
	20-2039-10	1.0g	800.00	
	1 μ mole columns	20-2039-41	Pack of 4	160.00
	0.2 μ mole columns	20-2039-42	Pack of 4	96.00
	10 μ mole column (ABI)	20-2039-13	Pack of 1	240.00
	15 μ mole column (Expedite)	20-2039-14	Pack of 1	360.00



Amino-Modifier C6 dC CPG



Amino-Modifier C6 dT CPG

CHEMICAL PHOSPHORYLATION

INTELLECTUAL PROPERTY

Chemical Phosphorylation Reagent II is covered by US Patent No.: 5,959,090 and European Patent: EP0816368.


(1) A. Guzaev, H. Salo, A. Azhayev, and H. Lonnberg, *Tetrahedron*, 1995, **51**, 9375-9384.

SEE ALSO


High load supports p27

Chemical Phosphorylation Reagent is most commonly used to phosphorylate the 5'-terminus of an oligonucleotide. Although this product is also successful in 3'-phosphorylation, 3'-Phosphate CPG allows direct preparation of oligonucleotides with a 3'-phosphate group. Chemical Phosphorylation Reagent II contains a DMT group on a side chain which is stable to base cleavage and can be left on the oligonucleotide for use in RP purification. The DMT group is later removed with aqueous acid and the side chain is eliminated after brief treatment with aqueous ammonium hydroxide to yield the 5'-phosphate.¹ Solid CPR II is similar in performance to CPR II but it is easier to prepare aliquots since it is a powder. Many researchers treat synthesis supports with a hindered base (e.g., diethylamine, diisopropylethylamine, or DBU) post-synthesis to eliminate and remove the cyanoethyl phosphate groups. In this way, the acrylonitrile formed in situ is removed from the support and is not available to alkylate dT residues at the N3 position in the oligos. Since the sulfonylethyl group in 3'-Phosphate CPG is also susceptible to β -elimination leading to oligo cleavage, this technique is not compatible with 3'-phosphate CPG. Using CPR II CPG, which is base labile but does not support β -elimination, the cyanoethyl groups can be removed from the oligo prior to cleavage and base deprotection. ABI-style vials and columns are supplied unless otherwise requested.


Item	Cat. No.	Pack	Price (\$)
Chemical Phosphorylation Reagent	10-1900-90	100 μ mole	50.00
	10-1900-02	0.25g	160.00
3'-Phosphate CPG	20-2900-01	0.1g	70.00
	20-2900-10	1.0g	480.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	60.00
	10 μ mole column (ABI)	Pack of 1	180.00
15 μ mole column (Expedite)	Pack of 1	280.00	
3'-Phosphate PS	26-2900-01	0.1g	75.00
	26-2900-10	1.0g	510.00
	200 nmole columns (AB 3900)	Pack of 10	150.00
40 nmole columns (AB 3900)	Pack of 10	150.00	
3'-Phosphate CPG (High Load)	25-2900-01	0.1g	85.00
	25-2900-10	1.0g	600.00
	2.5 μ mole columns	Pack of 4	120.00
Chemical Phosphorylation Reagent II (CPR II)	10-1901-90	100 μ mole	60.00
	10-1901-02	0.25g	200.00
Solid Chemical Phosphorylation Reagent II (Solid CPR II)	10-1902-90	100 μ mole	60.00
	10-1902-02	0.25g	200.00
3'-CPR II CPG	20-2903-01	0.1g	70.00
	20-2903-10	1.0g	480.00
	0.2 μ mole columns	Pack of 4	60.00
	1 μ mole columns	Pack of 4	100.00
	10 μ mole column (ABI)	Pack of 1	180.00
	15 μ mole column (Expedite)	Pack of 1	280.00




Chemical Phosphorylation Reagent




3'-Phosphate CPG



Chemical Phosphorylation Reagent II



Solid Chemical Phosphorylation Reagent II



3'-CPR II CPG


ALDEHYDE MODIFICATION

Aldehyde modifiers would be attractive electrophilic substitutions in oligonucleotides since they are able to react with amino groups to form a Schiff's base, with hydrazino groups to form hydrazones, and with semicarbazides to form semi-carbazones. The Schiff's base is unstable and must be reduced with sodium borohydride to form a stable linkage but hydrazones and semicarbazides are very stable linkages.


Our collaboration with Epoch Biosciences, a subsidiary of Nanogen, Inc., has allowed us to offer 5'-Aldehyde-Modifier C2 Phosphoramidite. The acetal protecting group is sufficiently hydrophobic for use in RP HPLC and cartridge purification and is readily removed after oligonucleotide synthesis under standard oligonucleotide detritylation conditions with 80% acetic acid / 20% water or 2% aqueous trifluoroacetic acid during cartridge purification.

A formylindole nucleoside analogue has been used to introduce aldehyde groups within an oligonucleotide or at the 5' terminus. This product has no protecting group on the aldehyde, which means that deprotection of the modified oligonucleotide can be done without changing preferred conditions.

<i>Item</i>	<i>Cat. No.</i>	<i>Pack</i>	<i>Price (\$)</i>
5'-Aldehyde-Modifier C2 Phosphoramidite	10-1933-90	100 μ mole	85.00
	10-1933-02	0.25g	325.00
Formylindole CE Phosphoramidite	10-1934-90	100 μ mole	85.00
	10-1934-02	0.25g	325.00



5'-Aldehyde-Modifier C2



Formylindole

INTELLECTUAL PROPERTY

These Products are for research purposes only, and may not be used for commercial, clinical, diagnostic or any other use. The Products are subject to proprietary rights of Epoch Biosciences, Inc. and are made and sold under license from Epoch Biosciences, Inc. There is no implied license for commercial use with respect to the Products and a license must be obtained directly from Epoch Biosciences, Inc. with respect to any proposed commercial use of the Products. "Commercial use" includes but is not limited to the sale, lease, license or other transfer of the Products or any material derived or produced from them, the sale, lease, license or other grant of rights to use the Products or any material derived or produced from them, or the use of the Products to perform services for a fee for third parties (including contract research).

A simple agreement must be signed before end-users and custom oligo services may purchase these products for use as defined above. <http://www.glenresearch.com/Reference/EpochProducts.pdf>

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type *Add*

Expedite	E
MerMade	M

Columns

For Instrument type *Add*

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)


SPACER MODIFIERS

SEE ALSO

PC Modifiers
Pyrrolidinep82
p59

The spacer phosphoramidites C3, 9, C12 and 18 are used to insert a spacer arm in an oligonucleotide. The compounds may be added in multiple additions when a longer spacer is required. 3'-Spacer C3 CPG may also act as a blocker of exonuclease and polymerase activity at the 3'-terminus. dSpacer is used to introduce a stable abasic site within an oligonucleotide. PC Spacer is a photocleavable C3 spacer modifier, part of our line of photocleavable (PC) modifiers.

Item	Cat. No.	Pack	Price (\$)
Spacer Phosphoramidite 9	10-1909-90	100 μ mole	75.00
	10-1909-02	0.25g	240.00
Spacer Phosphoramidite C3	10-1913-90	100 μ mole	75.00
	10-1913-02	0.25g	240.00
dSpacer CE Phosphoramidite	10-1914-90	100 μ mole	85.00
	10-1914-02	0.25g	295.00
Spacer Phosphoramidite 18	10-1918-90	100 μ mole	95.00
	10-1918-02	0.25g	240.00
Spacer C12 CE Phosphoramidite	10-1928-90	100 μ mole	95.00
	10-1928-02	0.25g	240.00
3'-Spacer C3 CPG	20-2913-01	0.1g	70.00
	20-2913-10	1.0g	480.00
	20-2913-41	Pack of 4	100.00
	20-2913-42	Pack of 4	60.00
	20-2913-13	Pack of 1	180.00
	20-2913-14	Pack of 1	280.00
PC Spacer Phosphoramidite	10-4913-90	100 μ mole	135.00
	10-4913-02	0.25g	395.00



DENDRIMERS


Dendrimers are discrete, highly branched, monodispersed polymers that possess patterns reminiscent of the branching of trees. Plain and mixed oligonucleotide dendrimers can be synthesized using novel doubling and trebling phosphoramidite synthons.^{1,2} Dendrimers offer the following advantages. Incorporation of label using γ -³²P-ATP and polynucleotide kinase increases in proportion to the number of 5'-ends. Fluorescent signal also increases in proportion to the number of 5'-ends, if spacers are incorporated between the labels and the ends of the branches. When using a dendrimeric oligonucleotide as a PCR primer, the strand bearing the dendrimer is resistant to degradation by T7 Gene 6 exonuclease making it easy to convert the double-stranded product of the PCR to a multiply labelled, single-stranded probe. Enhanced stability of DNA dendrimers makes them useful as building blocks for the 'bottom up' approach to nano-assembly. These features also suggest applications in DNA chip technology when higher temperatures are required, for example, to melt secondary structure in the target.

Item	Catalog No.	Pack	Price(\$)
Symmetric Doubler Phosphoramidite	10-1920-90	100 μ mole	150.00
	10-1920-02	0.25g	240.00
Trebler Phosphoramidite	10-1922-90	100 μ mole	180.00
	10-1922-02	0.25g	300.00
Long Trebler Phosphoramidite	10-1925-90	100 μ mole	200.00
	10-1925-02	0.25g	300.00


BRANCHING PHOSPHORAMIDITE

A branching monomer is required to construct comb-like oligonucleotide probes. The developers of the comb system from Chiron Corporation evaluated³ several protecting groups for the branch point and chose levulinyl (LEV), which is specifically removed using a reagent containing hydrazine hydrate, acetic acid and pyridine.


Item	Catalog No.	Pack	Price(\$)
5-Me-dC Brancher Phosphoramidite	10-1018-90	100 μ mole	205.00
	10-1018-02	0.25g	505.00




Symmetric Doubler



Trebler



Long Trebler



5-Me-dC Brancher

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers For Instrument type	Add
Expedite	E
MerMade	M

Columns
For Instrument type

Columns For Instrument type	Add
Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

REFERENCES

- (1) M.S. Shchepinov, I.A. Udalova, A.J. Bridgman, and E.M. Southern, *Nucleic Acids Res*, 1997, **25**, 4447-4454.
- (2) M.S. Shchepinov, K.U. Mir, J.K. Elder, M.D. Frank-Kamenetskii, and E.M. Southern, *Nucleic Acids Res*, 1999, **27**, 3035-41.
- (3) T. Horn, C.A. Chang, and M.S. Urdea, *Nucleic Acids Res*, 1997, **25**, 4842-4849.

INTELLECTUAL PROPERTY

Doublers and Treblers are supplied under license from ISIS Innovation Limited.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

INTELLECTUAL PROPERTY

Glen Research offers PC Biotin, PC Amino-Modifier and PC Spacer products in association with AmberGen, Inc. and Link Technologies, Ltd. For a commercial application license, please contact AmberGen, Inc., +617-923-9990, (sales@ambergenc.com), <http://www.ambergenc.com/>.

PC Linker phosphoramidite is available from Glen Research in association with Link Technologies Ltd (Scotland).

SEE ALSO

5'-Biotin Phosphoramidite p92

REFERENCES





- (1) P. Ordoukhanian and J-S. Taylor, *J. Am. Chem. Soc.*, **117**, 9570-9571, 1995.
- (2a) F. Hausch and A. Jäschke, *Nucleic Acids Research*, 2000, **28**, e35.
- (2b) F. Hausch and A. Jäschke, *Tetrahedron*, 2001, **57**, 1261-1268.
- (3) T. Wenzel, T. Elssner, K. Fahr, J. Bimmler, S. Richter, I. Thomas, and M. Kostrzewa, *Nucleosides, Nucleotides & Nucleic Acids*, 2003, **22**, 1579-1581.

PHOTOCLEAVABLE MONOMERS

PC Biotin Phosphoramidite can be used to prepare 5'-biotinylated oligonucleotides suitable for capture by streptavidin in a mode similar to our popular 5' Biotin Phosphoramidite. Amino- and thiol-modified oligonucleotides have proven to be very useful for the attachment of a variety of haptens and fluorophores, as well as for the tethering of the oligonucleotides to a diversity of beads and surfaces. **PC Amino-Modifier Phosphoramidite** is used to prepare 5'-amino-modified oligonucleotides suitable for subsequent photocleavage. **PC Spacer Phosphoramidite** can be used as an intermediary to attach any modification reagent, available as a phosphoramidite, to the terminus of oligonucleotides. After photocleavage, a 5'-phosphate is generated on the DNA, rendering it suitable for further biological transformations, such as gene construction and cloning after ligation.

A versatile photocleavable DNA building block has been described by researchers in Washington University, Missouri and used in phototriggered hybridization.¹ This reagent has also been used in the design of multifunctional DNA and RNA conjugates² for the in vitro selection of new molecules catalyzing biomolecular reactions. Researchers at Bruker Daltonik in Germany have also developed genoSNIP, a method for single-nucleotide polymorphism (SNP) genotyping by MALDI-TOF mass spectrometry.³ This method uses size reduction of primer extension products by incorporation of the photocleavable linker for phototriggering strand breaks near to the 3' end of the extension primer. **PC Linker** can be incorporated into oligonucleotides at any position by standard automated DNA synthesis methodology. **PC Linker Phosphoramidite** has the added advantage in that photocleavage results in monophosphate fragments at both the 3'- and 5'-termini of the oligonucleotide fragments.

Item	Catalog No.	Pack	Price(\$)
PC Biotin Phosphoramidite	10-4950-95	50 µmole	145.00
	10-4950-90	100 µmole	280.00
	10-4950-02	0.25g	675.00
PC Amino-Modifier Phosphoramidite	10-4906-90	100 µmole	135.00
	10-4906-02	0.25g	395.00
PC Spacer Phosphoramidite	10-4913-90	100 µmole	135.00
	10-4913-02	0.25g	395.00
PC Linker Phosphoramidite	10-4920-90	100 µmole	255.00
	10-4920-02	0.25g	795.00




CONJUGATION USING CLICK CHEMISTRY

The copper(I)-catalyzed azide-alkyne cycloaddition (CuAAC) reaction between azides and alkynes to form 1,2,3-triazoles, as reported¹ by Sharpless, was found to be so exquisitely regioselective and efficient at even the most mild conditions that Sharpless coined the term 'Click Chemistry' to describe it. The use of this method for DNA modification has been somewhat delayed by the fact that copper ions damage DNA, typically yielding strand breaks.² As these problems have now been overcome by the use of copper(I)-stabilizing ligands (e.g., tris(benzyltriazolylmethyl)amine, TBTA³), Carell et al. and Seela et al. discovered that the CuAAC reaction can be used to functionalize alkyne-modified DNA nucleobases with extremely high efficiency.⁴

Oligonucleotides bearing a single nucleosidic alkyne group can be prepared using a C8-Alkyne-dC or dT-CE Phosphoramidite. Purified oligonucleotides are usually modified with 2-5 equivalents of the corresponding marker-azide (e.g., fluorescent-dye azides). After the addition of precomplexed Cu(I), complete conversion to the labelled oligo is observed in a time span between 30 min and 4 hours. After a simple precipitation step, labelled oligonucleotides can be recovered in near quantitative yields. Using a combination of C8-Alkyne, C8-TIPS-Alkyne and C8-TMS-Alkyne, it is possible to label oligonucleotides in up to three separate click reactions. The alkyne groups on the last two monomers are protected, respectively, with triisopropylsilyl (TIPS) and trimethylsilyl (TMS) protecting groups.^{5,6} The first click reaction on solid phase on a C8-Alkyne yields the singly modified oligonucleotide with full retention of the TIPS and/or TMS protecting group. For double click, a C8-TIPS-Alkyne is used as the second nucleoside and the TIPS protecting group is cleaved with tetrabutylammonium fluoride (TBAF) without causing any damage to the DNA. The second click reaction in solution yields the doubly modified oligonucleotide in excellent yield. For the introduction of three different labels, all three nucleosides are introduced into oligonucleotides. The first click reaction is performed directly on the resin. The singly modified oligonucleotide is subsequently cleaved from the support with concomitant cleavage of the TMS group and retention of the TIPS protecting group. The second click reaction is performed in solution. Precipitation of the doubly modified oligonucleotide, cleavage of the TIPS group with TBAF, and a subsequent third click reaction in solution furnishes the desired triply modified oligonucleotide in excellent overall yield.

5-Ethynyl-dU offers convenient click conjugation with an azide to generate a label rigidly attached to one of the oligonucleotide bases.

Item	Catalog No.	Pack	Price(\$)
C8-Alkyne-dT-CE Phosphoramidite	10-1540-95	50 µmole	165.00
	10-1540-90	100 µmole	315.00
	10-1540-02	0.25g	900.00
C8-TIPS-Alkyne-dC-CE Phosphoramidite	10-1541-95	50 µmole	295.00
	10-1541-90	100 µmole	575.00
	10-1541-02	0.25g	1275.00
C8-TMS-Alkyne-dC-CE Phosphoramidite	10-1542-95	50 µmole	270.00
	10-1542-90	100 µmole	525.00
	10-1542-02	0.25g	1275.00
C8-Alkyne-dC-CE Phosphoramidite	10-1543-95	50 µmole	225.00
	10-1543-90	100 µmole	435.00
	10-1543-02	0.25g	1125.00



REFERENCES

- [1] C.W. Tornøe, C. Christensen, M. Meldal, *J. Org. Chem.* 2002, **67**, 3057-3064; V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem.* 2002, **114**, 2708-2711; *Angew. Chem. Int. Ed.* 2002, **41**, 2596-2599.
- [2] C. J. Burrows, J. G. Muller, *Chem. Rev.* 1998, **98**, 1109 - 1151.
- [3] T. R. Chan, R. Hilgraf, K. B. Sharpless, V. V. Fokin, *Org. Lett.* 2004, **6**, 2853 - 2855.
- [4] J. Gierlich, G. A. Burley, P. M. E. Gramlich, D. M. Hammond, T. Carell, *Org. Lett.* 2006, **8**, 3639-3642. F. Seela, V. R. Sirivolu, *Chem. Biodiversity* 2006, **3**, 509-514.
- [5] P. M. E. Gramlich, S. Warncke, J. Gierlich, T. Carell, *Angew. Chem.* 2008, **120**, 3491-3493; *Angew. Chem. Int. Ed.* 2008, **47**, 3442-3444.
- [6] P. M. E. Gramlich, C. T. Wirges, A. Manetto, T. Carell, *Angew. Chem. Int. Ed.* 2008, **47**, 8350-8358.

INTELLECTUAL PROPERTY

All products of baseclick are patent protected and available in collaboration with baseclick.

Baseclick GmbH has filed the following patent applications:

1. WO2006/117161, New labelling strategies for the sensitive detection of analytes
2. WO2008/952775, Click Chemistry for the production of reporter molecules

Baseclick GmbH holds a worldwide license for the research market of the "Click Chemistry" patent from "The Scripps Research Institute":

3. WO03/101972, Copper-catalysed ligation of azides and acetylenes

CONJUGATION USING CLICK CHEMISTRY (CONT.)

Item	Catalog No.	Pack	Price(\$)
C8-TIPS-Alkyne-dT-CE Phosphoramidite	10-1544-95	50 μ mole	220.00
	10-1544-90	100 μ mole	425.00
	10-1544-02	0.25g	1020.00
C8-TMS-Alkyne-dT-CE Phosphoramidite	10-1545-95	50 μ mole	205.00
	10-1545-90	100 μ mole	395.00
	10-1545-02	0.25g	1050.00
5-Ethynyl-dU-CE Phosphoramidite	10-1554-95	50 μ mole	130.00
	10-1554-90	100 μ mole	245.00
	10-1554-02	0.25g	775.00
THPTA Ligand (Water soluble)	50-1004-92	25 μ mole	50.00
	50-1004-90	100 μ mole	180.00
Click-Solution (DMSO/t-BuOH)	50-1002-11	10 x 1.0mL	185.00

SEE ALSO

3'-Propargyl-5-Me-dC CPG p60

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M


(Please inquire for availability of vials and columns for other instrument types.)

OLIGO-CLICK KITS

Oligo-Click Kits contain an air-stable, insoluble Cu(I) source in pellet form in a pre-loaded and ready-to-use vial. Within the kit, the TBTA ligand is replaced by an activator which is compatible with both aqueous and organic solvents. This innovative combination of catalyst and ligand/activator results in a much easier labelling work-flow of only three simple steps. The preparation of the oligonucleotide labelling via CuAAC now requires only a minimal hands-on time of a few minutes or even less and can be carried out in air without any additional precautions. Glen Research is offering the following kits in collaboration with baseclick GmbH.

- Oligo-Kit M Reload: This kit has sufficient reagents for conjugating up to nine alkyne-containing oligonucleotides on a 100 nmole scale or a single oligonucleotide on a 1 μ mole scale. *The user must supply the azide and a solvent such as DMSO for dissolving the azide.*
- Oligo-Kit M Biotin, Oligo-Kit M Fluorescein and Oligo-Kit M TAMRA: Each kit has sufficient reagents for conjugating up to seven alkyne-containing oligonucleotides on a 100 nmole scale or a single oligonucleotide on a 1 μ mole scale. *Each kit contains all of the ingredients necessary, including the azide and DMSO solvent.*







Item	Catalog No.	Pack	Price(\$)
baseclick Oligo-Click-M-Reload	50-2100-01	each	120.00
baseclick Oligo-Click-M-Biotin	50-2101-01	each	200.00
baseclick Oligo-Click-M-Fluorescein	50-2102-01	each	240.00
baseclick Oligo-Click-M-TAMRA	50-2103-01	each	270.00



CONJUGATION USING CLICK CHEMISTRY (CONT.)

Oligonucleotides prepared using 5'-Hexynyl Phosphoramidite are stable to standard deprotection conditions and exhibit a slightly increased retention time on RP HPLC. Azides are not compatible with oligonucleotide synthesis using phosphoramidites so a post-synthesis reaction is required. Azidobutyrate NHS Ester is used¹ for azido-modification of amines at either the 3'-end or the 5'-end of an oligo and it can even be used for internal modification on an Amino-Modifier-C6 dX residue within the sequence. Specific to the 5'-terminus, 5'-Bromohexyl Phosphoramidite is added in the last cycle. This modifier can then be easily transformed into a 5'-azido group by displacement of bromide using sodium azide.² Alkyne NHS ester allows the functionalization of an amino moiety in a variety of molecules, including DNA and RNA oligonucleotides as well as peptides or proteins. We also offer two products for use in Click Chemistry based upon our 1,3-diol product portfolio with the serinol backbone - a phosphoramidite for adding an alkyne group at the 5' terminus or within the sequence, and a synthesis support for labelling the 3' terminus of oligonucleotides with an alkyne group. A 5'-iodo-modified oligonucleotide (prepared using 5'-Iodo-dT) can be quantitatively converted to the corresponding 5'-azide.

Item	Catalog No.	Pack	Price(\$)	
5'-Hexynyl Phosphoramidite	10-1908-90	100 µmole	60.00	
	10-1908-02	0.25g	200.00	
Azidobutyrate NHS Ester (Dissolve 2.3mg in 60µL of DMSO)	50-1904-23	2.3mg	60.00	
	50-1904-24	23mg	300.00	
5'-Bromohexyl Phosphoramidite	10-1946-90	100 µmole	60.00	
	10-1946-02	0.25g	200.00	
Alkyne-NHS Ester (Dissolve 2.3mg in 60µL of DMSO)	50-1905-23	2.3mg	60.00	
	50-1905-24	23mg	300.00	
Alkyne-Modifier Serinol Phosphoramidite	10-1992-95	50 µmole	100.00	
	10-1992-90	100 µmole	185.00	
	10-1992-02	0.25g	575.00	
3'-Alkyne-Modifier Serinol CPG	20-2992-01	0.1g	105.00	
	20-2992-10	1.0g	800.00	
	0.2 µmole columns	20-2992-42	Pack of 4	100.00
	1 µmole columns	20-2992-41	Pack of 4	175.00
	10 µmole column (ABI)	20-2992-13	Pack of 1	260.00
15 µmole column (Expedite)	20-2992-14	Pack of 1	390.00	
5'-I-dT-CE Phosphoramidite	10-1931-90	100 µmole	85.00	
	10-1931-02	0.25g	295.00	



REFERENCES

- (1) R. Kumar, et al., *Journal of the American Chemical Society*, 2007, **129**, 6859-6864.
- (2) J. Lietard, A. Meyer, J.J. Vasseur, and F. Morvan, *Tetrahedron Letters*, 2007, **48**, 8795-8798.


SEE ALSO

Serinol Products

p89

STABILITY NOTES

Oligonucleotides containing a 5'-iodo group are prepared conventionally with the exception that deprotection is carried out in ammonium hydroxide at room temperature for 24 hours. Under these conditions, degradation of the iodo group was less than 2%.



COPPER-FREE CLICK CHEMISTRY

At Glen Research, our goal was to offer a copper-free click phosphoramidite reagent with the following properties:

- Simple to use
- Stable in solution on the synthesizer
- Stable to ammonium hydroxide and AMA
- Excellent click performance in 17 hours or less at room temperature

From the variety of cyclooctyne-based copper-free click reagents so far described, we have chosen to offer compounds based on a dibenzo-cyclooctyne (DBCO) structure. We are offering 5'-DBCO-TEG Phosphoramidite for preparing oligos with a 5'-DBCO modification and DBCO-dT-CE Phosphoramidite for inserting a DBCO group at any position within the oligonucleotide. DBCO-sulfo-NHS Ester is also offered for post-synthesis conjugation reactions. DBCO-modified oligos may be conjugated with azides in organic solvents, such as DMSO, or aqueous buffers. Depending on the azide used, the reaction will go to completion in 4-17 hours at room temperature. Simple desalting on a Glen Gel-Pak™ leads to a product with virtually quantitative conjugation efficiency.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M


Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

Item	Catalog No.	Pack	Price(\$)
5'-DBCO-TEG Phosphoramidite	10-1941-95	50 μmole	125.00
	10-1941-90	100 μmole	230.00
	10-1941-02	0.25g	775.00
DBCO-dT-CE Phosphoramidite	10-1539-95	50 μmole	250.00
	10-1539-90	100 μmole	485.00
	10-1539-02	0.25g	975.00
DBCO-sulfo-NHS Ester (Dissolve 5.2mg in 60μL water or DMSO)	50-1941-23	5.2mg	60.00
	50-1941-24	52mg	300.00



CONJUGATION USING CLICK CHEMISTRY (CONT.)


Glen Research is offering first our most popular labels for general interest and, subsequently, we will add azide products that are not compatible with phosphoramidite chemistry.

Biotin is still our most commonly used label and biotinTEG, with its hydrophilic triethylene glycol spacer, is the most popular biotin product. Desthiobiotin is a biotin analogue that is well captured by streptavidin but the captured product can be easily released by applying a biotin solution to the streptavidin beads. 6-FAM is our most popular fluorescein derivative and we offer azides of both 6-FAM and pivaloyl-protected 6-FAM for situations where subsequent reactions require the 6-FAM to be protected. In both 6-FAM products, the hydrophilic TEG spacer is again used. The azides are offered in 25 and 100 μ mole packs for convenient oligonucleotide labelling.


7-Hydroxycoumarin, also known as umbelliferone, is a highly fluorescent, pH-sensitive fluorophore that emits in the blue region of the spectrum. However, its fluorescence is strongly quenched if the hydroxyl is alkylated or phosphorylated, making it useful in high-throughput screening for phosphatases and lipases. Interestingly, it was found that the 3-azido derivative is also highly quenched but, upon reaction with an alkyne in the presence of copper to form the triazole, the fluorescence is restored.¹ The clicked coumarin emits at a lambda max of 480 nm and absorbs at 358 nm.

HEX and TET are two of our most popular fluorescein-based dyes for labelling oligonucleotides. We are happy to offer 6-HEX and 6-TET Azides for use in click conjugations.


Item	Catalog No.	Pack	Price(\$)
BiotinTEG Azide	50-2000-92	25 μ mole	150.00
	50-2000-90	100 μ mole	450.00
DesthiobiotinTEG Azide	50-2001-92	25 μ mole	135.00
	50-2001-90	100 μ mole	400.00
Dipivaloyl 6-FAM-TEG Azide	50-2002-92	25 μ mole	230.00
	50-2002-90	100 μ mole	690.00
6-FAM-TEG Azide	50-2003-92	25 μ mole	180.00
	50-2003-90	100 μ mole	540.00
Coumarin Azide	50-2004-92	25 μ mole	115.00
	50-2004-90	100 μ mole	350.00
6-HEX Azide	50-2005-92	25 μ mole	150.00
	50-2005-90	100 μ mole	450.00
6-TET Azide	50-2006-92	25 μ mole	150.00
	50-2006-90	100 μ mole	450.00




BiotinTEG Azide




DesthiobiotinTEG Azide




Coumarin Azide




Dipivaloyl 6-FAM-TEG Azide



6-FAM-TEG Azide



6-HEX Azide



6-TET Azide

REFERENCE

- (1) J. Gierlich, G.A. Burley, P.M. Gramlich, D.M. Hammond, and T. Carell, *Org Lett*, 2006, **8**, 3639-42.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)


CONJUGATION USING CLICK CHEMISTRY (CONT.)

Two nitroxide spin labels, TEMPO Azide and TEMPO-TEG Azide, for site directed spin labelling (SDSL) are now offered.

Click Chemistry with psoralen azide and one of our many nucleosidic and non-nucleosidic alkyne derivatives has the potential to generate a variety of practical cross-linkers. The well known reversible cross-linking behavior of psoralen with an adjacent thymidine residue could be very useful.

To better address applications in near-infrared (NIR) imaging, Glen Research is offering a water soluble Disulfo-Cyanine 7 azide that can be easily conjugated to DNA and RNA through standard click chemistry. This long wavelength dye offers the benefits of improved solubility, reduced aggregation, and improved stability in the near-infrared spectrum along with the convenience of click chemistry.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
TEMPO Azide	50-2007-92	25 μ mole	115.00
	50-2007-90	100 μ mole	350.00
TEMPO-TEG Azide	50-2008-92	25 μ mole	135.00
	50-2008-90	100 μ mole	400.00
Psoralen Azide	50-2009-92	25 μ mole	115.00
	50-2009-90	100 μ mole	350.00
Disulfo-Cyanine 7 Azide	50-2010-92	25 μ mole	325.00
	50-2010-90	100 μ mole	975.00




SERINOL REAGENTS FOR MODIFICATION AND LABELLING

Most popular non-nucleosidic phosphoramidites for modification and labelling are based on two structural types: 1,2-diols and 1,3-diols. Products based on a 1,2-diol backbone were first described to allow amino-modification and biotin labelling. Technically, the 1,2-diol backbone has some drawbacks relative to the 1,3-diol backbone. The 1,2-diol backbone can participate in a dephosphorylation reaction since the 1,2-diol can form a favored 5-membered cyclic phosphate intermediate. This reaction is competitive with simple hydrolysis of the protecting groups and leads to some loss of label. However, the degree of loss at the 3' terminus can be limited by the removal of the cyanoethyl protecting group using DBU or diethylamine prior to the cleavage and deprotection steps. Similarly, loss at the 5' terminus can be eliminated by retaining the DMT group until the oligo is fully deprotected. Fortunately, the elimination reaction is virtually non-existent in the 1,3-diol backbone since the cyclic intermediate would be a 6-membered ring which is not favored for a cyclic phosphate intermediate.

IVD customers have requested a new backbone based on a 1,3-diol that would overcome any technical or IP issues surrounding our current products. We now offer a line of products based on the serinol backbone, which have been developed in close collaboration between Glen Research and Nelson Biotechnologies. Protected Biotin Serinol Phosphoramidite and CPG are protected with a *t*-butylbenzoyl group on the biotin ring. This group is designed to stop any phosphoramidite reactions at this active position in biotin. This protection avoids branching when using nucleophilic activators like DCI. The protecting group is easily removed during oligonucleotide cleavage and deprotection. The BiotinLC versions are similarly protected and should be useful for the synthesis of highly sensitive biotinylated probes. 6-Fluorescein Serinol Phosphoramidite and CPG are designed to prepare oligonucleotides containing one or several 6-Fluorescein (6-FAM) residues. Amino-Modifier Serinol Phosphoramidite and CPG are used to add amino groups into one or several positions in oligonucleotides. The amino group is protected with Fmoc, which may be removed on the synthesis column prior to solid-phase conjugation to the amino groups, or which may be removed during deprotection for subsequent solution phase conjugation to the amino groups.

We offer two products for use in Click Chemistry based upon our 1,3-diol product portfolio with the serinol backbone - a phosphoramidite for adding an alkyne group at the 5' terminus or within the sequence, and a synthesis support for labelling the 3' terminus of oligonucleotides with an alkyne group.

Item	Catalog No.	Pack	Price(\$)
Protected Biotin Serinol Phosphoramidite	10-1993-95	50 µmole	165.00
	10-1993-90	100 µmole	295.00
	10-1993-02	0.25g	675.00
6-Fluorescein Serinol Phosphoramidite	10-1994-95	50 µmole	165.00
	10-1994-90	100 µmole	295.00
	10-1994-02	0.25g	595.00
Protected BiotinLC Serinol Phosphoramidite	10-1995-95	50 µmole	205.00
	10-1995-90	100 µmole	365.00
	10-1995-02	0.25g	675.00
Amino-Modifier Serinol Phosphoramidite	10-1997-95	50 µmole	125.00
	10-1997-90	100 µmole	225.00
	10-1997-02	0.25g	595.00



SERINOL REAGENTS FOR MODIFICATION AND LABELLING (CONT.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M


Columns

For Instrument type Add


Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)


Item	Catalog No.	Pack	Price(\$)
Alkyne-Modifier Serinol Phosphoramidite	10-1992-95	50 µmole	100.00
	10-1992-90	100 µmole	185.00
	10-1992-02	0.25g	575.00
3'-Protected Biotin Serinol CPG	20-2993-01	0.1g	120.00
	20-2993-10	1.0g	995.00
	0.2 µmole columns	Pack of 4	120.00
	1 µmole columns	Pack of 4	200.00
	10 µmole column (ABI)	Pack of 1	300.00
	15 µmole column (Expedite)	Pack of 1	450.00
3'-6-Fluorescein Serinol CPG	20-2994-01	0.1g	120.00
	20-2994-10	1.0g	995.00
	0.2 µmole columns	Pack of 4	120.00
	1 µmole columns	Pack of 4	200.00
	10 µmole column (ABI)	Pack of 1	300.00
	15 µmole column (Expedite)	Pack of 1	450.00
3'-Protected BiotinLC Serinol CPG	20-2995-01	0.1g	120.00
	20-2995-10	1.0g	995.00
	0.2 µmole columns	Pack of 4	120.00
	1 µmole columns	Pack of 4	200.00
	10 µmole column (ABI)	Pack of 1	300.00
	15 µmole column (Expedite)	Pack of 1	450.00
3'-Amino-Modifier Serinol CPG	20-2997-01	0.1g	95.00
	20-2997-10	1.0g	675.00
	0.2 µmole columns	Pack of 4	85.00
	1 µmole columns	Pack of 4	140.00
	10 µmole column (ABI)	Pack of 1	250.00
	15 µmole column (Expedite)	Pack of 1	375.00
3'-Alkyne-Modifier Serinol CPG	20-2992-01	0.1g	105.00
	20-2992-10	1.0g	800.00
	0.2 µmole columns	Pack of 4	100.00
	1 µmole columns	Pack of 4	175.00
	10 µmole column (ABI)	Pack of 1	260.00
	15 µmole column (Expedite)	Pack of 1	390.00




Alkyne-Modifier Serinol Phosphoramidite




3'-Alkyne-Modifier Serinol CPG




Protected Biotin Serinol CPG



Protected BiotinLC Serinol CPG



Amino-Modifier Serinol CPG




6-Fluorescein Serinol CPG


DABCYL LABELLING

A molecular beacon probe¹ has its natural fluorescence quenched in solution unless it is hybridized to the target sequence. Consequently, the design of a molecular beacon requires a fluorophore to be in one part of the sequence and the quencher molecule to be in another, with both molecules being separated from the oligonucleotide by a hydrocarbon spacer. The DabcyI group has been found to be a universal quencher. 3'-DabcyI CPG and 3'-DabcyI CPG are used to prepare probes with the quencher blocking the 3'-terminus. 5'-DabcyI Phosphoramidite locates the quencher at the 5'-terminus and DabcyI-dT places it within the sequence, leaving the 3'-terminus available for polymerase extension.


Item	Catalog No.	Pack	Price(\$)	
3'-DabcyI CPG	20-5911-01	0.1g	120.00	
	20-5911-10	1.0g	975.00	
	1 μmole columns	20-5911-41	Pack of 4	200.00
	0.2 μmole columns	20-5911-42	Pack of 4	120.00
	10 μmole column (ABI)	20-5911-13	Pack of 1	350.00
15 μmole column (Expedite)	20-5911-14	Pack of 1	500.00	
3'-DabcyI CPG	20-5912-01	0.1g	120.00	
	20-5912-10	1.0g	975.00	
	1 μmole columns	20-5912-41	Pack of 4	200.00
	0.2 μmole columns	20-5912-42	Pack of 4	120.00
	10 μmole column (ABI)	20-5912-13	Pack of 1	350.00
15 μmole column (Expedite)	20-5912-14	Pack of 1	500.00	
3'-DabcyI PS	26-5912-01	0.1g	125.00	
	26-5912-10	1.0g	1025.00	
	200 nmole columns (AB 3900)	26-5912-52	Pack of 10	300.00
	40 nmole columns (AB 3900)	26-5912-55	Pack of 10	300.00
DabcyI-dT	10-1058-95	50 μmole	180.00	
	10-1058-90	100 μmole	325.00	
	10-1058-02	0.25g	675.00	
5'-DabcyI Phosphoramidite	10-5912-95	50 μmole	125.00	
	10-5912-90	100 μmole	225.00	
	10-5912-02	0.25g	650.00	




DabcyI CPG



DabcyI CPG



DabcyI-dT




5'-DabcyI Phosphoramidite

REFERENCE

(1) S. Tyagi and F.R. Kramer, *Nature Biotechnology*, 1996, **4**, 303-308.

DYE QUENCHER PLOT



http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf

BIOTIN LABELLING (CONT.)

Biotin-dT can replace dT residues within the oligonucleotide sequence. 5'-Biotin phosphoramidite can be added ONLY ONCE to the 5'-terminus of an oligonucleotide. However, the DMT group on the biotin can be used in RP cartridge and HPLC purification techniques. PC Biotin is a photocleavable 5'-biotin phosphoramidite. BiotinTEG CPG and Protected BiotinLC Serinol CPG are designed for the direct synthesis of oligonucleotides containing biotin at the 3' terminus.


Desthiobiotin is a biotin analogue that exhibits lower binding to biotin-binding proteins such as streptavidin. This biotin analogue is lacking the sulfur group from the molecule and has a dissociation constant (Kd) several orders of magnitude less than biotin/streptavidin. As a result, biomolecules containing desthiobiotin are dissociated from streptavidin simply in the presence of buffered solutions of biotin. We offer desthiobiotinTEG phosphoramidite and the corresponding CPG.

ABI-style vials and columns are supplied unless otherwise requested (see note box).


Item	Catalog No.	Pack	Price (\$)
5'-Biotin Phosphoramidite	10-5950-95	50 μ mole	125.00
	10-5950-90	100 μ mole	225.00
	10-5950-02	0.25g	650.00
Biotin-dT	10-1038-95	50 μ mole	167.50
	10-1038-90	100 μ mole	325.00
	10-1038-02	0.25g	625.00
PC Biotin Phosphoramidite	10-4950-95	50 μ mole	145.00
	10-4950-90	100 μ mole	280.00
	10-4950-02	0.25g	675.00
DesthiobiotinTEG Phosphoramidite	10-1952-95	50 μ mole	185.00
	10-1952-90	100 μ mole	335.00
	10-1952-02	0.25g	775.00

SEE ALSO


PC Biotin Phosphoramidite p82




Biotin-dT



PC Biotin Phosphoramidite



5'-Biotin Phosphoramidite



DesthiobiotinTEG Phosphoramidite

BIOTIN LABELLING (CONT.)

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
3'-BiotinTEG CPG	20-2955-01	0.1g	120.00
	20-2955-10	1.0g	995.00
	0.2 μ mole columns	Pack of 4	120.00
	1 μ mole columns	Pack of 4	200.00
	10 μ mole column (ABI)	Pack of 1	300.00
15 μ mole column (Expedite)	20-2955-14	Pack of 1	450.00
3'-BiotinTEG PS	26-2955-01	0.1g	125.00
	26-2955-10	1.0g	1025.00
	200 nmole columns (AB 3900)	Pack of 10	300.00
	40 nmole columns (AB 3900)	Pack of 10	300.00
3'-Protected Biotin Serinol CPG	20-2993-01	0.1g	120.00
	20-2993-10	1.0g	995.00
	0.2 μ mole columns	Pack of 4	120.00
	1 μ mole columns	Pack of 4	200.00
	10 μ mole column (ABI)	Pack of 1	300.00
15 μ mole column (Expedite)	20-2993-14	Pack of 1	450.00
3'-Protected BiotinLC Serinol CPG	20-2995-01	0.1g	120.00
	20-2995-10	1.0g	995.00
	0.2 μ mole columns	Pack of 4	120.00
	1 μ mole columns	Pack of 4	200.00
	10 μ mole column (ABI)	Pack of 1	300.00
15 μ mole column (Expedite)	20-2995-14	Pack of 1	450.00
DesthiobiotinTEG CPG	20-2952-01	0.1g	140.00
	20-2952-10	1.0g	1150.00
	0.2 μ mole columns	Pack of 4	140.00
	1 μ mole columns	Pack of 4	230.00
	10 μ mole column (ABI)	Pack of 1	345.00
15 μ mole column (Expedite)	20-2952-14	Pack of 1	520.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add


Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)



FLUORESCIN LABELLING


5'-Fluorescein phosphoramidite contains no 4,4'-dimethoxytrityl (DMT) group and can be added only once at the 5'-terminus, thereby terminating synthesis. This product is prepared using the 6-carboxyfluorescein derivative. The tetrachloro-, hexachloro- and dichloro-dimethoxy-fluorescein (TET, HEX and JOE, respectively) phosphoramidites are designed to take advantage of the multicolor detection capability of modern DNA sequencers and genetic analyzers. Fluorescein phosphoramidite is designed to produce the same fluorescein-type structure as had been previously prepared using fluorescein isothiocyanate (FITC). Our fluorescein phosphoramidite also contains a DMT group to allow quantification of coupling. The analogous structure, 6-Fluorescein Phosphoramidite, prepared using 6-FAM, is also available, along with 6-Fluorescein Serinol Phosphoramidite. Fluorescein-dT can be inserted into the desired sequence as a replacement for a dT residue.

We offer five fluorescein supports. Fluorescein CPG has traditionally been used to add the fluorescein label at the 3'-terminus. The analogous structure, 3'-(6-Fluorescein) CPG, prepared using 6-FAM, is now also available, along with 6-Fluorescein Serinol CPG. We also offer 3'-(6-FAM) CPG and Fluorescein-dT CPG, both derivatives of 6-carboxyfluorescein (6-FAM). Both are single isomers and use an amide linkage which is stable during cleavage and deprotection and does not allow isomer formation. 3'-(6-FAM) CPG allows effective blockage of the 3'-terminus from polymerase extension as well as exonuclease digestion. Fluorescein-dT CPG allows both of these enzymatic activities to proceed. Normal cleavage and deprotection with ammonium hydroxide readily generates the fluorescein labelled oligos.


The spectral characteristics of these dyes are detailed on the following page.

Item	Cat. No.	Pack	Price (\$)
5'-Fluorescein Phosphoramidite (6-FAM)	10-5901-95	50 µmole	110.00
	10-5901-90	100 µmole	215.00
	10-5901-02	0.25g	575.00
5'-Hexachloro-Fluorescein Phosphoramidite (HEX)	10-5902-95	50 µmole	190.00
	10-5902-90	100 µmole	375.00
	10-5902-02	0.25g	875.00
5'-Tetrachloro-Fluorescein Phosphoramidite (TET)	10-5903-95	50 µmole	180.00
	10-5903-90	100 µmole	350.00
	10-5903-02	0.25g	875.00
5'-Dichloro-dimethoxy-Fluorescein Phosphoramidite II (JOE)	10-5906-95	50 µmole	105.00
	10-5906-90	100 µmole	198.00
	10-5906-02	0.25g	495.00


DYE QUENCHER PLOT




http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf




5'-Fluorescein Phosphoramidite



5'-Hexachloro-Fluorescein Phosphoramidite







5'-Tetrachloro-Fluorescein Phosphoramidite




5'-Dichloro-dimethoxy-Fluorescein Phosphoramidite II

FLUORESC EIN LABELLING (CONT.)

Item	Cat. No.	Pack	Price (\$)	
3'-Fluorescein CPG	20-2963-01	0.1g	120.00	
	20-2963-10	1.0g	995.00	
	1 μ mole columns	20-2963-41	Pack of 4	200.00
	0.2 μ mole columns	20-2963-42	Pack of 4	120.00
	10 μ mole column (ABI)	20-2963-13	Pack of 1	300.00
15 μ mole column (Expedite)	20-2963-14	Pack of 1	450.00	
3'-(6-Fluorescein) CPG	20-2964-01	0.1g	120.00	
	20-2964-10	1.0g	995.00	
	1 μ mole columns	20-2964-41	Pack of 4	200.00
	0.2 μ mole columns	20-2964-42	Pack of 4	120.00
	10 μ mole column (ABI)	20-2964-13	Pack of 1	300.00
15 μ mole column (Expedite)	20-2964-14	Pack of 1	450.00	
3'-(6-FAM) CPG	20-2961-01	0.1g	120.00	
	20-2961-10	1.0g	995.00	
	1 μ mole columns	20-2961-41	Pack of 4	200.00
	0.2 μ mole columns	20-2961-42	Pack of 4	120.00
	10 μ mole column (ABI)	20-2961-13	Pack of 1	300.00
15 μ mole column (Expedite)	20-2961-14	Pack of 1	450.00	
3'-(6-FAM) PS	26-2961-01	0.1g	130.00	
	26-2961-10	1.0g	1045.00	
	200 nmole columns (AB 3900)	26-2961-52	Pack of 10	300.00
40 nmole columns (AB 3900)	26-2961-55	Pack of 10	300.00	
3'-6-Fluorescein Serinol CPG	20-2994-01	0.1g	120.00	
	20-2994-10	1.0g	995.00	
	0.2 μ mole columns	20-2994-42	Pack of 4	120.00
	1 μ mole columns	20-2994-41	Pack of 4	200.00
	10 μ mole column (ABI)	20-2994-13	Pack of 1	300.00
15 μ mole column (Expedite)	20-2994-14	Pack of 1	450.00	



DYE QUENCHER PLOT



http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf

FLUORESCIN LABELLING (CONT.)

Item	Cat. No.	Pack	Price (\$)
3'-Fluorescein-dT CPG	20-2056-01	0.1g	120.00
	20-2056-10	1.0g	995.00
1 μ mole columns	20-2056-41	Pack of 4	200.00
0.2 μ mole columns	20-2056-42	Pack of 4	120.00
10 μ mole column (ABI)	20-2056-13	Pack of 1	300.00
15 μ mole column (Expedite)	20-2056-14	Pack of 1	450.00

FLUORESCIN LABELLING (SIMA)

Dichloro-diphenyl-fluorescein, SIMA (HEX) exhibits virtually identical absorbance and emission spectra to HEX. SIMA (HEX) is much more stable to basic deprotection conditions than HEX and oligonucleotides can be deprotected using ammonium hydroxide at elevated temperatures and even ammonium hydroxide/methylamine (AMA) at room temperature or 65°C for 10 minutes. SIMA absorption maximum was 3 nm blue-shifted compared to HEX at pH 7. The absorbance is broader, so the extinction coefficient is smaller than that of HEX, but when exciting at 500 nm where the absorbance was normalized, the emission was still 90% of HEX and the emission was red-shifted by 5 nm. A second SIMA (HEX) product, SIMA (HEX)-dT, can be used to introduce SIMA (HEX) in the synthetic oligonucleotide sequence, usually as a replacement for the native dT linkage. Again, this product is fully compatible with deprotection schemes using ammonium hydroxide at elevated temperatures or AMA at room temperature and 65°C.

Item	Cat. No.	Pack	Price (\$)
SIMA (HEX) Phosphoramidite	10-5905-95	50 μ mole	90.00
	10-5905-90	100 μ mole	175.00
	10-5905-02	0.25g	400.00
SIMA (HEX)-dT Phosphoramidite	10-5945-95	50 μ mole	345.00
	10-5945-90	100 μ mole	675.00
	10-5945-02	0.25g	995.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers
For Instrument type Add


Expedite	E
MerMade	M

Columns
For Instrument type Add




Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

DYE QUENCHER PLOT




http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf




RHODAMINE (TAMRA) LABELLING

Rhodamine derivatives are not sufficiently stable to survive conventional deprotection and these must be attached to amino-modified oligonucleotides using post-synthesis labelling techniques. Because Tetramethyl Rhodamine (TAMRA) is not base stable, the procedure to cleave and deprotect the labelled oligonucleotide must be carefully considered. Using the UltraMILD monomers and deprotection with potassium carbonate in methanol, TAMRA oligonucleotides can be fairly conveniently isolated. To streamline the preparation of TAMRA oligos, we offer 3'-TAMRA CPG for 3' labelling and TAMRA-dT for labelling within the sequence. We also offer TAMRA NHS ester for labelling amino-modified oligonucleotides.


Item	Cat. No.	Pack	Price (\$)
3'-TAMRA CPG	20-5910-01	0.1g	120.00
	20-5910-10	1.0g	995.00
	1 μ mole columns	Pack of 4	200.00
	0.2 μ mole columns	Pack of 4	120.00
3'-TAMRA PS	26-5910-01	0.1g	130.00
	26-5910-10	1.0g	1045.00
	200 nmole columns (AB 3900)	Pack of 10	300.00
	40 nmole columns (AB 3900)	Pack of 10	300.00
TAMRA-dT	10-1057-95	50 μ mole	250.00
	10-1057-90	100 μ mole	495.00
	10-1057-02	0.25g	975.00
TAMRA NHS Ester (Solution in anhydrous DMSO)	50-5910-66	60 μ L	240.00



TAMRA CPG



TAMRA-dT



TAMRA NHS Ester

SEE ALSO

UltraMILD monomers p68
Oligo-Click-M-TAMRA p84

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M


Columns

For Instrument type Add

Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)

DYE QUENCHER PLOT



http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf

CYANINE LABELLING

Two cyanine derivatives, Cyanine 3 and Cyanine 5, which differ in structure simply by the number of carbons in the conjugated poly-ene linkage, are joined by the closely related analogues, Cyanine 3.5 and Cyanine 5.5, and are available as phosphoramidites. Cyanine dyes are normally added once at the 5'-terminus and the MMT group should be removed on the synthesizer. The absorbance of the MMT cation (yellow) is noticeably different from the DMT cation (orange), and so, absorbance-based trityl monitors will detect it incorrectly as a low coupling. On the other hand, conductivity detectors will interpret the release more correctly. Cyanine dye phosphoramidites have also been used successfully adjacent to the 3'-terminus. Cyanine 3 and Cyanine 5 supports are also offered to allow simpler production of 3' cyanine dye-labelled oligonucleotides.

Deprotection of oligos containing Cyanine dyes may be carried out with ammonium hydroxide at room temperature, regardless of the base protecting groups on the monomers used. If there is a need to use ammonium hydroxide at elevated temperature, Cyanine 3 and Cyanine 3.5 are more stable than Cyanine 5 and Cyanine 5.5. However, it is always prudent to use monomers with base labile protecting groups to limit the exposure time to 2 hours or less at 65°C during deprotection.


To better address applications in near-infrared (NIR) imaging, Glen Research is offering a water soluble Disulfo-Cyanine 7 azide that can be easily conjugated to DNA and RNA through standard click chemistry. This long wavelength dye offers the benefits of improved solubility, reduced aggregation, and improved stability in the near-infrared spectrum along with the convenience of click chemistry.

SPECTRAL DATA FOR CYANINE DYES

	Absorbance Maximum	Emission Maximum	Color
Cyanine 3	546nm	563nm	Red
Cyanine 3.5	588nm	604nm	Purple
Cyanine 5	646nm	662nm	Violet
Cyanine 5.5	683nm	707nm	Dark Blue
Cyanine 7	750nm	773nm	Dark Green


(Measured in an oligo in 0.1M TEAA buffer, pH7.)

DYE QUENCHER PLOT




http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf


Item	Cat. No.	Pack	Price (\$)
Cyanine 3 Phosphoramidite	10-5913-95	50 µmole	205.00
	10-5913-90	100 µmole	375.00
	10-5913-02	0.25g	925.00
Cyanine 3.5 Phosphoramidite	10-5914-95	50 µmole	220.00
	10-5914-90	100 µmole	400.00
	10-5914-02	0.25g	925.00
Cyanine 5 Phosphoramidite	10-5915-95	50 µmole	205.00
	10-5915-90	100 µmole	375.00
	10-5915-02	0.25g	925.00
Cyanine 5.5 Phosphoramidite	10-5916-95	50 µmole	245.00
	10-5916-90	100 µmole	450.00
	10-5916-02	0.25g	925.00




Cyanine 3 Phosphoramidite



Cyanine 3.5 Phosphoramidite



Cyanine 5 Phosphoramidite



Cyanine 5.5 Phosphoramidite






CYANINE LABELLING (CONT.)

Cyanine 3 CPG	20-5913-01	0.1g	160.00
	20-5913-10	1.0g	1250.00
1 μmole columns (TWIST format only)	20-5913-41	Pack of 4	250.00
0.2 μmole columns	20-5913-42	Pack of 4	70.00
Cyanine 5 CPG	20-5915-01	0.1g	160.00
	20-5915-10	1.0g	1250.00
1 μmole columns (TWIST format only)	20-5915-41	Pack of 4	250.00
0.2 μmole columns	20-5915-42	Pack of 4	70.00
Disulfo-Cyanine 7 Azide	50-2010-92	25 μmole	325.00
	50-2010-90	100 μmole	975.00

DYLIGHT™ DYES

DyLight™ dyes, DY547 and DY647, are phosphoramidite alternatives to Cyanine 3 and Cyanine 5. The performance of these two dyes is similar to the equivalent Cyanine dyes and the absorption and emission spectra are virtually identical. A comparison of the emission or absorbance spectra of oligos shows that the spectra are virtually identical for Cyanine 3 and DY547. Similarly, Cyanine 5 and DY 647 are virtually identical. When oligos containing DY547 were excited at 488 nm, the quantum yield (QY) was 12% greater than oligos containing Cyanine 3. When oligos containing DY647 were excited at 580 nm, the QY was 5% greater and the emission was blue-shifted by only 1 nm compared to Cyanine 5.

Item	Catalog No.	Pack	Price(\$)
DyLight DY547 Phosphoramidite	10-5917-95	50 μmole	175.00
	10-5917-90	100 μmole	335.00
	10-5917-02	0.25g	995.00
DyLight DY647 Phosphoramidite	10-5918-95	50 μmole	175.00
	10-5918-90	100 μmole	335.00
	10-5918-02	0.25g	995.00



INTELLECTUAL PROPERTY

DyLight is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

EPOCH DYES AND QUENCHER

Glen Research's agreement with Epoch Biosciences, Inc., a subsidiary of Nanogen, Inc., allows us to distribute several of Epoch's proprietary products designed for the synthesis of novel DNA probes. We are pleased to offer products based on Epoch's Redmond Red™, Yakima Yellow™ and Gig Harbor Green™ fluorophores and Eclipse™ non-fluorescent quencher. Under our agreement we also supply PPG, a modified nucleoside and 5'-Aldehyde-Modifier C2 Phosphoramidite. The two fluorescent dyes, Yakima Yellow and Redmond Red, are available as phosphoramidites and supports. Yakima Yellow has an absorbance maximum at 530 nm and emission maximum at 549 nm, while Redmond Red's absorbance and emission maxima are at 579 nm and 595 nm, respectively. Gig Harbor Green and 6-FAM are based on the same fluorescein core structure but Gig Harbor Green is 15-20% brighter than FAM.

The Eclipse quencher from Epoch solves most of the problems inherent in the synthesis of molecular beacon and FRET probes. The Eclipse molecule is highly stable and can be used safely in all common oligo deprotection schemes. The absorbance maximum for Eclipse Quencher is at 522 nm, compared to 479 nm for dabcyI. In addition, the structure of the Eclipse Quencher is substantially more electron deficient than that of dabcyI and this leads to better quenching over a wider range of dyes, especially those with emission maxima at longer wavelengths (red shifted) such as Redmond Red and Cy5. In addition, with an absorption range from 390 nm to 625 nm, the Eclipse Quencher is capable of effective performance in a wide range of colored FRET probes.

SEE ALSO

PPG p52
5'-Aldehyde-Modifier C2 p102

FLUORESCENT DYES

	Absorbance Maximum	Emission Maximum	Color
Gig Harbor Green	494nm	525nm	Green
Yakima Yellow	530nm	549nm	Yellow
Redmond Red	579nm	595nm	Red





INTELLECTUAL PROPERTY

These Products are for research purposes only, and may not be used for commercial, clinical, diagnostic or any other use. The Products are subject to proprietary rights of Epoch Biosciences, Inc. and are made and sold under license from Epoch Biosciences, Inc. There is no implied license for commercial use with respect to the Products and a license must be obtained directly from Epoch Biosciences, Inc. with respect to any proposed commercial use of the Products. "Commercial use" includes but is not limited to the sale, lease, license or other transfer of the Products or any material derived or produced from them, the sale, lease, license or other grant of rights to use the Products or any material derived or produced from them, or the use of the Products to perform services for a fee for third parties (including contract research).

A simple agreement must be signed before end-users and custom oligo services may purchase these products for use as defined above. <http://www.glenresearch.com/Reference/EpochProducts.pdf>


Redmond Red, Yakima Yellow, Gig Harbor Green, and Eclipse are trademarks of Epoch Biosciences, Inc.

Item	Cat. No.	Pack	Price (\$)
Epoch Redmond Red™ Phosphoramidite	10-5920-95	50 µmole	220.00
	10-5920-90	100 µmole	420.00
	10-5920-02	0.25g	1045.00
Epoch Yakima Yellow™ Phosphoramidite	10-5921-95	50 µmole	230.00
	10-5921-90	100 µmole	440.00
	10-5921-02	0.25g	1045.00
Epoch Gig Harbor Green™ Phosphoramidite	10-5922-95	50 µmole	182.50
	10-5922-90	100 µmole	345.00
	10-5922-02	0.25g	920.00
Epoch Eclipse™ Quencher Phosphoramidite	10-5925-95	50 µmole	250.00
	10-5925-90	100 µmole	480.00
	10-5925-02	0.25g	1185.00




EPOCH DYES AND QUENCHER (CONT.)


Item	Cat. No.	Pack	Price (\$)	
Epoch Redmond Red™ CPG	20-5920-01	0.1g	180.00	
	20-5920-10	1.0g	1500.00	
	1 μmole columns	20-5920-41	Pack of 4	300.00
	0.2 μmole columns	20-5920-42	Pack of 4	150.00
	10 μmole column (ABI)	20-5920-13	Pack of 1	750.00
15 μmole column (Expedite)	20-5920-14	Pack of 1	1125.00	
Epoch Yakima Yellow™ CPG	20-5921-01	0.1g	180.00	
	20-5921-10	1.0g	1500.00	
	1 μmole columns	20-5921-41	Pack of 4	300.00
	0.2 μmole columns	20-5921-42	Pack of 4	150.00
	10 μmole column (ABI)	20-5921-13	Pack of 1	750.00
15 μmole column (Expedite)	20-5921-14	Pack of 1	1125.00	
Epoch Eclipse™ Quencher CPG	20-5925-01	0.1g	230.00	
	20-5925-10	1.0g	1925.00	
	1 μmole columns	20-5925-41	Pack of 4	350.00
	0.2 μmole columns	20-5925-42	Pack of 4	175.00
	10 μmole column (ABI)	20-5925-13	Pack of 1	995.00
15 μmole column (Expedite)	20-5925-14	Pack of 1	1495.00	



Epoch Redmond Red™ CPG



Epoch Yakima Yellow™ CPG



Epoch Eclipse™ Quencher CPG

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M


Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

DYE QUENCHER PLOT



http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf

BLACK HOLE QUENCHER DYES

TABLE 1: BLACK HOLE QUENCHERS

Quencher	λ_{max} (nm)	E260 (L/mol.cm)	E _{max} (L/mol.cm)
BHQ-1	534	8,000	34,000
BHQ-2	579	8,000	38,000
BHQ-3	672	13,000	42,700

REFERENCES

- S.A.E. Marras, F.R. Kramer, and S. Tyagi, *Nucleic Acids Res.*, 2002, **30**, E122.
- M.K. Johansson, H. Fidder, D. Dick, and R.M. Cook, *J Am Chem Soc*, 2002, **124**, 6950-6956.

SEE OTHER QUENCHERS





Dabcyl	p91
Eclipse™	p102
BBQ-650®	p106

INTELLECTUAL PROPERTY

"Black Hole Quencher", "BHQ-0", "BHQ-1", "BHQ-2" and "BHQ-3" are trademarks of Biosearch Technologies, Inc., Novato, CA. The BHQ dye technology is the subject of pending patents and is licensed and sold under agreement with Biosearch Technologies, Inc.. Products incorporating the BHQ dye moiety are sold exclusively for R&D use by the end-user. They may not be used for clinical or diagnostic purposes and they may not be re-sold, distributed or re-packaged.

With the growing popularity of red and near-infrared dyes, we are offering the Black Hole Quencher™ dyes (BHQs), whose physical properties are detailed in Table 1. BHQ dyes are robust dark quenchers that very nicely complement our existing product line. They are compatible with ammonium hydroxide deprotection, exhibit excellent coupling efficiencies, have large extinction coefficients and are completely non-fluorescent. Their absorbances are well-tuned to quench a variety of popular fluorophores – even those far into the red, such as Cy3 and Cy5. The dark quencher most typically used in a Molecular Beacon is Dabcyl. Because the quenching does not involve FRET, there is little, if any, dependence upon donor-acceptor spectral overlap. In a comprehensive paper by Marras, Kramer and Tyagi,¹ the ability of BHQ-1 and BHQ-2 to quench 22 different fluorophores was evaluated. For shorter wavelength fluorophores such as fluorescein, the quenching efficiency was roughly the same as Dabcyl (91% – 93%). However, for dyes emitting in the far red, such as Cy5, the BHQ dyes were far superior – quenching the Cy5 with 96% efficiency, compared to 84% with Dabcyl. This may reflect the BHQ's ability to form stable, non-fluorescent complexes which can be a plus even in FRET probes. Indeed, recent work suggests that these non-fluorescent complexes which can be a plus even in FRET probes. Indeed, recent work suggests that these non-fluorescent complexes will form even in the absence of a hairpin stem structure used by Molecular Beacons.²

Item	Cat. No.	Pack	Price (\$)
5'-BHQ-1 Phosphoramidite	10-5931-95	50 μ mole	100.00
	10-5931-90	100 μ mole	200.00
	10-5931-02	0.25g	700.00
5'-BHQ-2 Phosphoramidite	10-5932-95	50 μ mole	100.00
	10-5932-90	100 μ mole	200.00
	10-5932-02	0.25g	700.00
BHQ-1-dT	10-5941-95	50 μ mole	265.00
	10-5941-90	100 μ mole	525.00
	10-5941-02	0.25g	925.00
BHQ-2-dT	10-5942-95	50 μ mole	265.00
	10-5942-90	100 μ mole	525.00
	10-5942-02	0.25g	925.00



BLACK HOLE QUENCHER DYES (CONT.)

Item	Cat. No.	Pack	Price (\$)
3'-BHQ-1 CPG	20-5931-01	0.1g	190.00
	20-5931-10	1.0g	1500.00
	1 μ mole columns	Pack of 4	300.00
	0.2 μ mole columns	Pack of 4	80.00
	10 μ mole column (ABI)	Pack of 1	575.00
15 μ mole column (Expedite)	20-5931-14	Pack of 1	825.00
3'-BHQ-2 CPG	20-5932-01	0.1g	190.00
	20-5932-10	1.0g	1500.00
	1 μ mole columns	Pack of 4	300.00
	0.2 μ mole columns	Pack of 4	80.00
	10 μ mole column (ABI)	Pack of 1	575.00
15 μ mole column (Expedite)	20-5932-14	Pack of 1	825.00
3'-BHQ-3 CPG	20-5933-01	0.1g	190.00
	20-5933-10	1.0g	1500.00
	1 μ mole columns	Pack of 4	300.00
	0.2 μ mole columns	Pack of 4	80.00
	10 μ mole column (ABI)	Pack of 1	575.00
15 μ mole column (Expedite)	20-5933-14	Pack of 1	825.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M


Columns

For Instrument type Add




Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

DYE QUENCHER PLOT



http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf




BLACKBERRY® QUENCHER (BBQ-650®)

INTELLECTUAL PROPERTY

BlackBerry® Quencher technology: US Patent 7,879,986. The purchase of BlackBerry® Quencher reagents includes a limited license to use these reagents exclusively for research and development purposes. They may not be used for clinical or diagnostic purposes and they may not be re-sold, distributed, or re-packaged without prior agreement and consent of Berry & Associates, Inc. Subsequent sale of products that are derived from BlackBerry® Quencher reagents is permitted so long as the following written disclaimer is included in written and electronic catalogs, in commercial advertisement, and in packages with containers of such derivative products: "BlackBerry is a trademark of Berry & Associates, Inc. Products derived from BlackBerry® Quencher reagents are sold exclusively for research and development use by the purchaser. They may not be used for clinical or diagnostic purposes without prior agreement and consent of Berry & Associates, Inc."

DYE QUENCHER PLOT




http://www.glenresearch.com/ProductFiles/Dye_Quencher_Plot.pdf

We are happy to offer several products containing the BlackBerry® Quencher (BBQ-650®), which exhibits a broad absorption profile from 550nm to 750nm, centered at 650nm. This range offers more effective quenching of some of our popular long wavelength dyes like TAMRA, Redmond Red, Cy dyes and DyLight dyes. We offer BBQ-650 products for the 3' and 5' termini, as well as BBQ-650-dT for inclusion within the oligonucleotide sequence, with the following properties:

- Quenches the fluorescence of long wavelength dyes
- Quenches in FRET and contact mode
- Absorbance maximum at ~650nm
- Quenching range – 550–750nm
- Compatible with standard oligo synthesis chemistry
- Compatible with regular deprotection but requires mild deprotection with AMA at room temperature
- Available for 3', 5', and internal substitution
- More stable than BHQ-3

Item	Cat. No.	Pack	Price (\$)
5'-BBQ-650® Phosphoramidite	10-5934-95	50 µmole	160.00
	10-5934-90	100 µmole	305.00
	10-5934-02	0.25g	925.00
BBQ-650®-dT	10-5944-95	50 µmole	280.00
	10-5944-90	100 µmole	545.00
	10-5944-02	0.25g	925.00
3'-BBQ-650® CPG	20-5934-01	0.1g	190.00
	20-5934-10	1.0g	1500.00
	20-5934-41	Pack of 4	300.00
	20-5934-42	Pack of 4	80.00
	20-5934-13	Pack of 1	575.00
	20-5934-14	Pack of 1	825.00

- 1 µmole columns
- 0.2 µmole columns
- 10 µmole column (ABI)
- 15 µmole column (Expedite)



ACRIDINE LABELLING

Acridine phosphoramidite is designed to produce an oligonucleotide containing acridine at any position in the molecule. Acridine CPG is used to label the 3'-terminus. Acridine is an effective intercalating agent.

<i>Item</i>	<i>Cat. No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Acridine Phosphoramidite	10-1973-95	50 µmole	165.00
	10-1973-90	100 µmole	295.00
	10-1973-02	0.25g	675.00
3'-Acridine CPG	20-2973-01	0.1g	120.00
	20-2973-10	1.0g	995.00
	1 µmole columns	Pack of 4	200.00
	0.2 µmole columns	Pack of 4	120.00
	10 µmole column (ABI)	Pack of 1	300.00
15 µmole cloumn (Expedite)	20-2973-14	Pack of 1	450.00

DNP LABELLING

An analytical test based on detection of 2,4-dinitrophenyl (DNP) labelled oligonucleotides with anti-DNP antibodies has been proposed. We have chosen the branched triethylene glycol (TEG) spacer in our version of DNP phosphoramidite since it can be added once or several times to the 3' or 5' terminus.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
DNP-TEG Phosphoramidite	10-1985-95	50 µmole	165.00
	10-1985-90	100 µmole	295.00
	10-1985-02	0.25g	675.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add


Expedite	E
MerMade	M

Columns


For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M


(Please inquire for availability of vials and columns for other instrument types.)



Acridine



Acridine CPG



DNP-TEG

CHOLESTEROL LABELLING

Potential therapeutic oligonucleotides must permeate the cell membrane for optimal activity. The addition of lipophilic groups to an oligonucleotide would be expected to enhance cellular uptake/membrane permeation. The use of cholesteryl oligos and the consequent improvement in activity has been described. We have designed our Cholesteryl products with triethyleneglycol (TEG) spacers for maximum solubility.

Item	Catalog No.	Pack	Price(\$)
Cholesteryl-TEG Phosphoramidite	10-1975-95	50 μ mole	140.00
	10-1975-90	100 μ mole	265.00
	10-1975-02	0.25g	545.00
5'-Cholesteryl-TEG Phosphoramidite	10-1976-95	50 μ mole	95.00
	10-1976-90	100 μ mole	175.00
	10-1976-02	0.25g	525.00
3'-Cholesteryl-TEG CPG	20-2975-01	0.1g	85.00
	20-2975-10	1.0g	700.00
	20-2975-41	Pack of 4	140.00
	20-2975-42	Pack of 4	84.00
	20-2975-13	Pack of 1	210.00
	20-2975-14	Pack of 1	315.00
1 μ mole columns			
0.2 μ mole columns			
10 μ mole column (ABI)			
15 μ mole column (Expedite)			

SEE ALSO

Spermine

p43

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

TOCOPHEROL LABELLING


Vitamin E is both lipophilic and non-toxic even at high doses so would be an excellent candidate as a lipophilic carrier for oligonucleotides. Therefore, as an addition to our cholesteryl product line, we offer simple α -tocopheryl (vitamin E) labelling. Totally synthetic α -tocopherol is racemic at its three chiral centers and is used to prepare this product.

Item	Catalog No.	Pack	Price(\$)
α -Tocopherol-TEG Phosphoramidite	10-1977-95	50 μ mole	160.00
	10-1977-90	100 μ mole	300.00
	10-1977-02	0.25g	575.00

STEARYL LABELLING

We now offer a simple C18 lipid as an economical and effective carrier molecule. We envisage that the 5'-stearyl group will become a favored lipophilic carrier for experimentation with synthetic oligonucleotides.


Item	Catalog No.	Pack	Price(\$)
5'- Stearyl Phosphoramidite	10-1979-90	100 μ mole	45.00
	10-1979-02	0.25g	180.00




PSORALEN LABELLING

Psoralen C2 at the 5'-terminus of an oligonucleotide serves effectively as a cross-linking reagent in double-stranded oligonucleotides. The 6 atom spacer arm of Psoralen C6 allows cross-linking with a triplex oligonucleotide strand. Click Chemistry with psoralen azide and one of our many nucleosidic and non-nucleosidic alkyne derivatives has the potential to generate a variety of practical cross-linkers. The well known reversible cross-linking behavior of psoralen with an adjacent thymidine residue could be very useful.


<i>Item</i>	<i>Cat. No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Psoralen C2 Phosphoramidite	10-1982-90	100 µmole	195.00
	10-1982-02	0.25g	495.00
Psoralen C6 Phosphoramidite	10-1983-90	100 µmole	195.00
	10-1983-02	0.25g	495.00
Psoralen Azide	50-2009-92	25 µmole	115.00
	50-2009-90	100 µmole	350.00



Psoralen C2



Psoralen C6



Psoralen Azide

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

EDTA LABELLING

EDTA-C2-dT phosphoramidite contains the triethyl ester of EDTA which allows sequence-specific cleavage of single- and double-stranded DNA and RNA. The cleavage reaction is only initiated once Fe(II) and dithiothreitol are added and so is readily controlled. Coupling of EDTA-dT is normal but cleavage and deprotection should be carried out with sodium hydroxide in aqueous methanol (0.4M NaOH in methanol/water 4:1) overnight at room temperature.

Item	Cat. No.	Pack	Price (\$)
EDTA-C2-dT-CE Phosphoramidite	10-1059-95	50 µmole	250.00
	10-1059-90	100 µmole	495.00
	10-1059-02	0.25g	975.00

FERROCENE LABELLING


With an excellent stability profile, ferrocene has always attracted considerable interest for DNA labelling to generate probes for electrochemical detection. Based on our Amino-Modifier C6-dT structure, Ferrocene-dT is easily added to oligonucleotides with no disruption of regular hybridization behavior. Multiple incorporations into an oligonucleotide probe are also simply achieved. Oligonucleotides are deprotected using standard techniques. Ferrocene oligonucleotides should be stored under Argon and aqueous solutions should be degassed immediately.

Item	Cat. No.	Pack	Price (\$)
Ferrocene-dT-CE Phosphoramidite	10-1576-95	50 µmole	170.00
	10-1576-90	100 µmole	330.00
	10-1576-02	0.25g	670.00

METHYLENE BLUE LABELLING

Dimethylaminophenothiazin-5-ium chloride, Methylene Blue (MB), has many uses in chemistry and biology. For decades, this compound has been used for applications including redox indicator, photosensitizer, dye for cellular staining procedures, antiseptic and in medicine against Alzheimer's disease. For the detection of biological analytes, MB has also been used as a redox reporter bound to DNA probes. Methylene Blue C3 Phosphoramidite is an asymmetric MB derivative specially designed for oligo synthesis. It must be used and deprotected with 0.05M potassium carbonate in methanol using UltraMild conditions. We also offer Methylene Blue NHS Ester to allow simple conjugation to the variety of amino-modifiers we supply.

Item	Cat. No.	Pack	Price (\$)
Methylene Blue C3 Phosphoramidite	10-5960-95	50 µmole	315.00
	10-5960-90	100 µmole	610.00
	10-5960-02	0.25g	1500.00
Methylene Blue NHS Ester (Dissolve 5.4mg in 60µL of DMSO)	50-1960-23	5.4mg	540.00



INTELLECTUAL PROPERTY

Methylene Blue C3 Phosphoramidite is covered by PCT application number WO2013128099 and is sold under license from the University of Lyon.

LABELLING WITH METAL CHELATES


2,2'-Dipicolylamine is a versatile metal-coordinating ligand capable of forming complexes with common metal ions including Zn²⁺, Ni²⁺, Cu²⁺, or Ag⁺. A tremendous advantage of dipicolylamine is complete compatibility with standard DNA synthesis, cleavage and purification protocols. Other chelating ligands may require nonstandard conditions or additional protection and deprotection steps. This product was manufactured and developed by Syntrix Biosystems Inc. Patents Pending. For Research Use Only.

Item	Cat. No.	Pack	Price (\$)
2,2'-Dipicolylamine Phosphoramidite	10-5801-95	50 µmole	105.00
	10-5801-90	100 µmole	200.00
	10-5801-02	0.25g	625.00


LABELLING WITH POLYAROMATIC HYDROCARBONS

Pyrene and **perylene** are fluorescent polycyclic aromatic hydrocarbons that have the ability to form 'excited state dimers' known as excimers. This unstructured, long-wavelength emission arises from the formation of a charge-transfer complex between the excited state and the ground state of two fluorescent molecules. In Pyrene-dU and perylene-dU, the hydrocarbon is attached at the 5 position of deoxyuridine through a triple bond and is electronically coupled to the deoxyuridine base. This electronic coupling of the base and the hydrocarbon makes the fluorescence sensitive to the base pairing of the dU portion of the molecule, allowing the discrimination between perfect and one base mismatched targets.


Item	Cat. No.	Pack	Price (\$)
Pyrene-dU-CE Phosphoramidite	10-1590-95	50 µmole	105.00
	10-1590-90	100 µmole	210.00
	10-1590-02	0.25g	550.00
Perylene-dU-CE Phosphoramidite	10-1591-95	50 µmole	150.00
	10-1591-90	100 µmole	300.00
	10-1591-02	0.25g	720.00



2,2'-Dipicolylamine



Pyrene-dU



Perylene-dU

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers
For Instrument type Add

Expedite	E
MerMade	M

Columns
For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

FLUORESCENT DYES

	Absorbance Maximum	Emission Maximum	Excimer
Pyrene-dU	402nm	472nm	486nm
Perylene-dU	473nm	490nm	Not Determined

PUROMYCIN CPG

One of the most challenging requirements associated with combinatorial chemistry is the recovery of sequence information of the oligonucleotide or peptide selected by the screening assay. A method¹ has been developed to generate a fusion product between mRNA and the polypeptide it encodes using *in vitro* translation of synthetic RNAs 3'-labeled with puromycin, an antibiotic that mimics transfer RNA. Puromycin binds in the ribosome's A site, forms a peptide bond with the growing peptide chain, and blocks further peptide elongation. By linking puromycin to mRNA, a peptide-RNA fusion product results from the translation of the message linking the encoding mRNA with its peptide product.

REFERENCE

(1) R.W. Roberts and J.W. Szostak, *Proc. Natl. Acad. Sci. USA*, 1997, **94**, 12297-302.

Item	Catalog No.	Pack	Price(\$)
Puromycin CPG	20-4040-01	0.1g	120.00
	20-4040-10	1.0g	995.00
1 μmole columns	20-4140-41	Pack of 4	200.00
0.2 μmole columns	20-4140-42	Pack of 4	120.00
10 μmole column (ABI)	20-4140-13	Pack of 1	360.00
15 μmole columns (Expedite)	20-4140-14	Pack of 1	540.00

QUENCHED AUTOLIGATION (QUAL) PROBES

QUAL probes¹ consist of two oligonucleotides, the first containing a nucleophilic group at the 3'-terminus, while the second has an electrophilic group at the 5'-terminus. When the probe pair finds the target, the oligos line up with the 3'-terminus of the first directly adjacent to the 5'-terminus of the second. An autoligation reaction then takes place to combine the two oligos into a single probe. As usual, the 3' nucleophilic group is the 3-thiophosphate, easily prepared using 3'-phosphate CPG with a sulfurizing step in the first cycle. In this case, the electrophilic group is a 5'-dabsyl group, which is an excellent leaving group as well as a fine quencher of fluorescence. The second oligo, therefore, contains a fluorophore which is quenched by the dabsyl group. A popular choice for fluorophore is fluorescein-dT but it is easy to imagine that a variety of fluorophores could be attached to any of the commercially available amino-modified nucleoside phosphoramidites.


REFERENCE

(1) S. Sando and E.T. Kool, *J Amer Chem Soc*, 2002, **124**, 2096-2097.


Item	Catalog No.	Pack	Price(\$)
5'-Dabsyl-dT-CE Phosphoramidite	10-1532-90	100 μmole	250.00
	10-1532-02	0.25g	775.00

SEE ALSO

3'-Phosphate CPG p78
Sulfurizing Reagents p37
Fluorescein-dT p96



Puromycin CPG



5'-Dabsyl-dT

LABELLING FOR PHOTO-REGULATION OF OLIGONUCLEOTIDES


Photo-control, the use of ultraviolet or visible light to control a reaction, has a number of advantages over other external stimuli:

- Light does not introduce contaminants into the reaction system,
- Excitation wavelength can be controlled through the design of the photo-responsive molecule, and
- It is now straightforward to control irradiation time and/or local excitation.

When a photo-responsive molecule is directly attached to DNA as a receptor, photo-regulation of the bioprocess regulated by that DNA molecule could, in principle, be achieved. Such photo-responsive DNA could also be used as a switch in a DNA-based nano-machine. Professor Hiroyuki Asanuma and his group at the department of Molecular Design and Engineering of the Graduate School of Engineering of the Nagoya University (Japan) have developed an efficient method to achieve this goal. They have attached azobenzene to DNA and made it photo-responsive^{1,2}. Azobenzene is a typical photo-responsive molecule that isomerizes from its planar *trans*-form to the non-planar *cis*-form after UV-light irradiation with a wavelength between 300 nm and 400 nm (λ_{max} is around 330 nm). Interestingly, the system reverts from the *cis*-form to the *trans*-form after further irradiation with visible light (wavelength over 400 nm). This process is completely reversible, and the azobenzene group does not decompose or induce undesirable side reactions even on repeated *trans*-*cis* isomerization. By introducing azobenzenes into DNA through D-threoninol as a linker, Asanuma and co-workers succeeded in achieving photo-regulation of:

- Formation and dissociation of a DNA duplex^{3,4} and
- Transcription by T7-RNA polymerase reaction^{5,6,7}.

Item	Catalog No.	Pack	Price(\$)
Azobenzene Phosphoramidite	10-5800-95	50 μ mole	105.00
	10-5800-90	100 μ mole	200.00
	10-5800-02	0.25g	550.00



Azobenzene Phosphoramidite

REFERENCES

- (1) H. Asanuma, et al., *Angew Chem Int Ed*, 2001, **40**, 2671-2673.
- (2) T. Takarada, et al., *Chem Lett.*, 2001, **30**, 732.
- (3) H. Asanuma, X.G. Liang, T. Yoshida, and M. Komiyama, *Chembiochem*, 2001, **2**, 39-44.
- (4) H. Asanuma, D. Matsunaga, and M. Komiyama, *NUCLEIC ACIDS SYMP SER (OXF)*, 2005, **49**, 35.
- (5) H. Asanuma, et al., *Chembiochem*, 2002, **3**, 786.
- (6) M. Liu, H. Asanuma, and M. Komiyama, *J. Amer. Chem. Soc.*, 2006, **128**, 1009.
- (7) H. Asanuma, et al., *Nature Protocols*, 2007, **2**, 203-212.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type	Add
Expedite	E
MerMade	M

Columns

For Instrument type	Add
Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)




RNA SUPPORTS FOR 3' MODIFICATION

Glen Research offers RNA supports in which protected ribonucleosides are attached to CPG. With 5'-DMT protection, and all other protecting groups base-labile, the use of these supports is identical to DNA supports. These supports are suitable for use in producing oligodeoxynucleotides modified at the 3'-terminus or oligoribonucleotides. ABI-style columns are supplied unless otherwise requested (see note box).

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>	
Bz-A-RNA-CPG	20-3303-01	0.1g	40.00	
	20-3303-02	0.25g	95.00	
	20-3303-10	1.0g	355.00	
	1 μ mole columns	20-3403-41	Pack of 4	100.00
	0.2 μ mole columns	20-3403-42	Pack of 4	75.00
	10 μ mole columns (ABI)	20-3403-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3403-14	Pack of 1	300.00
Ac-C-RNA-CPG	20-3315-01	0.1g	40.00	
	20-3315-02	0.25g	95.00	
	20-3315-10	1.0g	355.00	
	1 μ mole columns	20-3415-41	Pack of 4	100.00
	0.2 μ mole columns	20-3415-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3415-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3415-14	Pack of 1	300.00
Ac-G-RNA-CPG	20-3324-01	0.1g	40.00	
	20-3324-02	0.25g	95.00	
	20-3324-10	1.0g	355.00	
	1 μ mole columns	20-3424-41	Pack of 4	100.00
	0.2 μ mole columns	20-3424-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3424-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3424-14	Pack of 1	300.00
U-RNA-CPG	20-3330-01	0.1g	40.00	
	20-3330-02	0.25g	95.00	
	20-3330-10	1.0g	355.00	
	1 μ mole columns	20-3430-41	Pack of 4	100.00
	0.2 μ mole columns	20-3430-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3430-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3430-14	Pack of 1	300.00

ABBREVIATIONS

Ac = Acetyl
 Bz = Benzoyl
 CNEt = Cyanoethyl
 CPG = Controlled Pore Glass
 DMT = 4,4'-Dimethoxytrityl




TOM-PROTECTED RNA PHOSPHoramIDITES

RNA synthesis using monomers containing the 2'-O-TriisopropylsilyloxyMethyl (TOM) group (TOM-Protecting-Group™) is characterized by very high coupling efficiency along with fast, simple deprotection. High coupling efficiency is achieved because the TOM-Protecting-Group exhibits lower steric hindrance than the 2'-O-t-butyl dimethylsilyl (TBDMS) group used in our alternative RNA monomers. Fast and reliable deprotection is achieved using methylamine in ethanol/water at room temperature. A further feature of the TOM-Protecting-Group is that during basic steps it can not undergo 2' to 3' migration. This migration under basic conditions leads to non-biologically active 2'-5' linkages when using the TBDMS group. These features allow the TOM-Protected monomers to produce longer oligonucleotides. TOM-Protected RNA monomers are also fully compatible with minor bases with 2'-O-TBDMS protection.


Item	Catalog No.	Pack	Price (\$)
A-TOM-CE Phosphoramidite	10-3004-02	0.25g	75.00
	10-3004-05	0.5g	150.00
	10-3004-10	1.0g	275.00
C-TOM-CE Phosphoramidite	10-3014-02	0.25g	75.00
	10-3014-05	0.5g	150.00
	10-3014-10	1.0g	275.00
G-TOM-CE Phosphoramidite	10-3024-02	0.25g	75.00
	10-3024-05	0.5g	150.00
	10-3024-10	1.0g	275.00
U-TOM-CE Phosphoramidite	10-3034-02	0.25g	75.00
	10-3034-05	0.5g	150.00
	10-3034-10	1.0g	275.00

RNA SUPPORTS FOR TOM RNA SYNTHESIS


Item	Catalog No.	Pack	Price (\$)	
Ac-A-RNA-CPG	20-3304-01	0.1g	40.00	
	20-3304-02	0.25g	95.00	
	20-3304-10	1.0g	355.00	
	1 µmole columns	20-3404-41	Pack of 4	100.00
	0.2 µmole columns	20-3404-42	Pack of 4	75.00
10 µmole column (ABI)	20-3404-13	Pack of 1	225.00	
15 µmole column (Expedite)	20-3404-14	Pack of 1	300.00	




A-TOM



C-TOM




G-TOM



U-TOM

INTELLECTUAL PROPERTY



TOM-Protecting-Group™

TOM-RNA Phosphoramidites are supplied under agreement with QIAGEN. RNA synthesis using the TOM-Protecting-Group is covered by US Patent No. 5,986,084.

TOM-Protecting-Group is a trademark of QIAGEN.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

RNA SUPPORTS FOR TOM RNA SYNTHESIS (CONT.)

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>	
Ac-C-RNA-CPG	20-3315-01	0.1g	40.00	
	20-3315-02	0.25g	95.00	
	20-3315-10	1.0g	355.00	
	1 μ mole columns	20-3415-41	Pack of 4	100.00
	0.2 μ mole columns	20-3415-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3415-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3415-14	Pack of 1	300.00
Ac-G-RNA-CPG	20-3324-01	0.1g	40.00	
	20-3324-02	0.25g	95.00	
	20-3324-10	1.0g	355.00	
	1 μ mole columns	20-3424-41	Pack of 4	100.00
	0.2 μ mole columns	20-3424-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3424-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3424-14	Pack of 1	300.00
U-RNA-CPG	20-3330-01	0.1g	40.00	
	20-3330-02	0.25g	95.00	
	20-3330-10	1.0g	355.00	
	1 μ mole columns	20-3430-41	Pack of 4	100.00
	0.2 μ mole columns	20-3430-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3430-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3430-14	Pack of 1	300.00

TBDMS-PROTECTED RNA PHOSPHoramIDITES

Glen Research CE (β -cyanoethyl) Phosphoramidites for RNA synthesis are produced and packaged to ensure the highest performance on commercial synthesizers. Every batch is accompanied by a Certificate of Analysis and an HPLC trace, showing the results of our QC testing. RNA Phosphoramidites are synthesis-tested with a minimum coupling efficiency of 97%. Glen Research RNA monomers are packaged in industry standard vials which are specially cleaned to eliminate particulate contamination. These monomers are available in a variety of packs, including high throughput (HT) and low cost (LC). An UltraMild set is also available for situations where sensitive bases are in use. Dmf-G (10-3029) has been discontinued and may be substituted with Ac-G (10-3025).


Item	Catalog No.	Pack	Price (\$)
Bz-A-CE Phosphoramidite	10-3003-02	0.25g	40.00
	10-3003-05	0.5g	80.00
	10-3003-10	1.0g	160.00
Ac-C-CE Phosphoramidite	10-3015-02	0.25g	40.00
	10-3015-05	0.5g	80.00
	10-3015-10	1.0g	160.00
Ac-G-CE Phosphoramidite	10-3025-02	0.25g	40.00
	10-3025-05	0.5g	80.00
	10-3025-10	1.0g	160.00
U-CE Phosphoramidite	10-3030-02	0.25g	40.00
	10-3030-05	0.5g	80.00
	10-3030-10	1.0g	160.00

HT RNA PHOSPHoramIDITES


Bz-A-CE Phosphoramidite	10-3003-2HT	2.0g
Ac-C-CE Phosphoramidite	10-3015-2HT	2.0g
Ac-G-CE Phosphoramidite	10-3025-2HT	2.0g
U-CE Phosphoramidite	10-3030-2HT	2.0g

LC RNA PHOSPHoramIDITES


Bz-A-CE Phosphoramidite	10-3003-10	1.0g
Ac-C-CE Phosphoramidite	10-3015-10	1.0g
Ac-G-CE Phosphoramidite	10-3025-10	1.0g
U-CE Phosphoramidite	10-3030-10	1.0g




Bz-A-CE Phosphoramidite



Ac-C-CE Phosphoramidite



Ac-G-CE Phosphoramidite



U-CE Phosphoramidite

ABBREVIATIONS

Bz = Benzoyl
 CNEt = Cyanoethyl
 CPG = Controlled Pore Glass
 dmf = Dimethylformamide
 DMT = 4,4'-Dimethoxytrityl
 iPr = Isopropyl
 lcaa = long chain alkylamino
 Pac = Phenoxyacetyl
 PhOAc = Phenoxyacetyl
 TBDMS = t-Butyl-dimethylsilyl

HT RNA

Minimum order: 40g total
 Products may be combined to 40g total. Please request a quote.

LC RNA

Minimum order: 20g total
 Products may be combined to 20g total. Please request a quote.


ULTRAMILD TBDMS RNA PHOSPHoramIDITES

Item	Catalog No.	Pack	Price (\$)
Pac-A-CE Phosphoramidite	10-3000-02	0.25g	75.00
	10-3000-05	0.5g	150.00
	10-3000-10	1.0g	275.00
Ac-C-CE Phosphoramidite	10-3015-02	0.25g	40.00
	10-3015-05	0.5g	80.00
	10-3015-10	1.0g	160.00
iPr-Pac-G-CE Phosphoramidite	10-3021-02	0.25g	75.00
	10-3021-05	0.5g	150.00
	10-3021-10	1.0g	275.00
U-CE Phosphoramidite	10-3030-02	0.25g	40.00
	10-3030-05	0.5g	80.00
	10-3030-10	1.0g	160.00


TBDMS RNA SUPPORTS

ABI-style columns are supplied for 1 μ mole and 0.2 μ mole scales unless otherwise requested (see note box).


Item	Catalog No.	Pack	Price (\$)	
Pac-A-RNA-CPG	20-3300-01	0.1g	40.00	
	20-3300-02	0.25g	95.00	
	20-3300-10	1.0g	355.00	
	1 μ mole columns	20-3400-41	Pack of 4	100.00
	0.2 μ mole columns	20-3400-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3400-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3400-14	Pack of 1	300.00
Bz-A-RNA-CPG	20-3303-01	0.1g	40.00	
	20-3303-02	0.25g	95.00	
	20-3303-10	1.0g	355.00	
	1 μ mole columns	20-3403-41	Pack of 4	100.00
	0.2 μ mole columns	20-3403-42	Pack of 4	75.00
	10 μ mole column (ABI)	20-3403-13	Pack of 1	225.00
	15 μ mole column (Expedite)	20-3403-14	Pack of 1	300.00




Pac-A-CE Phosphoramidite



Ac-C-CE Phosphoramidite



iPr-Pac-G-CE Phosphoramidite



U-CE Phosphoramidite

TBDMS RNA SUPPORTS (CONT.)

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
Ac-C-RNA-CPG	20-3315-01	0.1g	40.00
	20-3315-02	0.25g	95.00
	20-3315-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
10 μ mole column (ABI)	20-3415-13	Pack of 1	225.00
15 μ mole column (Expedite)	20-3415-14	Pack of 1	300.00
iPr-Pac-G-RNA-CPG	20-3321-01	0.1g	40.00
	20-3321-02	0.25g	95.00
	20-3321-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
10 μ mole column (ABI)	20-3421-13	Pack of 1	225.00
15 μ mole column (Expedite)	20-3421-14	Pack of 1	300.00
Ac-G-RNA-CPG	20-3324-01	0.1g	40.00
	20-3324-02	0.25g	95.00
	20-3324-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
10 μ mole column (ABI)	20-3424-13	Pack of 1	225.00
15 μ mole column (Expedite)	20-3424-14	Pack of 1	300.00
U-RNA-CPG	20-3330-01	0.1g	40.00
	20-3330-02	0.25g	95.00
	20-3330-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
10 μ mole column (ABI)	20-3430-13	Pack of 1	225.00
15 μ mole column (Expedite)	20-3430-14	Pack of 1	300.00

ULTRAMILD SOLVENTS/REAGENTS

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Cap Mix A</i> THF/Pyridine/Pac ₂ O (Applied Biosystems)	40-4210-52	200mL	140.00
	40-4210-57	450mL	300.00
THF/Pac ₂ O (Expedite)	40-4212-52	200mL	140.00
	40-4212-57	450mL	300.00
<i>Deprotection Solution</i> 0.05M Potassium Carbonate in Methanol	60-4600-30	30mL	30.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

MINOR RNA PHOSPHoramIDITES (TOM PROTECTED)

SEE ALSO

Minor TBDMS monomers
Pyrrolo-CTP

p122
p124


Glen Research offers minor RNA phosphoramidites with either TOM or TBDMS protecting groups. 4-Thio-U, 5-Methyl-Cytidine, and 2-Amino-Adenosine are useful for analyzing RNA structure and activity relationships, for example, in ribozyme studies.

Pyrrolo-C is a fluorescent nucleoside whose fluorescence is sensitive to its environment and is ideal for probing RNA structure. It base-pairs as a normal C nucleotide. It is highly fluorescent and its excitation and emission are well to the red of most fluorescent nucleotide analogs, which eliminates or reduces background fluorescence from proteins. Pyrrolo-CTP has potential uses in biological assay development.


rSpacer is used to introduce an abasic site to an RNA sequence.

The protecting scheme for 2,6-Diaminopurine has been changed and the original product (10-3084) has been replaced with the optimized product (10-3085) below.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
4-Thio-U-TOM-CE Phosphoramidite	10-3052-95	50 μ mole	212.50
	10-3052-90	100 μ mole	425.00
	10-3052-02	0.25g	975.00
5-Me-C-TOM-CE Phosphoramidite	10-3064-95	50 μ mole	95.00
	10-3064-90	100 μ mole	190.00
	10-3064-02	0.25g	475.00
2,6-Diaminopurine-TOM-CE Phosphoramidite (2-amino-A)	10-3085-95	50 μ mole	212.50
	10-3085-90	100 μ mole	425.00
	10-3085-02	0.25g	975.00



4-Thio-U-TOM



5-Me-C-TOM



2,6-diaminopurine-TOM

MINOR RNA PHOSPHoramIDITES (TOM PROTECTED) (CONT.)

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
Pyrrolo-C-TOM-CE Phosphoramidite	10-3017-95	50 μ mole	212.50
	10-3017-90	100 μ mole	425.00
	10-3017-02	0.25g	975.00
rSpacer CE Phosphoramidite	10-3914-95	50 μ mole	90.00
	10-3914-90	100 μ mole	180.00
	10-3914-02	0.25g	495.00

RNA SEQUENCE MODIFIER (TOM PROTECTED)

Amino-Modifier C6-U has been added to the growing family of sequence modifiers and we envisage applications in RNA structural studies as well as for labelling siRNA to probe uptake and cellular distribution.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
Amino-Modifier C6-U Phosphoramidite	10-3039-95	50 μ mole	360.00
	10-3039-90	100 μ mole	720.00
	10-3039-02	0.25g	1475.00

SEE ALSO

Pyrrolo-dC p64
Pyrrolo-CTP p124

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type *Add*


Expedite E
 MerMade M

Columns

For Instrument type *Add*

Expedite E
 Applied Biosystems 3900 A
 MerMade M

(Please inquire for availability of vials and columns for other instrument types.)



MINOR RNA PHOSPHoramidites (TBDMS PROTECTED)

SEE ALSO


Minor TOM monomers p120

REFERENCES


- (1) C.J. Adams, J.B. Murray, M.A. Farrow, J.R.P. Arnold, and P.G. Stockley, *Tetrahedron Lett.*, 1995, **36**, 5421-5424.
 (2) D.A. Berry, et al., *Tetrahedron Lett.*, 2004, **45**, 2457-2461.

Inosine and 5-Methyl-Uridine are useful for analyzing RNA structure and activity relationships. 5-Bromo-Uridine and 5-Iodo-Uridine have been used for crystallography studies and cross-linking experiments. 6-Thioguanosine (6-thio-G) has applications in ribozyme and siRNA research, as well as in RNA-protein interactions. The removal of the silyl protecting group without interfering with the sulfur is critical. This is removed cleanly by triethylamine trihydrofluoride in DMSO but t-butylammonium fluoride (TBAF) leads to degradation of the thio-nucleotide analogue and should not be used. 2-Aminopurine riboside is useful for analyzing RNA structure and activity relationships, for example, in ribozyme studies.


Item	Catalog No.	Pack	Price(\$)
I-CE Phosphoramidite	10-3040-95	50 μ mole	95.00
	10-3040-90	100 μ mole	190.00
	10-3040-02	0.25g	475.00
5-Me-U-CE Phosphoramidite (T)	10-3050-95	50 μ mole	95.00
	10-3050-90	100 μ mole	190.00
	10-3050-02	0.25g	475.00
Br-U-CE Phosphoramidite	10-3090-95	50 μ mole	98.00
	10-3090-90	100 μ mole	195.00
	10-3090-02	0.25g	475.00
I-U-CE Phosphoramidite	10-3091-95	50 μ mole	98.00
	10-3091-90	100 μ mole	195.00
	10-3091-02	0.25g	475.00
6-Thio-G-CE Phosphoramidite	10-3072-95	50 μ mole	250.00
	10-3072-90	100 μ mole	500.00
	10-3072-02	0.25g	1200.00
2-Aminopurine-CE Phosphoramidite	10-3070-95	50 μ mole	212.50
	10-3070-90	100 μ mole	425.00
	10-3070-02	0.25g	975.00




Inosine




5-Me-U




Br-U



I-U



6-Thio-G



2-Aminopurine







MINOR RNA (TBDMS PROTECTED) (CONT.)

7-Deaza-Adenosine is lacking nitrogen at the 7-position, which is replaced by carbon. The N7 position in adenosine takes part in non-Watson and Crick hydrogen bonding, which may be relevant to RNA folding and subsequent activity. This Adenosine analogue is also known as Tubercidin. 8-Aza-7-deaza-Adenosine is an isomer of Adenosine with virtually identical electron density. Again, the N7 nitrogen is not available for hydrogen bonding. Nebularine or Purine Nucleoside can be viewed as an Adenosine derivative that is lacking the exocyclic amino group. This molecule allows researchers to determine the relevance of the exocyclic amine of Adenosine to RNA structure and function.

Ribozyme activity is substantially affected by the substitution of modified pyrimidine bases. Zebularine (pyrimidin-2-one ribonucleoside) may be regarded as a Cytidine derivative lacking the exocyclic amino group. Zebularine and Pyridin-2-one Ribonucleoside, the 3-deaza analogue of Zebularine, are prime candidates for use in evaluating ribozyme activity and function. It should be noted that Zebularine is mildly fluorescent, absorbing at 298nm and emitting at 367nm.

PseudoUridine is one of the most common modified nucleosides found in RNA. The availability of a phosphoramidite will allow detailed research into the effects of this modified base on RNA structure and activity.

Item	Catalog No.	Pack	Price(\$)
7-Deaza-A-CE Phosphoramidite (Discontinued)	10-3001		
Zebularine-CE Phosphoramidite	10-3011-95	50 µmole	125.00
	10-3011-90	100 µmole	250.00
	10-3011-02	0.25g	650.00
Pyridin-2-one-CE Phosphoramidite	10-3012-95	50 µmole	210.00
	10-3012-90	100 µmole	420.00
	10-3012-02	0.25g	1200.00
Nebularine-CE Phosphoramidite (Purine Ribonucleoside) (Discontinued)	10-3041		
PseudoUridine-CE Phosphoramidite	10-3055-95	50 µmole	175.00
	10-3055-90	100 µmole	350.00
	10-3055-02	0.25g	995.00
8-Aza-7-deaza-A-CE Phosphoramidite	10-3083-95	50 µmole	300.00
	10-3083-90	100 µmole	600.00
	10-3083-02	0.25g	1500.00



OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers For Instrument type Add

Expedite	E
MerMade	M

Columns For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

MINOR RNA (TBDMS PROTECTED) (CONT.)

Methylation of adenosine at position 1 produces a drastic functional change in the nucleobase. 1-Methyladenosine (pK_a 8.25) is a much stronger base than adenosine (pK_a 3.5). N-1 methylation excludes participation of the adenine base in canonical Watson-Crick base pairing and provides a positive charge to the nucleobase. This modification also alters the hydrophobicity of the base, the stacking properties, the ordering of water molecules and the chelation properties. The base may become involved in non-canonical hydrogen bonding, in electrostatic interactions and, in general, it may contribute to the conformational dynamics of the tRNA.


In the central dogma of molecular biology, genetic information flows from DNA to RNA and then to protein. Reversible epigenetic modifications on genomic DNA and histone have been known to substantially regulate gene expression. On the other hand, there exists more than 100 naturally occurring chemical modifications in RNA; however, the functions of these RNA modifications are largely unknown. Whether some of these modifications in RNA can be reversed and could impact gene expression in the central dogma was unknown until the recent discovery of N6-methyladenosine (N6-Me-A) as the first example of reversible RNA methylation.¹ We offer the N6-Me-A RNA monomer with a phenoxyacetyl protecting group to minimize potential branching. We have shown N6-Me-A-CE Phosphoramidite to be completely compatible with all popular RNA synthesis and deprotection methods, from UltraMild to the most popular procedure using AMA for deprotection.

Item	Catalog No.	Pack	Price(\$)
1-Me-A-CE Phosphoramidite	10-3501-95	50 μ mole	190.00
	10-3501-90	100 μ mole	380.00
	10-3501-02	0.25g	975.00
N6-Me-A-CE Phosphoramidite	10-3005-95	50 μ mole	285.00
	10-3005-90	100 μ mole	550.00
	10-3005-02	0.25g	1295.00


MINOR RNA TRIPHOSPHATES

Pyrrolo-dC is a fluorescent nucleoside that codes as dC and base pairs efficiently with dG. Preliminary evidence indicates that pyrrolo-dC triphosphate is an excellent substrate for Taq, Pfu and Vent polymerases and is incorporated specifically opposite dG. Pyrrolo-dCTP has been available for some time and is in use in biological assays. Pyrrolo-CTP is a fluorescent ribonucleotide with fluorescence exquisitely sensitive to its environment and is of great interest for RNA structural research. The pyrrolo-C project is a joint development by Berry and Associates, Inc. and Glen Research Corporation.


Item	Catalog No.	Pack	Price(\$)
Pyrrolo-CTP 10mM	81-3017-01	100 μ L	270.00



1-Me-A



N6-Me-A



Pyrrolo-CTP

REFERENCE

- (1) Y. Fu, D. Dominissini, G. Rechavi, and C. He, *Nat Rev Genet*, 2014, **15**, 293-306.

SEE ALSO

Pyrrolo-dC
Pyrrolo-C

p64
p121


2'-OME-RNA PHOSPHoramIDITES

Glen Research 2'-OMe-RNA CE (β -cyanoethyl) Phosphoramidites are designed to produce synthetic oligonucleotides containing nuclease resistant 2'-O-methyl ribonucleotide linkages. Deprotection, isolation and handling of 2'-O-methyl oligonucleotides are identical to the procedures for oligodeoxynucleotides.


Item	Catalog No.	Pack	Price(\$)
2'-OMe-A-CE Phosphoramidite	10-3100-90	100 μ mole	20.00
	10-3100-02	0.25g	50.00
	10-3100-05	0.5g	100.00
	10-3100-10	1.0g	200.00
2'-OMe-C-CE Phosphoramidite	10-3110-90	100 μ mole	20.00
	10-3110-02	0.25g	50.00
	10-3110-05	0.5g	100.00
	10-3110-10	1.0g	200.00
2'-OMe-Ac-C-CE Phosphoramidite	10-3115-90	100 μ mole	20.00
	10-3115-02	0.25g	50.00
	10-3115-05	0.5g	100.00
	10-3115-10	1.0g	200.00
2'-OMe-iBu-G-CE Phosphoramidite	10-3120-90	100 μ mole	20.00
	10-3120-02	0.25g	50.00
	10-3120-05	0.5g	100.00
	10-3120-10	1.0g	200.00
2'-OMe-G-CE Phosphoramidite	10-3121-90	100 μ mole	20.00
	10-3121-02	0.25g	50.00
	10-3121-05	0.5g	100.00
	10-3121-10	1.0g	200.00
2'-OMe-U-CE Phosphoramidite	10-3130-90	100 μ mole	20.00
	10-3130-02	0.25g	50.00
	10-3130-05	0.5g	100.00
	10-3130-10	1.0g	200.00

HT 2'-OME-RNA PHOSPHoramIDITES


2'-OMe-Bz-A-CE Phosphoramidite	10-3100-2HT	2.0g
2'-OMe-Ac-C-CE Phosphoramidite	10-3115-2HT	2.0g
2'-OMe-iBu-G-CE Phosphoramidite	10-3120-2HT	2.0g
2'-OMe-U-CE Phosphoramidite	10-3130-2HT	2.0g




2'-OMe-A




2'-OMe-C




2'-OMe-Ac-C



2'-OMe-iBu-G



2'-OMe-G



2'-OMe-U

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

HT 2'-OMe-RNA

Minimum order: 40g total
Products may be combined to 40g total. Please request a quote.

ULTRAMILD 2'-OME-RNA

The use of UltraMild monomers in oligonucleotide synthesis has allowed very sensitive dyes like TAMRA, HEX and Cy5 to be used virtually routinely. The DNA and RNA monomers are currently available and we also provide this set of 2'-OMe-RNA monomers. In our version of this chemistry, we use as protecting groups phenoxyacetyl (Pac) for A, acetyl (Ac) for C, and isopropyl-phenoxyacetyl (iPr-Pac) for G.

It has become clear that acetic anhydride in the conventional capping mix can cause transamidation in situations where an amine protecting group is quite labile. This leads to acetyl protection on the amino group that may be slow to be removed. Consequently, if many dG residues are included in the oligonucleotide, we recommend the use of phenoxyacetic anhydride (Pac₂O) in Cap A. This modification removes the possibility of exchange of the iPr-Pac protecting group on the dG with acetate from the acetic anhydride capping mix.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
2'-OMe-Pac-A-CE Phosphoramidite	10-3601-02	0.25g	62.50
	10-3601-05	0.5g	125.00
	10-3601-10	1.0g	250.00
2'-OMe-Ac-C-CE Phosphoramidite	10-3115-02	0.25g	50.00
	10-3115-05	0.5g	100.00
	10-3115-10	1.0g	200.00
2'-OMe-iPr-Pac-G-CE Phosphoramidite	10-3621-02	0.25g	62.50
	10-3621-05	0.5g	125.00
	10-3621-10	1.0g	250.00

ULTRAMILD SOLVENTS/REAGENTS




Cap Mix A

THF/Pyridine/Pac ₂ O	40-4210-52	200mL	140.00
(Applied Biosystems)	40-4210-57	450mL	300.00

THF/Pac ₂ O	40-4212-52	200mL	140.00
(Expedite)	40-4212-57	450mL	300.00

Deprotection Solution

0.05M Potassium Carbonate in Methanol	60-4600-30	30mL	30.00
---------------------------------------	------------	------	-------



2'-OME-RNA SUPPORTS

ABI-style columns are supplied for 1 μ mole and 0.2 μ mole scales unless otherwise requested (see note box).

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
2'-OMe-A-RNA-CPG	20-3600-01	0.1g	40.00
	20-3600-02	0.25g	95.00
	20-3600-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
	10 μ mole column (ABI)	Pack of 1	225.00
	15 μ mole column (Expedite)	Pack of 1	300.00
2'-OMe-C-RNA-CPG	20-3610-01	0.1g	40.00
	20-3610-02	0.25g	95.00
	20-3610-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
	10 μ mole column (ABI)	Pack of 1	225.00
	15 μ mole column (Expedite)	Pack of 1	300.00
2'-OMe-Ac-C-RNA-CPG	20-3615-01	0.1g	40.00
	20-3615-02	0.25g	95.00
	20-3615-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
	10 μ mole column (ABI)	Pack of 1	225.00
	15 μ mole column (Expedite)	Pack of 1	300.00
2'-OMe-G-RNA-CPG	20-3621-01	0.1g	40.00
	20-3621-02	0.25g	95.00
	20-3621-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
	10 μ mole column (ABI)	Pack of 1	225.00
	15 μ mole column (Expedite)	Pack of 1	300.00
2'-OMe-U-RNA-CPG	20-3630-01	0.1g	40.00
	20-3630-02	0.25g	95.00
	20-3630-10	1.0g	355.00
	1 μ mole columns	Pack of 4	100.00
	0.2 μ mole columns	Pack of 4	75.00
	10 μ mole column (ABI)	Pack of 1	225.00
	15 μ mole column (Expedite)	Pack of 1	300.00

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type *Add*

Expedite	E
MerMade	M

Columns

For Instrument type *Add*

Expedite	E
Applied Biosystems 3900	A
MerMade	M


(Please inquire for availability of vials and columns for other instrument types.)

MINOR 2'-OME-RNA PHOSPHoramIDITES


To aid in the evaluation of the structures of 2'-OMe-RNA complexes, we offer the CE phosphoramidites listed below. 2'-OMe-T is useful in triplex studies while the 2-aminopurine derivative may be tested in ribozyme studies. By supporting an additional hydrogen bond, 2,6-diaminopurine (2-amino-adenosine) binds more strongly with uridine than does adenosine. Oligonucleotides containing 2'-OMe-5-Me-C and 2'-OMe-I would be of interest to researchers involved in triplex and antisense studies using 2'-OMe-RNA. The uses of 2'-OMe-5-bromo-U phosphoramidite range from crystallographic studies due to the heavy atom to cross-linking because of its photolability. 5-Fluoro-pyrimidine nucleosides have been useful as therapeutic agents and their effect on the structure and activity of oligonucleotides may be examined using the 2'-OMe-RNA derivatives. The 2,4,6-trimethylphenyl (TMP) protected 2'-OMe-U derivative is a convertible nucleoside and reaction with ammonia leads to the 5-fluoro-dC analogue. 2'-OMe-3-deaza-5-aza-C (Reverse C) derivative has the potential to mimic in oligonucleotides 5-azacytidine, a DNA methylase inhibitor. Its ability to bind as a C will likely be diminished.

ABI-style vials are supplied unless otherwise requested (see note box).


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
2'-OMe-2-Aminopurine- CE Phosphoramidite (N-dmf-AP)	10-3123-95	50 μ mole	177.50
	10-3123-90	100 μ mole	355.00
	10-3123-02	0.25g	975.00
2'-OMe-2,6-Diaminopurine- CE Phosphoramidite (2-amino-A)	10-3124-95	50 μ mole	177.50
	10-3124-90	100 μ mole	355.00
	10-3124-02	0.25g	975.00
2'-OMe-5-Me-U-CE Phosphoramidite (2'-OMe-T)	10-3131-90	100 μ mole	150.00
	10-3131-02	0.25g	360.00



2'-OMe-2-AP




2'-OMe-2-amino-A




2'-OMe-5-Me-U

MINOR 2'-OME-RNA PHOSPHoramidites (CONT.)


Item	Catalog No.	Pack	Price(\$)
2'-OMe-I-CE Phosphoramidite	10-3140-90	100 μ mole	150.00
	10-3140-02	0.25g	360.00
2'-OMe-5-Me-C-CE Phosphoramidite	10-3160-90	100 μ mole	240.00
	10-3160-02	0.25g	675.00
2'-OMe-5-Br-U-CE Phosphoramidite	10-3190-90	100 μ mole	240.00
	10-3190-02	0.25g	675.00
2'-OMe-TMP-5-F-U-CE Phosphoramidite (2'-OMe-5-F-C Precursor)	10-3111-95	50 μ mole	177.50
	10-3111-90	100 μ mole	355.00
	10-3111-02	0.25g	975.00
2'-OMe-5-F-U-CE Phosphoramidite	10-3132-95	50 μ mole	177.50
	10-3132-90	100 μ mole	355.00
	10-3132-02	0.25g	975.00
2'-OMe-3-deaza-5-aza-C-CE Phosphoramidite	10-3116-95	50 μ mole	177.50
	10-3116-90	100 μ mole	355.00
	10-3116-02	0.25g	975.00




2'-OMe-I




2'-OMe-5-Me-C




2'-OMe-5-Br-U



2'-OMe-TMP-5-F-U



2'-OMe-5-F-U



2'-OMe-3-deaza-5-aza-C

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add

Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

2'-F-RNA MONOMERS

2'-Deoxy-2'-fluoro-nucleosides adopt an RNA-type sugar conformation, presumably due to the high electronegativity of fluorine. Because of this sugar conformation, RNA duplexes (A-form) are generally more thermodynamically stable than DNA duplexes (B-form). As expected, the addition of 2'-F-RNA residues to oligodeoxynucleotides progressively increases the thermal stability of their duplexes with RNA. The stabilization is additive at approximately 2° per residue. This compares favorably with 2'-OMe-RNA at around 1.5° and RNA at 1.1° per residue. In the meantime, base pair specificity remains intact.

2'-F-RNA phosphodiester linkages are not nuclease resistant, although the corresponding phosphorothioate linkages are highly resistant. Researchers usually design antisense oligonucleotides to form duplexes with RNA, which are then substrates for RNase H. Uniformly modified 2'-F-RNA/RNA duplexes are not substrates for RNase H. However, it is straightforward to prepare chimeric 2'-F-RNA/DNA phosphorothioate oligonucleotides which exhibit enhanced binding to the RNA target, are substrates for RNase H, and are highly nuclease resistant.

STABILITY NOTE

Synthetic oligonucleotides containing 2'-F-RNA linkages may be deprotected with ammonium hydroxide as normal. Deprotection using AMA at 65°C leads to some degradation and so we recommend the use of AMA at room temperature for 2 hours.

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite E
MerMade M


Columns

For Instrument type Add


Expedite E
Applied Biosystems 3900 A
MerMade M

(Please inquire for availability of vials and columns for other instrument types.)


Item	Catalog No.	Pack	Price(\$)
2'-F-A-CE Phosphoramidite	10-3400-02	0.25g	100.00
	10-3400-05	0.5g	200.00
2'-F-Ac-C-CE Phosphoramidite	10-3415-02	0.25g	50.00
	10-3415-05	0.5g	100.00
2'-F-G-CE Phosphoramidite	10-3420-02	0.25g	100.00
	10-3420-05	0.5g	200.00
2'-F-U-CE Phosphoramidite	10-3430-02	0.25g	50.00
	10-3430-05	0.5g	100.00




2'-F-A



2'-F-Ac-C



2'-F-G



2'-F-U


2'-F-ARABINONUCLEIC ACID (2'-F-ANA)

Arabinonucleosides are epimers of ribonucleosides with the chiral switch being at the 2' position of the sugar residue. 2'-F-ANA adopts a more DNA-like B-type helix conformation, not through the typical C2'-endo conformation but, rather, through an unusual O4'-endo (east) pucker. However, the presence of the electronegative fluorine leads to a still significant increase (ΔT_m 1.2° C/mod) in melting temperature per modification.¹ 2'-F-ANA-containing oligonucleotides exhibit very high binding specificity to their targets. Indeed, a single mismatch in a 2'-F-ANA – RNA duplex leads to a ΔT_m of -7.2 °C and in a 2'-F-ANA – DNA duplex a ΔT_m of -3.9 °C.²


The presence of fluorine at the 2' position in 2'-F-ANA leads to increased stability to hydrolysis under basic conditions relative to RNA and even 2'-F-RNA.^{1,3} The stability of 2'-F-ANA to nucleases also makes this a useful modification for enhancing the stability of oligonucleotides in biological environments.² 2'-F-ANA hybridizes strongly to target RNA and, unlike most 2' modifications, induces cleavage of the target by RNase H. Phosphorothioate (PS) 2'-F-ANA is routinely used in these applications due to its increased nuclease resistance. Alternating 2'-F-ANA and DNA units provide among the highest potency RNase H-activating oligomers. Both the "altimer" and "gapmer" strand architectures consistently outperform PS-DNA and DNA/RNA gapmers.⁴

siRNA oligos were found to tolerate the presence of 2'-F-ANA linkages very well. High potency gene silencing was demonstrated⁵ with siRNA chimeras containing 2'-F-RNA and/or LNA and 2'-F-ANA. The high efficacy of these chimeras was attributed to the combination of the rigid RNA-like properties of 2'-F-RNA and LNA with the DNA-like properties of 2'-F-ANA.


Item	Catalog No.	Pack	Price(\$)
2'-F-A-ANA CE Phosphoramidite	10-3800-90	100 μ mole	150.00
	10-3800-02	0.25g	375.00
2'-F-Bz-C-ANA CE Phosphoramidite	10-3810-02	0.25g	200.00
	10-3810-05	0.5g	400.00
2'-F-G-ANA CE Phosphoramidite	10-3820-90	100 μ mole	165.00
	10-3820-02	0.25g	425.00
2'-F-U-ANA CE Phosphoramidite	10-3830-02	0.25g	125.00
	10-3830-05	0.5g	250.00




2'-F-A-ANA



2'-F-Bz-C-ANA



2'-F-G-ANA



2'-F-U-ANA

REFERENCES

1. E. Viazovkina, M.M. Mangos, M.I. Elzagheid, and M.J. Damha, *Curr Protoc Nucleic Acid Chem*, 2002, **Chapter 4**, Unit 4 15.
2. J.K. Watts, and M.J. Damha, *Can. J. Chem.*, 2008, **86**, 641-656.
3. J.K. Watts, A. Katolik, J. Viladoms, and M.J. Damha, *Org Biomol Chem*, 2009, **7**, 1904-10.
4. A. Kalota, *et al.*, *Nucleic Acids Res.*, 2006, **34**, 451.
5. G.F. Deleavey, *et al.*, *Nucleic Acids Res.*, 2010, **38**, 4547-4557, J.K. Watts, *et al.*, *Nucleic Acids Res.*, 2007, **35**, 1441-1451, T. Dowler, *et al.*, *Nucleic Acids Res.*, 2006, **34**, 1669-1675.

INTELLECTUAL PROPERTY


2'-F-ANA is covered by intellectual property. Key patents covering siRNA and antisense applications are as follows:

WO/2009/146556 (siRNA); WO 03064441 and WO 0220773 (antisense).


STABILITY NOTE

Synthetic oligonucleotides containing 2'-F-RNA linkages may be deprotected with ammonium hydroxide as normal. Deprotection using AMA at 65°C leads to some degradation and so we recommend the use of AMA at room temperature for 2 hours.


INTERFERENCE MAPPING

1. Incorporation of Phosphorothioate Tagged Patent Nucleotide (N) or Modified Analog (δ)


2. Selection of Functional RNAs



3. Cleavage of Phosphorothioate Linkage with Iodine



4. Electrophoresis and Autoradiography



Nucleotide Analog Interference Mapping (NAIM) is a chemogenetic approach that makes it possible to simultaneously, yet individually, probe the contribution of a particular functional group at almost every RNA nucleotide position in a single experiment¹. The method utilizes a series of 5'-O-(1-thio)nucleoside analog triphosphates in a modification interference procedure that is as simple as RNA sequencing. In a NAIM experiment the smallest mutable unit is not the base pair, but rather the individual functional groups that comprise the nucleotides. Because the modification or deletion of a particular functional group within an RNA can severely affect its activity, this approach makes it possible to efficiently determine the chemical basis of RNA structure and function.


Instead of synthesizing a series of RNAs with chemical substitutions at specific sites, NAIM utilizes a combinatorial approach. Each nucleotide analog is prepared as a triphosphate for incorporation into the RNA during DNA templated *in vitro* transcription. The nucleotide analogs used in NAIM include a specific chemical alteration to the base or sugar, and an α -phosphorothioate substitution which serves as a chemical tag. The nucleotide analog triphosphate is randomly incorporated into an RNA transcript, where the phosphorothioate linkage can be selectively cleaved by the addition of I_2 to produce a series of RNA cleavage products whose lengths correspond to the sites of analog incorporation². By radioactively or fluorescently tagging one end of the RNA transcript, cleaving the RNA with I_2 , and resolving the cleavage products on a denaturing polyacrylamide gel, the sites of analog incorporation throughout the RNA can be individually assayed and used for interference analysis. The phosphorothioate tagged nucleotide analogs make it possible for all of the positions in the RNA to be assayed individually for functional group modification in a single experiment.

Because the phosphorothioate chemical tag is independent of the nucleotide analog whose location it reports, NAIM is generalizable to any analog that can be incorporated into a transcript by an RNA polymerase. A typical NAIM experiment is comprised of four steps. (i) The phosphorothioate tagged nucleotide analog is randomly incorporated throughout the RNA to create a family of transcripts, each of which contains only a few substitutions. A different transcription reaction is performed for each analog. (ii) The functional RNA variants in the population are separated from the inactive transcripts. The exact nature of the activity assay is specific for the RNA being studied, but could include affinity chromatography, native gel electrophoresis, filter binding, selective radiolabeling, etc. (iii) The phosphorothioate linkages in the active and unselected RNA populations are cleaved by I_2 addition to mark the sites of analog incorporation within each molecule. (iv) The individual RNA fragments are resolved by gel electrophoresis and visualized by autoradiography. Sites of analog substitution that are detrimental to function are scored as gaps in the sequencing ladder among the active RNA variants. Because every position in the sequence is a unique and independent band on the sequencing gel, a single screen can define the effect a particular analog has at every incorporated position within the RNA. The approach is applicable to any RNA that can be transcribed *in vitro* and has an assayable function that can be used to distinguish active and inactive variants. RNA functions that are amenable to this approach include catalysis, folding, protein or ligand binding, and the ability to act as a reaction substrate.


A ANALOGS

NAIM utilizes α -phosphorothioate tagged nucleotide analogs, each of which includes an incremental chemical alteration in the base or ribose sugar. The most completely developed set of analogs are those of adenosine, for which eight different analogs have been utilized in NAIM.³ Five analogs modify the nucleotide base and three modify the ribose sugar. The base analogs include purine riboside (Pur α S), N6-methyladenosine (m⁶A α S), tubercidin (7dA α S), diaminopurine riboside (DAP α S), and 2-aminopurine riboside (2AP α S). The ribose sugar analogs all modify the 2'-OH group and include 2'-deoxyadenosine (dA α S), 2'-deoxy-2'-fluoroadenosine (fA α S), and 2'-O-methyladenosine (O^{me}A α S). All of the analogs can be randomly incorporated into an RNA transcript at an ideal 5% level of efficiency using either the wild-type T7 RNA polymerase or a Y639F RNA polymerase point mutant⁴. Each of these analogs provides specific information about the chemical basis of RNA activity at almost every incorporated position in the transcript.


Item	Catalog No.	Pack	Price(\$)
Adenosine α -thiotriphosphate (0.5mM)	80-3000-01	100 μ L	100.00
7-Deaza-Adenosine α -thiotriphosphate (1mM)	80-3303-01	100 μ L	100.00
N6-Me-Adenosine α -thiotriphosphate (4mM)	80-3302-01	100 μ L	100.00



Adenosine (A α S)



7-Deaza-Adenosine (7dA α S)



N6-Methyl-Adenosine (m⁶A α S)

REFERENCES

- (1) S. A. Strobel and K. Shetty, *Proc. Natl. Acad. Sci. U.S.A.*, 1997, **94**, 2903-2908.
- (2) G. Gish and F. Eckstein, *Science*, 1988, **240**, 1520-1522.
- (3) L. Ortoleva-Donnelly, A. A. Szewczak, R. R. Gutell and S. A. Strobel, *RNA*, 1998, **4**, 498-519.
- (4) R. Sousa and R. Padilla, *EMBO J.*, 1995, **14**, 4609-4621.


INTELLECTUAL PROPERTY

Products for Nucleotide Analog Interference Mapping (NAIM) are supplied under license.

A ANALOGS (CONT.)

Item	Catalog No.	Pack	Price(\$)
2,6-Diaminopurine riboside α -thiotriphosphate (0.25mM)	80-3305-01	100 μ L	100.00
2-Aminopurine riboside α -thiotriphosphate (10mM)	80-3304-01	100 μ L	100.00
Purine riboside α -thiotriphosphate (20mM)	80-3301-01	100 μ L	100.00
2'-deoxyAdenosine α -thiotriphosphate (15mM)	80-1000-01	100 μ L	100.00
2'-OMe-Adenosine α -thiotriphosphate (20mM)	80-1101-01	100 μ L	100.00
2'-Fluoro-Adenosine α -thiotriphosphate (10mM)	80-1102-01	100 μ L	100.00

α -Thiotriphosphates are sodium salts in TE buffer, pH7, 10X concentrates. The concentrations shown are optimal for incorporation during polymerase reactions.




NUCLEOSIDE α -THIOTRIPHOSPHATES


In addition to their applications in NAIM, there are many other uses for α -thiotriphosphates. So we have prepared, and will maintain, supplies of regular nucleoside and 2'-deoxy-nucleoside α -thiotriphosphates. These are currently offered in concentrations suitable for NAIM. Please inquire about specialized requirements.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
Cytidine α -thiotriphosphate (1.0mM)	80-3010-01	100 μ L	100.00
Guanosine α -thiotriphosphate (0.5mM)	80-3020-01	100 μ L	100.00
Uridine α -thiotriphosphate (0.5mM)	80-3040-01	100 μ L	100.00
5-Methyl-Uridine α -thiotriphosphate (10mM)	80-3093-01	100 μ L	100.00
Inosine α -thiotriphosphate (4mM)	80-3050-01	100 μ L	100.00
2'-deoxyCytidine α -thiotriphosphate (15mM)	80-1010-01	100 μ L	100.00
2'-deoxyGuanosine α -thiotriphosphate (5mM)	80-1020-01	100 μ L	100.00
Thymidine α -thiotriphosphate (15mM)	80-1030-01	100 μ L	100.00
2'-deoxyUridine α -thiotriphosphate (5mM)	80-1040-01	100 μ L	100.00


α -Thiotriphosphates are sodium salts in TE buffer, pH7, 10X concentrates. The concentrations shown are optimal for incorporation during polymerase reactions.




Cytidine




Guanosine




Uridine




5-Methyl Uridine




Inosine




2'-deoxyCytidine



2'-deoxyGuanosine




Thymidine




2'-deoxyUridine

NUCLEOSIDE α -THIOTRIPHOSPHATES (CONT.)


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
2'-OMe-Uridine α -thiotriphosphate (20mM)	80-1141-01	100 μ L	100.00
7-Deaza-Guanosine α -thiotriphosphate (1.0mM)	80-3321-01	100 μ L	100.00
Pseudo-Uridine α -thiotriphosphate (1.0mM)	80-3341-01	100 μ L	100.00



2'-OMe-Uridine



7-Deaza-Guanosine




Pseudo-Uridine

INTERNALLY QUENCHED NUCLEOTIDE FLUORESCENT REPORTERS

Several methods have been developed for enzymatic fluorescent labelling of nucleic acids. A dNTP analog can be used to incorporate a fluorophore by PCR, nick translation or random priming, either directly into DNA¹ or indirectly via a hapten such as biotin.² Though high incorporation efficiencies have been reported,³ all of these approaches require the separation of unincorporated label prior to downstream applications. A reagent called an Internally Quenched Nucleotide or IQN has been developed by Lawler Scientific, LLC. This reagent consists of a nucleoside triphosphate with a fluorescent reporter attached to the nucleobase and a quencher moiety attached to the gamma-phosphate. The nucleotide remains non-fluorescent until the quencher is enzymatically separated from the parent nucleotide. Since IQNs are non-fluorescent until incorporated into a nucleic acid, they do not give rise to the background fluorescence signals commonly observed when DNA labelled by standard means is inadequately purified.

The first generation IQN consists of a fluorescein-dUTP with a dabsyl quencher linked to the gamma phosphate. Fluorescein and dabsyl were selected because of their superior optical properties and because the photophysics governing their interaction is well described in the literature.⁴ In addition, this IQN is soluble and stable in aqueous solution.

Item	Catalog No.	Pack	Price(\$)
Fluorescein-dUTP-dabsyl 1mM, 25nmoles, 10mM Tris, 1mM EDTA	88-1056-01	25 µL	270.00



Fluorescein-dUTP-dabsyl

INTELLECTUAL PROPERTY

These products are subject to proprietary rights of Lawler Scientific, LLC and are made and sold under license from Lawler Scientific, LLC. There is no implied license for commercial use with respect to the Products and a license must be obtained directly from Lawler Scientific, LLC with respect to any proposed commercial use of the Products.

REFERENCES

- (1) H. Yu, J. Chao, D. Patek, R. Mujumdar, S. Mujumdar, and A.S. Waggoner, *Nucleic Acids Res.*, 1994, **22**, 3226-32.
- (2) X. Li, W.M. James, F. Traganos, and Z. Darzynkiewicz, *Biotech Histochem*, 1995, **70**, 234-42.
- (3) T. Tasara, et al., *Nucleic Acids Res.*, 2003, **31**, 2636-46.
- (4) S.A.E. Marras, F.R. Kramer, and S. Tyagi, *Nucleic Acids Res.*, 2002, **30**, E122.

GLEN-PAK™ PURIFICATION

Glen-Pak™ DNA and RNA cartridges have advantages over Poly-Pak cartridges in that a single loading of the diluted crude deprotection solution is all that is necessary. Also, the range of purification has been extended to 100+ using DMT-on oligos. Glen-Pak cartridges have similar performance to Fluoro-Pak cartridges but without the need for the fluororous DMT group at the 5' terminus and special phosphoramidites, so the cost is lower. In addition, Glen-Pak cartridges allow purification of virtually the complete range of dyes and modifiers. The Glen-Pak DNA Cartridge 3g is a large cartridge capable of purifying 10–20 µmole oligonucleotide syntheses using the standard DMT-on procedure and Glen-Pak DNA 30mg 96-Well Plates are for parallel purification of up to 50 nmole scale syntheses. Scale suggestions for the Glen-Pak DNA product line are shown below:

<i>Glen-Pak DNA Product</i>	<i>Catalog Number</i>	<i>Synthesis Scale Compatibility</i>
Glen-Pak DNA 50mg Purification Cartridge	60-5000-96	10 nmole – 200 nmole
Glen-Pak DNA Purification Cartridge	60-5100-XX and 60-5200-XX	10 nmole – 1.0 µmole
Glen-Pak DNA Cartridge 3G	60-5300-01	5 µmole – 20 µmole
Glen-Pak DNA 30 mg 96-Well Plate	60-5400-01	10 nmole – 50 nmole

A User Guide to *Glen-Pak™ Purification* describes in detail the process and several applications for DNA and RNA purification. This booklet is available online at: http://www.glenresearch.com/Technical/GlenPak_UserGuide.pdf.



SEE ALSO

Poly-Pak Reagents

p139

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>DNA Purification Cartridges</i>			
Glen-Pak™ 50mg DNA Purification Cartridge (For use in vacuum manifolds and high-throughput devices)	60-5000-96	Pack of 96	415.00
Glen-Pak™ DNA Purification Cartridge (For use in vacuum manifolds and high-throughput devices)	60-5100-10	Pack of 10	80.00
	60-5100-30	Pack of 30	200.00
	60-5100-96	Pack of 96	475.00
Glen-Pak™ DNA Purification Cartridge (For use with disposable syringes)	60-5200-01	each	8.00
	60-5200-10	Pack of 10	80.00
Glen-Pak™ DNA Cartridge 3g	60-5300-01	Pack of 1	150.00
Glen-Pak™ DNA 30mg 96-Well Plate	60-5400-01	Pack of 1	475.00
<i>RNA Purification Cartridges</i>			
Glen-Pak™ RNA Purification Cartridge (For use in vacuum manifolds and high-throughput devices)	60-6100-10	Pack of 10	95.00
	60-6100-30	Pack of 30	225.00
	60-6100-96	Pack of 96	575.00
Glen-Pak™ RNA Purification Cartridge (For use with disposable syringes)	60-6200-01	each	9.50
	60-6200-10	Pack of 10	95.00
<i>Reagents</i>			
RNA Quenching Buffer	60-4120-82	250mL	80.00
	60-4120-80	1L	200.00
<i>Racks and Seals</i>			
Adapter Rack (For use with 96 well manifolds)	60-0010-01	each	10.00
Seal for Adapter Rack (For use on 96 well adapter rack)	60-0020-01	each	20.00

Poly-Pak™ and Glen-Pak™ are trademarks of Glen Research Corporation

POLY-PAK™ PURIFICATION

The use of Poly-Pak™ packings in cartridges or barrels overcomes several disadvantages usually associated with reverse phase (RP) cartridges. The packing is stable in the pH range 1-13, thus the ammonium hydroxide solution, diluted with water, is loaded directly onto the packing. Also, after elution of failure sequences, the trityl group is removed and washed from the support-bound oligonucleotide. The fully deprotected product can then be eluted and isolated by lyophilization. Poly-Pak™ Cartridges may also be used for desalting normal or labelled oligonucleotides. The original Poly-Pak cartridge and barrel are designed for 0.2 μ mole syntheses or less. Poly-Pak II cartridges and barrels are designed for use with 1 μ mole syntheses. A booklet, *User Guide To Poly-Pak™ Cartridge Purification*, describes in detail the process and several applications. This booklet is available online at: <http://www.glenresearch.com/Technical/PolyPakBooklet.pdf>.

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Packing, Cartridges and Barrels</i>			
Poly-Pak™ Packing	60-1000-05	5g	70.00
	60-1000-25	25g	350.00
Poly-Pak™ Cartridge	60-1100-01	each	8.00
	60-1100-10	Pack of 10	80.00
Poly-Pak™ II Cartridge	60-3100-01	each	12.00
	60-3100-10	Pack of 10	120.00
<i>Reagents</i>			
2.0M Triethylamine Acetate (TEAA) HPLC Grade	60-4110-52	200mL	60.00
	60-4110-57	450mL	120.00
	60-4110-60	960mL	200.00
	60-4110-62	2 L	
400.00 2% Aqueous Trifluoroacetic Acid	60-4040-57	450mL	36.00



Poly-Pak Cartridge Used Manually

Poly-Pak™ is a trademark of Glen Research Corporation

FLUORO-PAK™ PURIFICATION

Glen Research has discontinued the line of Fluoro-Pak™ products. These may be purchased directly from Berry Et Associates.

We recommend Glen-Pak purification as an alternative.


<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price (\$)</i>
<i>Phosphoramidites</i> (Discontinued)			
F-DMT-dA-CE Phosphoramidite	10-1400		
F-DMT-dC-CE Phosphoramidite	10-1410		
F-DMT-dG-CE Phosphoramidite	10-1420		
F-DMT-dT-CE Phosphoramidite	10-1430		
Fluorous Chemical Phosphorylation Reagent II (F-CPR-II)	10-1904		
<i>Columns</i> (Discontinued)			
Fluoro-Pak™ columns	61-2100		
Fluoro-Pak™ II columns	61-4100		
<i>Reagents</i> (Discontinued)			
Loading Buffer	61-4120		

SEE ALSO

Glen-Pak Purification p138

INTELLECTUAL PROPERTY

“Fluoro-Pak” is a trademark of Berry & Associates, Inc. Products for Fluorous Affinity Purification of Oligonucleotides: Patents applied for, Berry & Associates, Inc. Further, the use of these products is licensed under U.S. Patents 6,673,539, 6,156,896; 5,859,247; and 5,777,121 and one or more pending patents owned or controlled by Fluorous Technologies, Inc.



GLEN GEL-PAK™ DESALTING

The principle of the Glen Research gel filtration column, Glen Gel-Pak™, is based on size exclusion chromatography that separates molecules according to the hydrodynamic volume of the molecule in aqueous solutions. In gel filtration, the mobile phase for size exclusion is an aqueous solution and the stationary phase is a porous resin. The pores of the resin are sized such that they allow small molecules to enter the pores, yet exclude larger molecules from the pores. The small molecules, such as salts and hydrolyzed protecting groups, diffuse into the pores of the resin and move slowly through the column. The larger molecules, such as DNA or proteins, are excluded from the pores and move quickly through the column. The end result is that the larger molecules elute first in the column void volume while the small molecules are still flowing through the resin of the column. Glen Gel-Pak columns are ideal for desalting and reaction clean up. They can be used for removal of the ammonium hydroxide deprotection solution and hydrolyzed protecting groups after deprotection. The columns can also be used for the clean up of NHS-labelling reactions to separate the labelled oligo and unlabelled oligo from the unreacted NHS ester, the hydrolyzed label, and n-hydroxysuccinimide, thereby greatly simplifying the downstream purification steps.

There are many benefits to Glen Gel-Pak columns:

Versatility:

- Ability to directly desalt oligonucleotides deprotected in either 30% ammonium hydroxide OR 50:50 ammonium hydroxide/40% aqueous methylamine (AMA)
- Easily exchange buffers
- Simple clean-up of labelling reactions
- Mild method for purification from salts and solvents such as DMSO and DMF

Capacity:

- Multiple column sizes (0.2 mL, 1.0 mL and 2.5 mL) are available to match synthesis scale
- Ability to efficiently desalt short and long oligos at different scales using the same protocol
- Suitable for oligos >10mer in length



Glen Gel-Pak 0.2

Glen Gel-Pak 2.5

Glen Gel-Pak 1.0

<i>Item</i>	<i>Catalog No.</i>	<i>Pack</i>	<i>Price(\$)</i>
Glen Gel-Pak™ 0.2 Desalting Column (0.2 mL Capacity)	61-5002-05	Pack of 5	30.00
	61-5002-50	Pack of 50	300.00
Glen Gel-Pak™ 1.0 Desalting Column (1.0 mL Capacity)	61-5010-05	Pack of 5	35.00
	61-5010-50	Pack of 50	350.00
Glen Gel-Pak™ 2.5 Desalting Column (2.5 mL Capacity)	61-5025-05	Pack of 5	45.00
	61-5025-25	Pack of 25	225.00

Glen Gel-Pak™ is a trademark of Glen Research Corporation

OLIGO-AFFINITY SUPPORT

OTHER INSTRUMENT TYPES

All minor bases, RNA products and modifiers are packaged in septum-capped vials suitable for ABI and other instruments. If you would like another type of vial/column add the following to the end of the catalog number.

Monomers

For Instrument type Add

Expedite	E
MerMade	M

Columns

For Instrument type Add


Expedite	E
Applied Biosystems 3900	A
MerMade	M

(Please inquire for availability of vials and columns for other instrument types.)

Oligo-affinity supports (OAS) should ideally be compatible with automated synthesis, should be non-friable, should not shrink or swell, and should have low non-specific binding of the proteins or DNA. On the support shown below is an Adenosine residue attached through the exocyclic amino group. In this way, synthesis progresses regularly on removal of the 5'-DMT group. However, on treatment with ammonium hydroxide, the oligo is not cleaved from the support. This matrix can then be used as an affinity support for a complementary segment of DNA or RNA. Alternatively, the complementary strand can be annealed to the support and the double stranded DNA can be used as an affinity support for purifying DNA binding proteins.

We expect that OAS PS will be used for purification of components from biological fluids.

Item	Catalog No.	Pack	Price (\$)
Oligo-Affinity Support (PS) (OAS PS)	26-4001-01	0.1g	180.00
	26-4001-02	0.25g	425.00
	26-4001-10	1.0g	1590.00
Oligo-Affinity Support (PS) 1 µmole TWIST columns	26-4101-41	Pack of 4	300.00



The physical data table contains information which is unique to each monomer phosphoramidite. The molecular weight (MW) is the formula weight of the fully-protected monomer phosphoramidite. The MW is used to calculate the volume of solvent required to dilute 0.25g of the monomer to give a final 0.1M concentration. This figure is also shown in the table. The unit molecular weight (Unit FW) is the formula weight of each monomer once inserted into an oligonucleotide with all protecting groups removed. To obtain the molecular weight of a specific oligonucleotide, the following formula is used:

$$\text{Oligonucleotide MW} = \text{Sum of Unit FW} - 61.96$$

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution 10.1M</i>
10-0001	dA-5'-CE Phosphoramidite	857.95	313.21	0.25g/2.91mL
10-0101	dC-5'-CE Phosphoramidite	833.93	289.18	0.25g/3.00mL
10-0301	dT-5'-CE Phosphoramidite	744.83	304.2	0.25g/3.36mL
10-1000	dA-CE Phosphoramidite	857.95	313.21	0.25g/2.91mL
10-1001	7-Deaza-dA-CE Phosphoramidite	856.96	312.22	0.25g/2.92mL
10-1003	N6-Me-dA-CE Phosphoramidite	767.86	327.24	0.25g/3.26mL
10-1004	3'-dA-CE Phosphoramidite	857.95	313.21	0.25g/2.91mL
10-1006	Etheno-dA-CE Phosphoramidite	777.86	337.23	0.25g/3.21mL
10-1007	8-Br-dA-CE Phosphoramidite	887.81	392.11	0.25g/2.82mL
10-1008	8-oxo-dA-CE Phosphoramidite	873.95	329.21	0.25g/2.86mL
10-1010	dC-CE Phosphoramidite	833.93	289.18	0.25g/3.00mL
10-1014	pdC-CE Phosphoramidite	907.1	327.23	0.25g/2.76mL
10-1015	Ac-dC-CE Phosphoramidite	771.85	289.18	0.25g/3.24mL
10-1016	TMP-F-dU-CE Phosphoramidite	866.97	307.18	0.25g/2.88mL
10-1017	Pyrolo-dC-CE Phosphoramidite	767.85	327.23	0.25g/3.26mL
10-1018	5-Me-dC Brancher Phosphoramidite	942.1	402.36	0.25g/2.65mL
10-1019	Amino-Modifier C6 dC	1049.14	457.42	0.25g/2.38mL
10-1020	dG-CE Phosphoramidite	839.92	329.21	0.25g/2.98mL
10-1021	7-deaza-dG-CE Phosphoramidite	823.93	328.22	0.25g/3.03mL
10-1027	8-Br-dG-CE Phosphoramidite	903.9	408.1	0.25g/2.77mL
10-1028	8-oxo-dG-CE Phosphoramidite	855.93	345.21	0.25g/2.92mL
10-1029	dmf-dG-CE Phosphoramidite	824.92	329.21	0.25g/3.03mL
10-1030	dT-CE Phosphoramidite	744.83	304.2	0.25g/3.36mL
10-1031	5'-OMe-dT-CE Phosphoramidite	456.48	318.22	0.25g/5.48mL
10-1032	O4-Me-dT-CE Phosphoramidite	758.85	318.22	0.25g/3.29mL
10-1034	4-Thio-dT-CE Phosphoramidite	813.95	320.26	0.25g/3.07mL
10-1035	Carboxy-dT	814.88	360.22	0.25g/3.07mL
10-1036	2-Thio-dT-CE Phosphoramidite	879.02	320.26	0.25g/2.84mL
10-1037	Amino-Modifier C2 dT	938.94	402.3	0.25g/2.66mL
10-1038	Biotin-dT	1285.55	684.7	0.25g/1.94mL
10-1039	Amino-Modifier C6 dT	995.05	458.41	0.25g/2.51mL
10-1040	di-CE Phosphoramidite	754.79	314.19	0.25g/3.31mL
10-1041	2'-DeoxyNebularine-CE Phosphoramidite (Purine)	738.82	298.19	0.25g/3.38mL
10-1042	O6-Phenyl-di-CE Phosphoramidite	830.92	Varies	0.25g/3.01mL
10-1044	5-Nitroindole-CE Phosphoramidite	780.86	340.23	0.25g/3.20mL
10-1046	2-Aminopurine-CE Phosphoramidite	809.01	313.21	0.25g/3.09mL
10-1047	dP-CE Phosphoramidite	771.85	330.23	0.25g/3.24mL
10-1048	dK-CE Phosphoramidite	853.96	358.25	0.25g/2.93mL
10-1050	dU-CE Phosphoramidite	730.8	290.17	0.25g/3.42mL
10-1051	O4-Triazolyl-dU-CE Phosphoramidite	781.84	varies	0.25g/3.20mL
10-1052	4-Thio-dU-CE Phosphoramidite	799.93	306.23	0.25g/3.13mL
10-1053	5-OH-dU-CE Phosphoramidite	788.83	306.17	0.25g/3.17mL
10-1054	pdU-CE Phosphoramidite	768.85	328.22	0.25g/3.25mL

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
10-1055	2'-deoxypseudoU-CE Phosphoramidite	730.8	290.17	0.25g/3.42mL
10-1056	Fluorescein-dT Phosphoramidite	1425.57	815.71	0.25g/1.75mL
10-1057	TAMRA-dT	1311.48	870.85	0.25g/1.91mL
10-1058	Dabcyl-dT	1150.32	709.7	0.25g/2.17mL
10-1059	EDTA-C2-dT-CE Phosphoramidite	1201.32	676.53	0.25g/2.08mL
10-1060	5-Me-dC-CE Phosphoramidite	847.9	303.21	0.25g/2.95mL
10-1061	5-Me-2'-deoxyZebularine-CE Phosphoramidite	728.82	288.19	0.25g/3.43mL
10-1062	5-Hydroxymethyl-dC-CE Phosphoramidite	917	319.21	0.25g/2.73mL
10-1063	5-OH-dC-CE Phosphoramidite	954.03	305.18	0.25g/2.62mL
10-1064	3'-dC-CE Phosphoramidite	833.92	289.18	0.25g/3.00mL
10-1065	dmf-5-Me-isodC-CE Phosphoramidite	798.91	303.21	0.25g/3.13mL
10-1066	5-Carboxy-dC-CE Phosphoramidite	905.97	333.19	0.25g/2.76mL
10-1068	N4-Et-dC-CE Phosphoramidite	757.87	317.42	0.25g/3.30mL
10-1070	06-Me-dG-CE Phosphoramidite	853.97	343.24	0.25g/2.93mL
10-1072	6-thio-dG-CE Phosphoramidite	934.97	345.26	0.25g/2.67mL
10-1073	7-Deaza-8-aza-dG-CE Phosphoramidite (PPG)	824.91	329.2	0.25g/3.03mL
10-1074	3'-dG-CE Phosphoramidite	824.92	329.21	0.25g/3.03mL
10-1076	7-deaza-dX-CE Phosphoramidite	769.83	329.21	0.25g/3.25mL
10-1078	dmf-isodG-CE Phosphoramidite	1020.13	329.21	0.25g/2.45mL
10-1079	8-Amino-dG-CE Phosphoramidite	895.01	344.22	0.25g/2.79mL
10-1080	5-Br-dC-CE Phosphoramidite	912.82	368.08	0.25g/2.74mL
10-1081	5-I-dC-CE Phosphoramidite	959.83	415.08	0.25g/2.60mL
10-1082	2-F-dI-CE Phosphoramidite	921.96	varies, 2F=332.18	0.25g/2.71mL
10-1083	7-deaza-8-aza-dA-CE Phosphoramidite	808.91	313.2	0.25g/3.09mL
10-1084	3'-dT-CE Phosphoramidite	744.83	304.2	0.25g/3.36mL
10-1085	2-Amino-dA-CE Phosphoramidite	1047.33	328.22	0.25g/2.39mL
10-1086	8-Amino-dA-CE Phosphoramidite	879.01	328.22	0.25g/2.84mL
10-1088	3-deaza-dA-CE Phosphoramidite	856.95	312.22	0.25g/2.92mL
10-1089	Amino-Modifier C6 dA	1068.14	427.4	0.25g/2.34mL
10-1090	5-Br-dU-CE Phosphoramidite	809.69	369.07	0.25g/3.09mL
10-1091	5-I-dU-CE Phosphoramidite	856.69	416.07	0.25g/2.92mL
10-1092	5-F-dU-CE Phosphoramidite	748.79	308.16	0.25g/3.34mL
10-1093	5-Hydroxymethyl-dU-CE Phosphoramidite	802.86	320.19	0.25g/3.11mL
10-1096	Thymidine Glycol CE Phosphoramidite	1007.36	338.21	0.25g/2.48mL
10-1097	AP-dC-CE Phosphoramidite	974.97	438.33	0.25g/2.56mL
10-1098	8,5'-Cyclo-dA CE Phosphoramidite	855.92	311.19	0.25g/2.92mL
10-1100	dA-Me Phosphoramidite	802.91	311.24	0.25g/3.11mL
10-1115	Ac-dC-Me Phosphoramidite	716.81	287.21	0.25g/3.49mL
10-1120	dG-Me Phosphoramidite	784.89	327.24	0.25g/3.19mL
10-1130	dT-Me Phosphoramidite	689.79	302.23	0.25g/3.62mL
10-1140	dA-PACE Phosphoramidite	928.02	354.24	0.25g/2.69mL
10-1150	Ac-dC-PACE Phosphoramidite	841.93	330.21	0.25g/2.97mL
10-1160	dG-PACE Phosphoramidite	910.01	370.24	0.25g/2.75mL
10-1170	dT-PACE Phosphoramidite	814.9	345.22	0.25g/3.07mL
10-1200	dA-H-Phosponate, TEA Salt	822.9	313.21	0.25g/3.04mL
10-1210	dC-H-Phosponate, DBU Salt	849.35	289.18	0.25g/2.94mL
10-1220	dG-H-Phosponate, TEA Salt	804.88	329.21	0.25g/3.11mL
10-1230	dT-H-Phosponate, TEA Salt	709.78	304.2	0.25g/3.52mL
10-1301	Pac-dA-Me Phosphoramidite	848.93	327.23 (Methyl triester)	0.25g/2.94mL
10-1315	Ac-dC-Me Phosphoramidite	732.81	303.21 (Methyl triester)	0.25g/3.41mL
10-1321	iPr-Pac-dG-Me Phosphoramidite	907.01	343.23 (Methyl triester)	0.25g/2.76mL
10-1330	dT-Me Phosphoramidite	705.79	318.22 (Methyl triester)	0.25g/3.54mL

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
10-1400	F-DMT-dA-CE Phosphoramidite	1304.05	313.21	0.25g/1.92mL
10-1410	F-DMT-dC-CE Phosphoramidite	1280.02	289.18	0.25g/1.95mL
10-1420	F-DMT-dG-CE Phosphoramidite	1286.03	329.21	0.25g/1.94mL
10-1430	F-DMT-dT-CE Phosphoramidite	1190.93	304.2	0.25g/2.10mL
10-1440	CleanAmp™-Pac-dA-CE Phosphoramidite	1045.25	523.56 (triester)	0.25g/2.39mL
10-1450	CleanAmp™-Ac-dC-CE Phosphoramidite	929.13	499.54 (triester)	0.25g/2.69mL
10-1460	CleanAmp™-Pac-dG-CE Phosphoramidite	1061.25	539.56 (triester)	0.25g/2.36mL
10-1470	CleanAmp™-dT-CE Phosphoramidite	902.11	514.55 (triester)	0.25g/2.77mL
10-1501	1-Me-dA-CE Phosphoramidite	814.31	328.24	0.25g/3.07mL
10-1503	N6-Ac-N6-Me-dA-CE Phosphoramidite	809.89	327.23	0.25g/3.09mL
10-1510	5-Hydroxymethyl-dC II-CE Phosphoramidite	785.82	319.21	0.25g/3.18mL
10-1511	5-aza-5,6-dihydro-dC-CE Phosphoramidite	787.89	292.18	0.25g/3.17mL
10-1513	N4-Ac-N4-Et-dC-CE Phosphoramidite	799.89	317.24	0.25g/3.13mL
10-1514	5-Formyl-dC-CE Phosphoramidite	915.96	317.19 (formyl) 349.23 (diol)	0.25g/2.73mL
10-1516	tC-CE Phosphoramidite	835.95	395.33	0.25g/2.99mL
10-1517	tC ^o -CE Phosphoramidite	819.88	379.26	0.25g/3.05mL
10-1518	tCnitro-CE Phosphoramidite	880.94	440.32	0.25g/2.84mL
10-1520	8-D-dG-CE Phosphoramidite	825.91	330.21	0.25g/3.03mL
10-1521	dDs-CE Phosphoramidite	819.95	379.33	0.25g/3.05mL
10-1522	Pac-ds-CE Phosphoramidite	970.08	395.33	0.25g/2.58mL
10-1523	dPa-CE Phosphoramidite	713.8	273.18	0.25g/3.50mL
10-1524	dDss-CE Phosphoramidite	902.07	461.451	0.25g/2.77mL
10-1529	N2-Amino-Modifier C6 dG	965.01	428.38	0.25g/2.59mL
10-1530	5,6-Dihydro-dT-CE Phosphoramidite	746.84	306.21	0.25g/3.35mL
10-1531	N3-Cyanoethyl-dT	797.88	357.26	0.25g/3.13mL
10-1532	5'-Dabsyl-dT-CE Phosphoramidite	729.78	591.53	0.25g/3.43mL
10-1534	N-POM Caged-dT-CE Phosphoramidite	967.99	527.38 (N-POM-dT)	0.25g/2.58mL
10-1535	NHS-Carboxy-dT	897.91	varies, -CO2H=360.22	0.25g/2.78mL
10-1536	Fmoc Amino-Modifier C6 dT	1121.28	458.41(NH2)	0.25g/2.23mL
10-1537	dX-CE Phosphoramidite	1069.1	330.19	0.25g/2.34mL
10-1538	S-Bz-Thiol-Modifier C6-dT	1091.26	546.53	0.25g/2.29mL
10-1539	DBCO-dT-CE Phosphoramidite	1214.57	773.77	0.25g/2.06mL
10-1540	C8-Alkyne-dT-CE Phosphoramidite	834.94	394.32	0.25g/2.99mL
10-1541	C8-TIPS-Alkyne-dC-CE Phosphoramidite	1094.4	393.33	0.25g/2.28mL
10-1542	C8-TMS-Alkyne-dC-CE Phosphoramidite	1010.24	393.33	0.25g/2.47mL
10-1543	C8-Alkyne-dC-CE Phosphoramidite	938.06	393.33	0.25g/2.67mL
10-1544	C8-TIPS-Alkyne-dT-CE Phosphoramidite	991.28	394.32	0.25g/2.52mL
10-1545	C8-TMS-Alkyne-dT-CE Phosphoramidite	907.12	394.32	0.25g/2.76mL
10-1550	5,6-Dihydro-dU-CE Phosphoramidite	732.81	292.19	0.25g/3.41mL
10-1554	5-Ethynyl-dU-CE Phosphoramidite	754.81	314.19	0.25g/3.31mL
10-1560	Ac-5-Me-dC-CE Phosphoramidite	785.86	303.21	0.25g/3.18mL
10-1564	5-Formyl dC III CE Phosphoramidite	950.02	317.19 375.27 (acetal)	0.25g/2.63mL
10-1576	Ferrocene-dT-CE Phosphoramidite	1125.07	684.45	0.25g/2.22mL
10-1590	Pyrene-dU-CE Phosphoramidite	955.04	514.42	0.25g/2.62mL
10-1591	Perylene-dU-CE Phosphoramidite	1005.1	564.48	0.25g/2.49mL
10-1598	8,5'-Cyclo-dG-CE Phosphoramidite	619.65	327.19	0.25g/4.03mL
10-1601	Pac-dA-CE Phosphoramidite	887.97	313.21	0.25g/2.82mL
10-1621	iPr-Pac-dG-CE Phosphoramidite	946.05	329.21	0.25g/2.64mL
10-1700	dA-Thiophosphoramidite	955.09	345.34(dithioate)	0.25g/1.75mL
10-1710	dC-Thiophosphoramidite	931.07	321.31(dithioate)	0.25g/1.79mL

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
10-1720	dG-Thiophosphoramidite	937.07	361.34(dithioate)	0.25g/1.78mL
10-1730	dT-Thiophosphoramidite	841.97	336.32(dithioate)	0.25g/1.98mL
10-1900	Chemical Phosphorylation Reagent	656.77	79.98	0.25g/3.81mL
10-1901	Chemical Phosphorylation Reagent II	722.82	79.98	0.25g/3.46mL
10-1902	Solid Chemical Phosphorylation Reagent II	692.79	79.98	0.25g/3.61mL
10-1904	Fluorous Chemical Phosphorylation Reagent II	1168.92	79.98	0.25g/2.14mL
10-1905	5'-Amino-Modifier 5	577.71	167.1	0.25g/4.33mL
10-1906	5'-Amino-Modifier C6	589.76	179.16	0.25g/4.24mL
10-1907	5'-DMS(O)MT-Amino-Modifier C6	681.34	179.16	0.25g/3.67mL
10-1908	5'-Hexynyl Phosphoramidite	298.36	160.11	0.25g/8.38mL
10-1909	Spacer Phosphoramidite 9	652.77	212.14	0.25g/3.83mL
10-1912	5'-Amino-Modifier C12	673.92	263.32	0.25g/3.71mL
10-1913	Spacer Phosphoramidite C3	578.69	138.06	0.25g/4.32mL
10-1914	dSpacer CE Phosphoramidite	620.73	180.1	0.25g/4.03mL
10-1915	Pyrrolidine-CE Phosphoramidite	841.97	178.1	0.25g/2.97mL
10-1916	5'-Amino-Modifier C6-TFA	413.42	179.16	0.25g/6.05mL
10-1917	5'-Amino-Modifier TEG CE-Phosphoramidite	489.47	255.21	0.25g/5.11mL
10-1918	Spacer Phosphoramidite 18	784.93	344.3	0.25g/3.18mL
10-1919	5'-Aminoxy-Modifier-11-CE Phosphoramidite	711.82	271.21	0.25g/3.51mL
10-1920	Symmetric Doubler Phosphoramidite	1095.32	351.31	0.25g/2.28mL
10-1922	Trebler Phosphoramidite	1417.72	370.33	0.25g/1.76mL
10-1923	5'-Amino-Modifier C3-TFA	371.34	137.08	0.25g/6.73mL
10-1925	Long Trebler Phosphoramidite	1475.78	428.41	0.25g/1.69mL
10-1926	5'-Thiol-Modifier C6	576.78	196.2	0.25g/4.33mL
10-1927	Abasic II Phosphoramidite	750.98	196.1	0.25g/3.33mL
10-1928	Spacer C12 CE Phosphoramidite	704.93	264.3	0.25g/3.55mL
10-1931	5'-I-dT-CE Phosphoramidite	552.35	414.09	0.25g/4.53mL
10-1932	5'-Amino-dT-CE Phosphoramidite	713.81	303.21	0.25g/3.50mL
10-1933	5'-Aldehyde-Modifier C2 Phosphoramidite	480.58	228.14	0.25g/5.20mL
10-1934	5-Formylindole-CE Phosphoramidite	763.86	323.24	0.25g/3.27mL
10-1935	5'-Carboxy-Modifier C10	485.56	varies, -CO ₂ H = 250.23	0.25g/5.15mL
10-1936	Thiol-Modifier C6 S-S	769.05	328.4 (disulfide) 196.2 (thiol)	0.25g/3.25mL
10-1938	5'-Maleimide-Modifier Phosphoramidite	437.47	299.22 (pre-retro-DA) 203.09 (maleimide)	0.25g/5.71mL
10-1939	Spermine Phosphoramidite	1233.17	408.52	0.25g/2.03mL
10-1941	5'-DBCO-TEG Phosphoramidite	708.82	570.57	0.25g/3.53mL
10-1946	5'-Bromoethyl Phosphoramidite	381.29	243.04 (bromide) 205.15 (azide)	0.25g/6.56mL
10-1947	5'-Amino-Modifier C6-PDA	478.57	179.15	0.25g/5.22mL
10-1948	5'-Amino-Modifier C12-PDA	562.7	263.32	0.25g/4.44mL
10-1949	5'-Amino-Modifier TEG PDA	554.62	255.21	0.25g/4.51mL
10-1952	DesthiobiotinTEG Phosphoramidite	980.19	539.56	0.25g/2.55mL
10-1953	Biotin Phosphoramidite	876.1	435.48	0.25g/2.85mL
10-1955	BiotinTEG Phosphoramidite	1010.24	569.61	0.25g/2.47mL
10-1963	Fluorescein Phosphoramidite	1207.5	598.56	0.25g/2.07mL
10-1964	6-Fluorescein Phosphoramidite	1176.35	566.48	0.25g/2.13mL
10-1973	Acridine Phosphoramidite	891.53	450.86	0.25g/2.80mL
10-1975	Cholesteryl-TEG Phosphoramidite	1196.6	755.97	0.25g/2.09mL
10-1976	5'-Cholesteryl-TEG Phosphoramidite	820.13	682.89	0.25g/3.05mL
10-1977	a-Tocopherol-TEG Phosphoramidite	1139.56	698.91	0.25g/2.19mL
10-1979	Stearyl Phosphoramidite	470.71	332.46	0.25g/5.31mL

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution 10.1M</i>
10-1982	Psoralen C2 Phosphoramidite	502.55	364.29	0.25g/4.97mL
10-1983	Psoralen C6 Phosphoramidite	558.65	420.4	0.25g/4.48mL
10-1985	DNP-TEG Phosphoramidite	950	509.41	0.25g/2.63mL
10-1986	5'-Trimethoxystilbene Cap Phosphoramidite	571.65	433.39	0.25g/4.37mL
10-1987	5'-Pyrene Cap Phosphoramidite	501.6	363.35	0.25g/4.98mL
10-1992	Alkyne-Modifier Serinol Phosphoramidite	758.88	318.26	0.25g/3.29mL
10-1993	Protected Biotin Serinol Phosphoramidite	1051.28	450.45	0.25g/2.38mL
10-1994	6-Fluorescein Serinol Phosphoramidite	1191.3	582.45	0.25g/2.10mL
10-1995	Protected BiotinLC Serinol Phosphoramidite	1298.57	697.74	0.25g/1.93mL
10-1997	Amino-Modifier Serinol Phosphoramidite	887.01	224.15	0.25g/2.82mL
10-3000	Pac-A-CE Phosphoramidite	1018.23	329.21	0.25g/2.46mL
10-3001	7-deaza-A-CE Phosphoramidite	987.2	328.22	0.25g/2.53mL
10-3003	Bz-A-CE Phosphoramidite	988.21	329.21	0.25g/2.53mL
10-3004	A-TOM-CE Phosphoramidite	998.24	329.21	0.25g/2.50mL
10-3011	Zebularine-CE Phosphoramidite	845.05	290.17	0.25g/2.96mL
10-3012	Pyridin-2-one-CE Phosphoramidite	844.06	289.18	0.25g/2.96mL
10-3014	C-TOM-CE Phosphoramidite	974.22	305.18	0.25g/2.57mL
10-3015	Ac-C-CE Phosphoramidite	902.11	305.18	0.25g/2.77mL
10-3017	Pyrrolo-C-TOM-CE Phosphoramidite	970.23	343.27	0.25g/2.58mL
10-3021	iPr-Pac-G-CE Phosphoramidite	1076.31	345.21	0.25g/2.32mL
10-3024	G-TOM-CE Phosphoramidite	1014.24	345.21	0.25g/2.46mL
10-3025	Ac-G-CE Phosphoramidite	941.43	345.21	0.25g/2.66mL
10-3030	U-CE Phosphoramidite	861.06	306.17	0.25g/2.90mL
10-3034	U-TOM-CE Phosphoramidite	933.17	306.17	0.25g/2.68mL
10-3039	Amino-Modifier C6-U Phosphoramidite	1197.41	474.4	0.25g/2.09mL
10-3040	I-CE Phosphoramidite	885.08	330.19	0.25g/2.82mL
10-3041	Nebularine-CE Phosphoramidite	869.07	314.19	0.25g/2.88mL
10-3050	5-Me-U-CE Phosphoramidite	875.08	320.19	0.25g/2.86mL
10-3052	4-Thio-U-TOM-CE Phosphoramidite	1002.29	322.22	0.25g/2.49mL
10-3055	PseudoUridine-CE Phosphoramidite	861.05	306.17	0.25g/2.90mL
10-3064	5-Me-C-TOM-CE Phosphoramidite	988.25	319.21	0.25g/2.53mL
10-3070	2-Aminopurine-TBDMS-CE Phosphoramidite	954.19	329.21	0.25g/2.62mL
10-3072	6-Thio-G-CE Phosphoramidite	1039.31	361.26	0.25g/2.41mL
10-3083	8-Aza-7-deaza-A-CE Phosphoramidite	939.16	329.21	0.25g/2.66mL
10-3085	2,6-Diaminopurine-TOM-CE Phosphoramidite	1113.36	344.22	0.25g/2.25mL
10-3090	Br-U-CE Phosphoramidite	939.96	385.06	0.25g/2.66mL
10-3091	5-I-U-CE Phosphoramidite	986.96	432.07	0.25g/2.53mL
10-3100	2'-OMe-A-CE Phosphoramidite	887.97	343.24	0.25g/2.82mL
10-3110	2'-OMe-C-CE Phosphoramidite	863.95	319.21	0.25g/2.89mL
10-3111	2'-OMe-TMP-5-F-U-CE Phosphoramidite	897.08	337.2	0.25g/2.79mL
10-3115	2'-OMe-Ac-C-CE Phosphoramidite	801.88	319.21	0.25g/3.12mL
10-3116	2'-OMe-3-deaza-5-aza-C-CE Phosphoramidite	816.91	319.21	0.25g/3.06mL
10-3120	2'-OMe-ibu-G-CE Phosphoramidite	869.97	359.24	0.25g/2.87mL
10-3121	2'-OMe-G-CE Phosphoramidite	854.93	359.24	0.25g/2.92mL
10-3123	2'-OMe-2-Aminopurine-CE Phosphoramidite	839.04	343.24	0.25g/2.98mL
10-3124	2'-OMe-2,6-Diaminopurine-CE Phosphoramidite	924.05	358.25	0.25g/2.71mL
10-3130	2'-OMe-U-CE Phosphoramidite	760.82	320.2	0.25g/3.29mL
10-3131	2'-OMe-5-Me-U-CE Phosphoramidite	774.84	334.22	0.25g/3.23mL
10-3132	2'-OMe-5-F-U-CE Phosphoramidite	778.78	338.19	0.25g/3.21mL
10-3140	2'-OMe-I-CE Phosphoramidite	784.85	344.22	0.25g/3.19mL
10-3160	2'-OMe-5-Me-C-CE Phosphoramidite	815.9	333.24	0.25g/3.06mL
10-3190	2'-OMe-5-Br-U-CE Phosphoramidite	839.72	399.09	0.25g/2.98mL

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
10-3400	2'-F-A-CE Phosphoramidite	875.93	331.2	0.25g/2.85mL
10-3415	2'-F-Ac-C-CE Phosphoramidite	789.84	307.18	0.25g/3.17mL
10-3420	2'-F-G-CE Phosphoramidite	857.91	347.19	0.25g/2.91mL
10-3430	2'-F-U-CE Phosphoramidite	748.79	308.16	0.25g/3.34mL
10-3501	1-Me-A-CE Phosphoramidite	944.57	344.24	0.25g/2.65mL
10-3601	2'-OMe-Pac-A-CE Phosphoramidite	917.99	343.24	0.25g/2.72mL
10-3621	2'-OMe-iPr-Pac-G-CE Phosphoramidite	976.07	359.24	0.25g/2.56mL
10-3800	2'-FANA-A-CE Phosphoramidite	875.93	331.2	0.25g/2.85mL
10-3810	2'-FANA-C-CE Phosphoramidite	851.91	307.17	0.25g/2.93mL
10-3820	2'-FANA-G-CE Phosphoramidite	857.91	347.19	0.25g/2.91mL
10-3830	2'-FANA-U-CE Phosphoramidite	748.79	308.16	0.25g/3.34mL
10-3914	rSpacer CE Phosphoramidite	823.09	196.09	0.25g/3.04mL
10-4410	UniCap Phosphoramidite	334.39		0.25g/7.48mL
10-4906	PC Amino-Modifier Phosphoramidite	605.59	371.32	0.25g/4.13mL
10-4913	PC Spacer Phosphoramidite	784.88	344.26	0.25g/3.19mL
10-4920	PC Linker Phosphoramidite	699.78	259.15	0.25g/3.57mL
10-4950	PC Biotin Phosphoramidite	1038.25	597.62	0.25g/2.41mL
10-5800	Azobenzene Phosphoramidite	815.94	375.32	0.25g/3.06mL
10-5801	2,2'-Dipicolylamine Phosphoramidite	775.91	335.29	0.25g/3.22mL
10-5901	5'-Fluorescein Phosphoramidite	843.95	537.46	0.25g/2.96mL
10-5902	5'-Hexachloro-Fluorescein Phosphoramidite	1050.62	744.13	0.25g/2.38mL
10-5903	5'-Tetrachloro-Fluorescein Phosphoramidite	981.73	675.24	0.25g/2.55mL
10-5905	SIMA (HEX) Phosphoramidite	1065.02	759.54	0.25g/2.35mL
10-5906	5'-Dichloro-dimethoxy-Fluorescein Phosphoramidite	11972.88	666.4	0.25g/2.57mL
10-5912	5'-DabcyI Phosphoramidite	568.69	430.18	0.25g/4.40mL
10-5913	Cyanine 3 Phosphoramidite	953.64	507.59	0.25g/2.62mL
10-5914	Cyanine 3.5 Phosphoramidite	1053.76	607.7	0.25g/2.37mL
10-5915	Cyanine 5 Phosphoramidite	979.68	533.63	0.25g/2.55mL
10-5916	Cyanine 5.5 Phosphoramidite	1171.25	633.74	0.25g/2.13mL
10-5917	DyLight DY547 Phosphoramidite	756.74	490.57	0.25g/3.30mL
10-5918	DyLight DY647 Phosphoramidite	827.07	516.61	0.25g/3.02mL
10-5920	Epoch Redmond Red™ Phosphoramidite	971.09	445.34	0.25g/2.57mL
10-5921	Epoch Yakima Yellow™ Phosphoramidite	1023.81	718.33	0.25g/2.44mL
10-5922	Epoch Gig Harbor Green™ Phosphoramidite	900.06	594.57	0.25g/2.78mL
10-5925	Epoch Eclipse™ Quencher Phosphoramidite	978.5	537.89	0.25g/2.55mL
10-5931	5'-BHQ-1 Phosphoramidite	676.75	538.49	0.25g/3.69mL
10-5932	5'-BHQ-2 Phosphoramidite	678.72	540.47	0.25g/3.68mL
10-5934	5'-BBQ-650®-CE Phosphoramidite	802.9	665.65	0.25g/3.11mL
10-5941	BHQ-1-dT	1401.56	960.93	0.25g/1.78mL
10-5942	BHQ-2-dT	1403.53	962.91	0.25g/1.78mL
10-5944	BBQ-650®-dT-CE Phosphoramidite	1441.57	1000.95	0.25g/1.73mL
10-5945	SIMA (HEX)-dT Phosphoramidite	1646.64	1037.79	0.25g/1.52mL
10-5950	5'-Biotin Phosphoramidite	846.08	405.45	0.25g/2.95mL
10-5960	Methylene Blue C3 Phosphoramidite	994.7	517.62	0.25g/2.51mL
10-7001	2',3'-ddA-CE Phosphoramidite	574.7	297.21	0.25g/4.35mL
10-7101	2',3'-ddC-CE Phosphoramidite	550.68	273.18	0.25g/4.54mL
10-7201	2',3'-ddG-CE Phosphoramidite	506.54	313.2	0.25g/4.94mL
10-7301	2',3'-ddT-CE Phosphoramidite	426.45	288.19	0.25g/5.86mL
10-9201	dmf-dG-5'-CE Phosphoramidite	824.92	329.21	0.25g/3.03mL
11-1330	Cis-syn Thymine Dimer Phosphoramidite	1024.01	608.39	0.25g/2.44mL

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
13-1000	AAA Trimer Phosphoramidite	1911.5		0.25g/1.31mL
13-1001	AAC Trimer Phosphoramidite	1887.5		0.25g/1.32mL
13-1011	ACC Trimer Phosphoramidite	1863.5		0.25g/1.34mL
13-1013	ACT Trimer Phosphoramidite	1774.5		0.25g/1.41mL
13-1020	AGA Trimer Phosphoramidite	1893.5		0.25g/1.32mL
13-1031	ATC Trimer Phosphoramidite	1774.5		0.25g/1.41mL
13-1032	ATG Trimer Phosphoramidite	1780.5		0.25g/1.40mL
13-1102	CAG Trimer Phosphoramidite	1869.5		0.25g/1.34mL
13-1103	CAT Trimer Phosphoramidite	1774.5		0.25g/1.41mL
13-1110	CCA Trimer Phosphoramidite	1863.5		0.25g/1.34mL
13-1112	CCG Trimer Phosphoramidite	1845.5		0.25g/1.35mL
13-1122	CGG Trimer Phosphoramidite	1851.5		0.25g/1.35mL
13-1123	CGT Trimer Phosphoramidite	1756.5		0.25g/1.42mL
13-1132	CTG Trimer Phosphoramidite	1756.5		0.25g/1.42mL
13-1200	GAA Trimer Phosphoramidite	1893.5		0.25g/1.32mL
13-1201	GAC Trimer Phosphoramidite	1869.5		0.25g/1.34mL
13-1203	GAT Trimer Phosphoramidite	1780.5		0.25g/1.40mL
13-1210	GCA Trimer Phosphoramidite	1869.5		0.25g/1.34mL
13-1212	GCG Trimer Phosphoramidite	1851.5		0.25g/1.35mL
13-1213	GCT Trimer Phosphoramidite	1756.5		0.25g/1.42mL
13-1223	GGT Trimer Phosphoramidite	1762.5		0.25g/1.42mL
13-1230	GTA Trimer Phosphoramidite	1780.5		0.25g/1.40mL
13-1233	GTT Trimer Phosphoramidite	1667.5		0.25g/1.50mL
13-1301	TAC Trimer Phosphoramidite	1774.5		0.25g/1.41mL
13-1313	TCT Trimer Phosphoramidite	1661.4		0.25g/1.50mL
13-1321	TGC Trimer Phosphoramidite	1756.5		0.25g/1.42mL
13-1322	TGG Trimer Phosphoramidite	1762.5		0.25g/1.42mL
13-1331	TTC Trimer Phosphoramidite	1661.4		0.25g/1.50mL
13-1333	TTT Trimer Phosphoramidite	1572.4		0.25g/1.59mL
20-0002	dA-5'-CPG		313.21	
20-0102	dC-5'-CPG		289.18	
20-0202	dG-5'-CPG		329.21	
20-0302	dT-5'-CPG		304.2	
20-2000	dA-CPG 500		313.21	
20-2001	dA-CPG 1000		313.21	
20-2002	dA-CPG 2000		313.21	
20-2004	3'-dA-CPG		313.21	
20-2010	dC-CPG 500		289.18	
20-2011	dC-CPG 1000		289.18	
20-2012	dC-CPG 2000		289.18	
20-2013	Ac-dC-CPG 500		289.18	
20-2015	Ac-dC-CPG 1000		289.18	
20-2017	2',3'-ddC-CPG		273.19	
20-2019	3'-Amino-Modifier C6 dC CPG		457.42	
20-2020	dG-CPG 500		329.21	
20-2021	dG-CPG 1000		329.21	
20-2022	dG-CPG 2000		329.21	
20-2029	dmf-dG-CPG		329.21	
20-2030	dT-CPG 500		304.2	
20-2031	dT-CPG 1000		304.2	
20-2032	dT-CPG 2000		304.2	

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
20-2040	dI-CPG 500		314.19	
20-2041	dI-CPG 1000		314.19	
20-2050	dU-CPG 500		290.17	
20-2051	dU-CPG 1000		290.17	
20-2056	3'-Fluorescein-dT CPG		815.71	
20-2060	5-Me-dC-CPG		303.21	
20-2064	3'-dC-CPG		289.18	
20-2074	3'-dG-CPG		329.21	
20-2084	3'-dT-CPG		304.2	
20-2090	5-Br-dU-CPG		369.07	
20-2101-61	dA-CPG 1000		313.21	
20-2101-62	dA-CPG 1000		313.21	
20-2101-65	dA-CPG 1000		313.21	
20-2115-61	Ac-dC-CPG 1000		289.18	
20-2115-62	Ac-dC-CPG 1000		289.18	
20-2115-65	Ac-dC-CPG 1000		289.18	
20-2129-61	dmf-dG-CPG		329.21	
20-2129-62	dmf-dG-CPG		329.21	
20-2129-65	dmf-dG-CPG		329.21	
20-2131-61	dT-CPG 1000		304.2	
20-2131-62	dT-CPG 1000		304.2	
20-2131-65	dT-CPG 1000		304.2	
20-2601	Pac-dA-CPG		313.21	
20-2621	iPr-Pac-dG-CPG		329.21	
20-2900	3'-Phosphate CPG		79.98	
20-2902	3'-Glyceryl CPG		154.06	
20-2903	3'-CPR II CPG		79.98	
20-2913	3'-Spacer C3 CPG		138.06	
20-2933	3'-Thiol-Modifier C3 S-S CPG	154.12 (thiol), 244.27 (disulfide)		
20-2938	3'-Thiol-Modifier 6 S-S CPG	198.18 (thiol), 332.37 (disulfide)		
20-2952	DesthiobiotinTEG-CPG		539.56	
20-2954	3'-PT-Amino-Modifier C3 CPG		137.07	
20-2955	3'-BiotinTEG CPG		569.61	
20-2956	3'-PT-Amino-Modifier C6 CPG		179.15	
20-2958	3'-Amino-Modifier C7 CPG 1000		209.18	
20-2961	3'-(6-FAM) CPG		569.46	
20-2963	3'-Fluorescein CPG		598.56	
20-2964	3'-(6-Fluorescein) CPG		566.48	
20-2973	3'-Acridine CPG		450.86	
20-2975	3'-Cholesteryl-TEG CPG		755.97	
20-2980	3'-Uaq Cap CPG		539.39	
20-2981	3'-Amino-dT CPG		303.21	
20-2982	3'-Propargyl-5-Me-dC CPG		341.26	
20-2992	3'-Alkyne-Modifier Serinol CPG		334.26	
20-2993	3'-Protected Biotin Serinol CPG		450.45	
20-2994	3'-6-Fluorescein Serinol CPG		584.47	
20-2995	3'-Protected BiotinLC Serinol CPG		697.74	
20-2997	3'-Amino-Modifier Serinol CPG		224.15	
20-3300	Pac-A-RNA-CPG		329.21	
20-3303	Bz-A-RNA-CPG		329.21	
20-3304	Ac-A-RNA-CPG		329.21	
20-3315	Ac-C-RNA-CPG		305.18	

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>Phosphoramidite MW</i>	<i>Unit FW</i>	<i>Dilution (0.1M)</i>
20-3321	iPr-Pac-G-RNA-CPG		345.21	
20-3324	Ac-G-RNA-CPG		345.21	
20-3330	U-RNA-CPG		306.17	
20-3600	2'-OMe-A-RNA-CPG		343.24	
20-3610	2'-OMe-C-RNA-CPG		319.21	
20-3615	2'-OMe-Ac-C-RNA-CPG		319.21	
20-3621	2'-OMe-G-RNA-CPG		359.24	
20-3630	2'-OMe-U-RNA-CPG		320.2	
20-4040	Puromycin-CPG		533.48	
20-5910	3'-TAMRA CPG		623.6	
20-5911	3'-Dabsyl CPG		498.49	
20-5912	3'-Dabeyl CPG		462.44	
20-5913	Cyanine 3 CPG		507.59	
20-5915	Cyanine 5 CPG		533.63	
20-5920	Epoch Redmond Red™ CPG		445.34	
20-5921	Epoch Yakima Yellow™ CPG		718.33	
20-5925	Epoch Eclipse™ Quencher CPG		537.89	
20-5931	3'-BHQ-1 CPG		554.49	
20-5932	3'-BHQ-2 CPG		556.47	
20-5933	3'-BHQ-3 CPG		597.63	
20-5934	BBQ-650® CPG		667.63	
21-2000	dA-Q-CPG 500		313.21	
21-2010	dC-Q-CPG 500		289.18	
21-2013	Ac-dC-Q-CPG 500		305.18	
21-2029	dmf-dG-Q-CPG 500		329.21	
21-2030	dT-Q-CPG 500		304.2	
25-2000	dA-High Load-CPG		313.21	
25-2010	dC-High Load-CPG		289.18	
25-2020	dG-High Load CPG		329.21	
25-2030	dT-High Load-CPG		304.2	
25-2900	3'-Phosphate CPG (High Load)		79.98	
26-2600	dA PS		313.21	
26-2610	dC PS		289.18	
26-2629	dmf-dG PS		329.21	
26-2630	dT-PS		304.2	
26-2900	3'-Phosphate PS		79.98	
26-2955	3'-BiotinTEG PS		569.61	
26-2956	3'-PT-Amino-Modifier C6 PS		179.15	
26-2961	3'-(6-FAM) PS		569.46	
26-5910	3'-TAMRA PS		623.6	
26-5912	3'-Dabeyl PS		462.44	

PHYSICAL DATA

<i>Cat. No.</i>	<i>Item</i>	<i>MW</i>	<i>Unit FW</i>
50-1904	Azidobutyrate NHS Ester	226.19	113.12
50-1905	Alkyne-NHS Ester	225.2	110.11
50-1941	DBCO-sulfo-NHS Ester	532.5	316.37
50-2000	BiotinTEG Azide	444.55	
50-2001	DesthiobiotinTEG Azide	414.5	
50-2002	Dipivaloyl 6-FAM-TEG Azide	744.79	
50-2003	6-FAM-TEG Azide	576.55	
50-2004	Coumarin Azide	203.15	
50-2005	6-HEX Azide	665.09	
50-2006	6-TET Azide	596.2	
50-2007	TEMPO Azide	197.26	
50-2008	TEMPO-TEG Azide	373.47	
50-2009	Psoralen Azide	283.28	
50-2010	Disulfo-Cyanine 7 Azide	829.08	
50-5910	TAMRA NHS Ester	527.53	413.45

Symbols**2'-5' Linkages**

- 3'-dA-CE Phosphoramidite 61
- 3'-dA-CPG 61
- 3'-dC-CE Phosphoramidite 61
- 3'-dC-CPG 61
- 3'-dG-CE Phosphoramidite 61
- 3'-dG-CPG 62
- 3'-dT-CE Phosphoramidite 61
- 3'-dT-CPG 62

2'-5' Linked Oligonucleotides 60,62**5' -> 3' SYNTHESIS 30,31**

- 5'-CE Phosphoramidites 30
- 5'-Supports 30

A**A**

- 1-Me-A-CE Phosphoramidite 124
- 1-Me-dA-CE Phosphoramidite 62
- 2'-F-A-ANA CE Phosphoramidite 131
- 2'-F-A-CE Phosphoramidite 130
- 2'-OMe-A-CE Phosphoramidite 125
- 2'-OMe-A-RNA 127
- 2'-OMe-Pac-A-CE Phosphoramidite 126
- 2',3'-ddA 51
- 3'-dA-CE Phosphoramidite 61
- 3'-dA-CPG 50,62
- 7-Deaza-dA-CE Phosphoramidite 52
- 8-Amino-dA-CE Phosphoramidite 54
- 8-Br-dA-CE Phosphoramidite 55
- 8-Oxo-dA-CE Phosphoramidite 57
- Ac-A-RNA-CPG 115
- A-TOM-CE Phosphoramidite 115,117
- Amino-Modifier C6 dA 73
- Bz-A-CE Phosphoramidite 117
- Bz-A-RNA-CPG 114,118
- CleanAmp™-Pac-dA-CE Phosphoramidite 49
- dA-CE Phosphoramidite 6,10,13,14,16,18
- dA-H-Phosphonate 35
- dA-Me Phosphoramidite 32
- dA-Me Phosphoramidite 34
- dA-PACE Phosphoramidite 33
- dA-Thiophosphoramidite 36
- dbf-dA-CE Phosphoramidite 58
- N6-Ac-N6-Me-dA-CE Phosphoramidite 43,62
- N6-Me-A-CE Phosphoramidite 124
- N6-Me-dA-CE Phosphoramidite 43,62
- Pac-A-CE Phosphoramidite 118
- Pac-A-RNA-CPG 118
- Pac-dA-CE Phosphoramidite 20,68

A-2-Amino

- 2'-OMe-2-Amino-A-CE Phosphoramidite 128
- 2-Amino-A-TOM-CE Phosphoramidite 120
- 2-Amino-dA-CE Phosphoramidite 42

Abasic Site 58,80

- Abasic II Phosphoramidite 58
- dSpacer Phosphoramidite 58
- Pyrrolidine-CE Phosphoramidite 59

Acridine Labelling

- 3'-Acridine CPG 107
- Acridine Phosphoramidite 107

Activator (Powder)

- 4,5-Dicyanoimidazole 28
- 5-Benzylthio-1H-tetrazole 28
- 5-Ethylthio-1H-tetrazole 24,28,66,104,106
- Saccharin 1-Methylimidazole 28

Adamantane Carbonyl Chloride 35**Affinity Chromatography 142****Åkta oligopilot 16****Aldehyde Modifier**

- 5'-Aldehyde-Modifier C2 Phosphoramidite 79
- 5-Formyl-dC-CE Phosphoramidite 45
- Formylindole CE Phosphoramidite 79

Alternative Solvents and Reagents 28**Amino-dA**

- 8-Amino-dA-CE Phosphoramidite 54

Amino-dG

- 8-Amino-dG-CE Phosphoramidite 58

Amino-Modifiers

- 3'-Amino-Modifier C6 dC CPG 77
- 3'-Amino-Modifier C6 dT CPG 77
- 3'-Amino-Modifier C7 CPG 75
- 3'-Amino-Modifier Serinol CPG 75
- 3'-PT-Amino-Modifier C3 CPG 75
- 3'-PT-Amino-Modifier C6 CPG 75
- 3'-PT-Amino-Modifier C6 PS 75
- 3'-PT-Amino-Modifier C6 PS 9
- 5'-Amino-dT-CE Phosphoramidite 50
- 5'-Amino-Modifier 5 70
- 5'-Amino-Modifier C3-TFA 71
- 5'-Amino-Modifier C6 71
- 5'-Amino-Modifier C6-PDA 71
- 5'-Amino-Modifier C6-TFA 71
- 5'-Amino-Modifier C12 70
- 5'-Amino-Modifier C3-TFA 70
- 5'-Amino-Modifier C6 70
- 5'-Amino-Modifier C6-TFA 70
- 5'-Amino-Modifier C12-PDA 71
- 5'-Amino-Modifier TEG 70
- 5'-Amino-Modifier-TEG-PDA 71
- 5'-DMS(O)MT-Amino-Modifier C6 70
- Amino-Modifier C2 dT 73
- Amino-Modifier C6-U Phosphoramidite 121
- Amino-Modifier C6 dA 73
- Amino-Modifier C6 dC 73
- Amino-Modifier C6 dT 73
- Amino-Modifier Serinol Phosphoramidite 74
- Fmoc-Amino-Modifier C6 dT 74
- N2-Amino-Modifier C6 dG 73
- PC Amino-Modifier Phosphoramidite 72,82

AminoOxy-Modifier

- 5'-AminoOxy-Modifier 11 72

Aminopurine

- 2'-OMe-2-Aminopurine-CE Phosphoramidite 128
- 2-Aminopurine-CE Phosphoramidite 53,122

Antraquinone

3'-Uaq Cap CPG 44

Applied Biosystems Instruments

AB 3900 1000Å CPG Columns 8,19

AB 3900 Polystyrene Columns 8,19

AB 3900 Polystyrene Modifier Columns 8

CE Phosphoramidites 6

Solvents and Reagents 6,7,11

Supports and Columns 7,8,12

Ara-C

Ara-C-CE Phosphoramidite 67

Aza-dC

5-Aza-5,6-dihydro-dC-CE Phosphoramidite 67

A a-Thiotriphosphate Analogs

2'-deoxyAdenosine 134

2'-Fluoro-Adenosine 134

2'-OMe-Adenosine 134

2,6-Diaminopurine riboside 134

2-Aminopurine riboside 134

7-Deaza-Adenosine 133

Adenosine 133

N6-Me-Adenosine 133

Purine riboside 134

Azidobutyrate NHS Ester 85**Azobenzene**

Azobenzene Phosphoramidite 113

B**Backbone Modification 32****Beaucage Reagent, See Sulfurizing Reagent****Benzylthio-1H-tetrazole 28****Biocompatible Chemical Ligation 60****Biological Mimics 58****Biosearch Cyclone**

Supports and Columns 11

Biotin Labelling

3'-BiotinTEG CPG 94

3'-BiotinTEG PS 9,94

3'-Protected BiotinLC Serinol CPG 90,94

3'-Protected Biotin Serinol CPG 90,94

5'-Biotin Phosphoramidite 93

Biotin-dT 93

Biotin PaTP 48

BiotinTEG Azide 87

BiotinTEG Phosphoramidite 92

Biotin Phosphoramidite 92

DesthiobiotinTEG Azide 87

DesthiobiotinTEG-CPG 94

DesthiobiotinTEG Phosphoramidite 93

PC Biotin Phosphoramidite 82,93

Protected BiotinLC Serinol Phosphoramidite 89,92

Protected Biotin Serinol Phosphoramidite 89,92

BlackBerry® Quencher

3'-BBQ-650® CPG 106

5'-BBQ-650® Phosphoramidite 106

BBQ-650®-dT 106

Black Hole Quencher™ Dyes

3'-BHQ-1 CPG 105

3'-BHQ-2 CPG 105

3'-BHQ-3 CPG 105,106

5'-BHQ-1 Phosphoramidite 104

5'-BHQ-2 Phosphoramidite 66,104,106

BHQ-1-dT 104,106

BHQ-2-dT 104

Br-dA

8-Br-dA-CE Phosphoramidite 55

Br-dC

5-Br-dC-CE Phosphoramidite 55

Br-dG

8-Br-dG-CE Phosphoramidite 55

Br-dU

5-Br-dU-CE Phosphoramidite 55

5-Br-dU-CPG 55

Bromohexyl Phosphoramidite 85**Br-U**

2'-OMe-5-Br-U-CE Phosphoramidite 129

5-Br-U-CE Phosphoramidite 122

Brancher Phosphoramidite

dC Brancher Phosphoramidite 81

C**C**

2'-F-Bz-C-ANA CE Phosphoramidite 131

2'-OMe-5-Me-C-CE Phosphoramidite 129

2'-OMe-Ac-C-CE Phosphoramidite 125,126

2'-OMe-Ac-C-RNA 127

2'-OMe-C-CE Phosphoramidite 125

2'-OMe-C-RNA 127

2',3'-ddC-CPG 51

2',3'-ddC 51

3'-Amino-Modifier C6 dC CPG 77

3'-dC-CE Phosphoramidite 61

3'-dC-CPG 50,61

5-Br-dC-CE Phosphoramidite 55

5-Carboxy-dC-CE Phosphoramidite 45

5-Formyl-dC-CE Phosphoramidite 45

5-Hydroxymethyl-dC-CE Phosphoramidite 45

5-Hydroxymethyl-dC II-CE Phosphoramidite 45

5-I-dC-CE Phosphoramidite 55

5-Me-dC-CE Phosphoramidite 42

5-Me-dC-CPG 42

5-OH-dC-CE Phosphoramidite 57

8-Oxo-G Clamp-CE Phosphoramidite 65

Ac-C-CE Phosphoramidite 117

Ac-C-RNA-CPG 114,116,119

Ac-dC-CE Phosphoramidite 6,10,13,14,16,18,20,68

Ac-dC-Me Phosphoramidite 32

Ac-dC-PACE Phosphoramidite 33

Amino-Modifier C6 dC 73

AP-dC 64

AP-dC-CE Phosphoramidite 42

C8-Alkyne-dC-CE Phosphoramidite 83,84

C8-TIPS-Alkyne-dC-CE Phosphoramidite 83

C8-TMS-Alkyne-dC-CE Phosphoramidite 83

CleanAmp™-Ac-dC-CE Phosphoramidite 49

- C-TOM-CE Phosphoramidite 115,117
dC-CE Phosphoramidite 6,10,13,14,16,18
dC-H-Phosphonate 35
dC-Me Phosphoramidite 34
dC Brancher Phosphoramidite 81
dC-Thiophosphoramidite 36
N4-Et-dC-CE Phosphoramidite 43
pdC-CE Phosphoramidite 42
Pyrrolo-dCTP 64
tC-CE Phosphoramidite 66
tC^o-CE Phosphoramidite 66
- Caged Nucleosides**
NPOM-Caged-dT 66
- Camphorsulfonyloxaziridine (CSO) 29,33**
- Cap CPG**
3'-Uaq Cap CPG 44,60
- Capping Reagent**
UniCap Phosphoramidite 29
- Cap Phosphoramidite**
5'-Pyrene Cap Phosphoramidite 44
5'-Trimethoxystilbene Cap Phosphoramidite 44
- Carboxy-dC**
5-Carboxy-dC-CE Phosphoramidite 45
- Carboxy-Modifiers**
5'-Carboxy-Modifier C5 72
5'-Carboxy-Modifier C10 72
Carboxy-dT 73
- Chain Terminators 50,51**
- Chelates**
2,2'-Dipicolylamine Phosphoramidite 111
EDTA-C2-dT-CE Phosphoramidite 110
- Chemical Phosphorylation 78**
- Cholesterol Labelling**
3'-Cholesteryl-TEG CPG 108
5'-Cholesteryl-TEG Phosphoramidite 108
Cholesteryl-TEG Phosphoramidite 71,108
- CleanAmp™ Technology**
CleanAmp™ Primers 49
- Click Chemistry 83**
1,2,3-triazoles 83
3'-Alkyne-Modifier Serinol CPG 85,90
3'-Propargyl-5-Me-dC CPG 60,84
5'-Bromohexyl Phosphoramidite 85
5'-Ethynyl-dU-CE Phosphoramidite 84,124
5'-Hexynyl Phosphoramidite 85
5'-I-dT-CE Phosphoramidite 85
Alkyne-Modifier Serinol Phosphoramidite 85,90
Alkyne-NHS Ester 85
Azides 85
Azidobutyrte NHS Ester 85
baseclick Oligo-Click-M-Biotin 84
baseclick Oligo-Click-M-Fluorescein 84
baseclick Oligo-Click-M-Reload 84
baseclick Oligo-Click-M-TAMRA 84
C8-Alkyne-dC-CE Phosphoramidite 83,84
C8-Alkyne-dT-CE Phosphoramidite 83
C8-TIPS-Alkyne-dC-CE Phosphoramidite 83,84
C8-TIPS-Alkyne-dT-CE Phosphoramidite 84
C8-TMS-Alkyne-dC-CE Phosphoramidite 83,84
C8-TMS-Alkyne-dT-CE Phosphoramidite 84
Click DNA and RNA Ligation 60
Copper-free Click Chemistry 86
THPTA Ligand 84
- Convertible 2-dG**
2-F-dl-CE Phosphoramidite 63
- Convertible dA**
06-Phenyl-dl-CE Phosphoramidite 63
- Convertible dU**
04-Triazolyl-dU-CE Phosphoramidite 63
- Convertible F-dC**
TMP-F-dU-CE Phosphoramidite 63
- Convertible Nucleosides 63**
- Copper-free Click Chemistry 84**
5'-DBCO-TEG Phosphoramidite 86
DBCO-dT-CE Phosphoramidite 86
DBCO-sulfo-NHS Ester 86
- Cross-linking 53,55,88,109**
- Custom Doping 46**
- Cyanine Labelling**
Cyanine 3.5 Phosphoramidite 100
Cyanine 3 CPG 101
Cyanine 3 Phosphoramidite 100
Cyanine 5.5 Phosphoramidite 100
Cyanine 5 CPG 101
Cyanine 5 Phosphoramidite 100
Disulfo-Cyanine 7 Azide 88,101
DyLight DY547 Phosphoramidite 101
DyLight DY647 Phosphoramidite 101
- Cyclo-dA**
5',8-Cyclo-dA CE Phosphoramidite 59
- Cyclo-dG**
5',8-Cyclo-dG CE Phosphoramidite 59
- Cytosine Arabanoside 67**
- D**
- Dabcyl Labelling**
3'-Dabcyl CPG 91
3'-Dabcyl PS 9,91
3'-Dabsyl CPG 66,91,104,106
5'-Dabcyl Phosphoramidite 91
Dabcyl-dT 91
- DBCO**
5'-DBCO-TEG Phosphoramidite 86
DBCO-dT-CE Phosphoramidite 86
DBCO-sulfo-NHS Ester 86
- DCI (4,5-Dicyanoimidazole) 28**
- DDTT, See Sulfurizing Reagent II**
- Deaza-5-aza-C**
2'-OMe-3-deaza-5-aza-C-CE Phosphoramidite 129
- Deaza-8-aza-A**
7-deaza-8-Aza-A-CE Phosphoramidite 123
7-Deaza-8-aza-dA-CE Phosphoramidite 52

- Deaza-8-aza-G
7-deaza-8-Aza-dG-CE Phosphoramidite 52
- Deaza-A
3-Deaza-dA-CE Phosphoramidite 52
7-Deaza-A-CE Phosphoramidite 123
7-Deaza-dA-CE Phosphoramidite 52
- Deaza-G
7-Deaza-dG-CE Phosphoramidite 52
- Deaza-X
7-Deaza-dX-CE Phosphoramidite 50 52
- Dendrimers
Doubler Phosphoramidite 81
Trebler Phosphoramidite 81
- Depurination-resistant dA 58
- Desthiobiotin
DesthiobiotinTEG-CPG 94
DesthiobiotinTEG Phosphoramidite 93
- Deuterated Nucleosides 55
- Diaminopurine 42,120,128
- Dicyanoimidazole 28
- Dideoxynucleoside, 2',3'- 50
- Dideoxynucleosides
2',3'-ddC-CPG 51
2',3'-ddA-CE Phosphoramidite 51
2',3'-ddC-CE Phosphoramidite 51
2',3'-ddG-CE Phosphoramidite 51
2',3'-ddT-CE Phosphoramidite 51
- Dihydro-dT
5,6-Dihydro-dT-CE Phosphoramidite 57
- Dihydro-dU
5,6-Dihydro-dU-CE Phosphoramidite 57
- Dipicolylamine
2,2'-Dipicolylamine Phosphoramidite 111
- Distributors 4
- Dithiol Phosphoramidite 72
- DNA Damage/Repair 56,58,59
- DNA Methylation
5-Carboxy-dC-CE Phosphoramidite 45
5-Formyl-dC-CE Phosphoramidite 45
5-Me-dC-CE Phosphoramidite 42
- DNP Labelling
DNP-TEG Phosphoramidite 107
- Doubler Phosphoramidite
Symmetric 81
- Dr. Oligo synthesizer
CE Phosphoramidites 18
Solvents and Reagents 18
Supports and Columns 19
- Ds
dDs-CE Phosphoramidite 48
- Dss
dDss-CE Phosphoramidite 48
- DTPA
3'-DTPA CPG 76
Dithiol Phosphoramidite 72
- Duplex Stabilization 42,43,44
- DyLight™ Dyes
DyLight DY547 Phosphoramidite 101
DyLight DY647 Phosphoramidite 101
- E**
- Eclipse™ Quencher
Eclipse™ Quencher CPG 103
Eclipse™ Quencher Phosphoramidite 102
- EDTA-dT
EDTA-C2-dT-CE Phosphoramidite 110
- Epigenetics 45,124
- Epoch Dyes and Quencher
Epoch Eclipse™ Quencher CPG 103
Epoch Eclipse™ Quencher Phosphoramidite 102
Epoch Gig Harbor Green™ Phosphoramidite 102
Epoch Redmond Red™ CPG 103
Epoch Redmond Red™ Phosphoramidite 102
Epoch Yakima Yellow™ CPG 103
Epoch Yakima Yellow™ Phosphoramidite 102
- Et-dC-CE Phosphoramidite 43
- Etheno-A
Etheno-dA-CE Phosphoramidite 64
- Ethylthiotetrazole 28
- Excimers 111
- Expedite Instruments
CE Phosphoramidites 10
Solvents and Reagents 10
Supports and Columns 11
- F**
- FAM
3'-(6-FAM) CPG 97
3'-(6-FAM) PS 97
6-FAM 95
6-FAM-TEG Azide 87
Dipivaloyl 6-FAM-TEG Azide 87
- F-ANA Monomers
2'-F-A-ANA CE Phosphoramidite 131
2'-F-Bz-C-ANA CE Phosphoramidite 131
2'-F-G-ANA CE Phosphoramidite 131
2'-F-U-ANA CE Phosphoramidite 131
- F-Arabinonucleic Acid
2'-F-Arabinonucleic Acid 131
- F-C
2'-F-Ac-C-CE Phosphoramidite 44
2'-OMe-5-F-C Precursor 129
F-dC Precursor 63
- F-dI
2-F-dI-CE Phosphoramidite 63
- F-dU
5-F-dU-CE Phosphoramidite 55
- Ferrocene Labelling
Ferrocene-dT-CE Phosphoramidite 110
- Filters
AB Columns
Replacement Frits (10µm) 8

- Expedite Columns
 Replacement Filters (40nm, 0.2 or 1µm) 12
 Replacement Frits (15µm) 12
- Fluorescein Labelling**
 3'-(6-FAM) CPG 97
 3'-(6-FAM) PS 9,97
 3'-(6-Fluorescein) CPG 97,101
 3'-6-Fluorescein Serinol CPG 90,97
 3'-Fluorescein CPG 97
 3'-Fluorescein-dT CPG 97
 5'-Dichloro-dimethoxy-Fluorescein 95
 5'-Fluorescein Phosphoramidite 95
 5'-Hexachloro-Fluorescein 95
 5'-Tetrachloro-Fluorescein 95
 6-Fluorescein Phosphoramidite 96
 6-Fluorescein Serinol Phosphoramidite 89,96
 Dichloro-diphenyl-fluorescein 98
 Fluorescein-dT Phosphoramidite 96
 SIMA (HEX) 98
- Fluorescent Nucleosides 64**
 2-Aminopurine-CE Phosphoramidite 53
 5-Me-2'-deoxyZebularine-CE Phosphoramidite 67
 AP-dC-CE Phosphoramidite 42
 Etheno-dA-CE Phosphoramidite 64
 Perylene-dU-CE Phosphoramidite 65
 Pyrene-dU-CE Phosphoramidite 65
 Pyrrolo-C-TOM-CE Phosphoramidite 121
 Pyrrolo-CTP 124
 Pyrrolo-dC-CE Phosphoramidite 64
 Pyrrolo-dCTP 64
 tC-CE Phosphoramidite 66
 tC°-CE Phosphoramidite 66
- Fluoro-Pak™ Purification**
 Fluoro-Pak™ columns 140
 Fluoro-Pak™ II columns 140
 Loading Buffer 140
 Phosphoramidites 140
- Formyl-dC**
 5-Formyl-dC-CE Phosphoramidite 45
 5-Formyl-dC III-CE Phosphoramidite 45
- Formylindole CE Phosphoramidite 79**
- F-RNA Monomers**
 2'-F-Ac-C-CE Phosphoramidite 130
 2'-F-A-CE Phosphoramidite 130
 2'-F-G-CE Phosphoramidite 130
 2'-F-U-CE Phosphoramidite 130
- F-U**
 2'-OMe-5-F-U-CE Phosphoramidite 129
 2'-OMe-TMP-5-F-U-CE Phosphoramidite 129
- Free Radicals 57**
- G**
- 2'-F-G-ANA CE Phosphoramidite 131
 2'-F-G-CE Phosphoramidite 130
 2'-OMe-G-CE Phosphoramidite 125
 2'-OMe-G-RNA 127
 2'-OMe-iBu-G-CE Phosphoramidite 125
 2'-OMe-iPr-Pac-G-CE Phosphoramidite 126
- 2',3'-ddG 51
 3'-dG-CE Phosphoramidite 61
 3'-dG-CPG 50,62
 6-Thio-dG-CE Phosphoramidite 53
 6-Thio-G-CE Phosphoramidite 122
 7-Deaza-8-aza-dG-CE Phosphoramidite 52
 7-Deaza-dG-CE Phosphoramidite 52
 8-Amino-dG-CE Phosphoramidite 58
 8-Br-dG-CE Phosphoramidite 55
 8-D-dG-CE Phosphoramidite 56
 8-Oxo-dG-CE Phosphoramidite 57
 Ac-G-CE Phosphoramidite 117
 Ac-G-RNA-CPG 116,119
 CleanAmp™-Pac-dG-CE Phosphoramidite 49
 dG-CE Phosphoramidite 6,10,13,14,16,18
 dG-H-Phosphonate 35
 dG-Me Phosphonamidite 32
 dG-Me Phosphoramidite 34
 dG-PACE Phosphoramidite 33
 dG-Thiophosphoramidite 36
 dmf-dG-5'-CE Phosphoramidite 30
 dmf-dG-CE Phosphoramidite 6,10,13,14,16,18
 G-TOM-CE Phosphoramidite 115,117
 iPr-Pac-dG-CE Phosphoramidite 20,68
 iPr-Pac-G-RNA-CPG 119
 N2-Amino-Modifier C6 dG 73
 O6-Me-dG-CE Phosphoramidite 62
- G-Clamp 42,64**
- GE Healthcare Life Sciences**
 CE Phosphoramidites 16
 Solvents and Reagents 17
- Gig Harbor Green™**
 Gig Harbor Green™ Phosphoramidite 102
- Glen Gel-Pak™ Purification**
 Glen Gel-Pak™ 141
- Glen-Pak™ Purification**
 Adapter Rack 138
 Glen-Pak™ DNA Purification Cartridge 138
 Glen-Pak™ RNA Purification Cartridge 138
 RNA Quenching Buffer 138
 Seal for Adapter Rack 138
- Glen UnySupport™**
 Glen UnySupport CPG 22,23
 Glen UnySupport FC CPG 23
 Glen UnySupport HybridCPG 25
 Glen UnySupport PS 9,22,23
- Glyceryl CPG 76**
- Gold Surface 72**
- H**
- HEX**
 6-HEX Azide 87,88
- Hexynyl Phosphoramidite 85**
- High Throughput**
 HT 2'-OMe-RNA Phosphoramidites 125
 HT DNA Phosphoramidites 13
 HT RNA Phosphoramidites 117

H-Phosphonate Chemistry

- Monomers 35
- Reagents for ABI synthesizers 35

Halogenated Nucleosides

- 2'-OMe-5-F-U-CE Phosphoramidite 129
- 2'-OMe-TMP-5-F-U-CE Phosphoramidite 129
- 5-Br-dC-CE Phosphoramidite 55
- 5-Br-dU-CE Phosphoramidite 55
- 5-Br-dU-CPG 55
- 5-Br-U-CE Phosphoramidite 122
- 5-F-dU-CE Phosphoramidite 55
- 5-I-dC-CE Phosphoramidite 55
- 5-I-dU-CE Phosphoramidite 55
- 5-I-U-CE Phosphoramidite 122
- 8-Br-dA-CE Phosphoramidite 55
- 8-Br-dG-CE Phosphoramidite 55

HEX 95**High Load CPG 27****HybridCPG™ 25****Hydrogen Bonding 52****Hydroxy-C**

- 5-OH-dC-CE Phosphoramidite 57

Hydroxymethyl-dC

- 5-Hydroxymethyl-dC-CE Phosphoramidite 45
- 5-Hydroxymethyl-dC II-CE Phosphoramidite 45

Hydroxy-U

- 5-OH-dU-CE Phosphoramidite 57

Hydroxymethyl-dU

- 5-Hydroxymethyl-dU-CE Phosphoramidite 57

I

- 2'-OMe-I-CE Phosphoramidite 129
- 2-F-dI-CE Phosphoramidite 63
- dI-CE Phosphoramidite 46
- dI-CPG 46
- I-CE Phosphoramidite 122
- O6-Phenyl-dI-CE Phosphoramidite 63

I-dC

- 5-I-dC-CE Phosphoramidite 55

I-dT

- 5'-I-dT-CE Phosphoramidite 85

I-dU

- 5-I-dU-CE Phosphoramidite 55

isodC

- dmf-5-Me-isodC-CE Phosphoramidite 48

isodG

- dmf-isodG-CE Phosphoramidite 48

I-U

- 5-I-U-CE Phosphoramidite 122

Introduction 4**Ionizing Radiation 57****Isopropyl Phosphite 35****J****JOE**

- 5'-Dichloro-dimethoxy-Fluorescein Phosphoramidite II 95

K**K**

- dK-CE Phosphoramidite 47

L**Labelling of MicroRNAs 60****Large Scale Synthesis**

- N3-Cyanoethyl-dT 67

Low Cost

- LC RNA Phosphoramidites 117

M**Maleimide-Modifier**

- 5'-Maleimide-Modifier Phosphoramidite 72

Me-C

- 2'-OMe-5-Me-C-CE Phosphoramidite 129
- 3'-Propargyl-5-Me-dC CPG 60
- 5-Me-C-TOM-CE Phosphoramidite 120
- 5-Me-dC-CE Phosphoramidite 42
- 5-Me-dC-CPG 42
- Ac-5-Me-dC-CE Phosphoramidite 42

Methylated Nucleosides

- 1-Me-A-CE Phosphoramidite 124
- 1-Me-dA-CE Phosphoramidite 62
- N6-Ac-N6-Me-dA-CE Phosphoramidite 43,62
- N6-Me-A-CE Phosphoramidite 124
- N6-Me-dA-CE Phosphoramidite 43,62
- O4-Me-dT-CE Phosphoramidite 43,62
- O6-Me-dG-CE Phosphoramidite 62

Methylene Blue

- Methylene Blue C3 Phosphoramidite 110
- Methylene Blue NHS Ester 110

Me-U

- 2'-OMe-5-Me-U-CE Phosphoramidite 128
- 5-Me-U-CE Phosphoramidite 122

MerMade Instruments

- CE Phosphoramidites 14
- Solvents and Reagents 14,15
- Supports 15

Methyl Phosphoramidites 32**MicroRNA 60****Minor 2'-OMe-RNA Phosphoramidites 128-129****Minor Groove 53****Mixed Base Combinations 46****Modifiers 70,71,74****Mutagenesis 62****N****NAIM**

- A Analogs 133,134

Nebularine

- 2'-DeoxyNebularine-CE Phosphoramidite 46
- Nebularine-CE Phosphoramidite 123

Nitroindole

- 5-Nitroindole-CE Phosphoramidite 47

O

Oligo-Affinity Support

OAS PS 142

OMe-RNA Synthesis

2'-OMe-RNA Phosphoramidites 125

2'-OMe-RNA Supports 127

Minor 2'-OMe-RNA Phosphoramidites 128,129

UltraMild 2'-OMe-RNA 125

OMe-T 128

Oxo-dA

8-Oxo-dA-CE Phosphoramidite 57

Oxo-dG

8-Oxo-dG-CE Phosphoramidite 57

Oxo-G Clamp-CE Phosphoramidite

65

P

P

dP-CE Phosphoramidite 47,49

Pa

Biotin PaTP 48

dPa-CE Phosphoramidite 48

PACE Phosphoramidites 33

PCR/Sequencing Utilities 46

Perylene

Perylene-dU-CE Phosphoramidite 65,111

Phenothiazine

tC-CE Phosphoramidite 66

Phenoxazine

tC^o-CE Phosphoramidite 66

Phosphonocarboxylate Monomers 33

Phosphorylation

3'-CPR II CPG 78

3'-Phosphate CPG 78

3'-Phosphate CPG - High Load 78

3'-Phosphate PS 78

3'-Phosphate PS 9

Chemical Phosphorylation Reagent 78

Chemical Phosphorylation Reagent II 78

CPR II 78

Solid CPR II 78

Photoaffinity Labelling 53

Photocleavable Monomers

PC Amino-Modifier Phosphoramidite 72,82

PC Biotin Phosphoramidite 82,93

PC Linker Phosphoramidite 82

PC Spacer Phosphoramidite 80,82

Photo cross-linking 53

Photo-responsive DNA

Azobenzene Phosphoramidite 113

Phthalimide (PT)

3'-PT-Amino-Modifier C3 CPG 75

3'-PT-Amino-Modifier C6 CPG 75

3'-PT-Amino-Modifier C6 PS 75

3'-PT-Amino-Modifier C6 PS 9

Poly-Pak™ Purification

Poly-Pak™ Cartridge 139

Poly-Pak™ II Cartridge 139

Poly-Pak™ Packing 138,139

Reagents 138,139

Polystyrene Supports

3'-(6-FAM) PS 9,97

3'-BiotinTEG PS 9,94

3'-Dabcyl PS 9,91

3'-Phosphate PS 9,78

3'-PT-Amino-Modifier C6 PS 9,75

3'-TAMRA PS 9,99

AB 3900 Polystyrene Columns 8,19

Glen UnySupport PS 9,22

Low Volume (LV) Polystyrene Columns 8

Universal Support III PS 9,24

PPG 52

Propyne Derivatives

pdC-CE Phosphoramidite 42

pdU-CE Phosphoramidite 42

Protein-DNA Interaction 53

PseudoU

2'-deoxypseudoU-CE Phosphoramidite 53

Psoralen Labelling

Psoralen Azide 88,109

Psoralen C2 Phosphoramidite 109

Psoralen C6 Phosphoramidite 109

Purification

Fluoro-Pak™ Purification 140

Glen-Pak™ Purification 138

Poly-Pak™ Purification 139

Purine

Purine 47

Purine Ribonucleoside 123

Puromycin

Puromycin CPG 112

Pyrene

5'-Pyrene Cap Phosphoramidite 44

Pyrene-dU-CE Phosphoramidite 65,66,111

Pyridin-2-one-CE Phosphoramidite 123

Pyrrolidine-CE Phosphoramidite 59

Pyrrolo-C

Pyrrolo-C-TOM-CE Phosphoramidite 121

Pyrrolo-CTP 124

Pyrrolo-dC-CE Phosphoramidite 64

Pyrrolo-dCTP 64

Q

Q-Supports 25,26

Quenched Autoligation (QUAL) Probes 112

5'-Dabsyl-dT-CE Phosphoramidite 112

R

Redmond Red

Redmond Red™ CPG 103

Redmond Red™ Phosphoramidite 102

Repair Enzyme 57

Rhodamine 99

RNA Supports

for 3' DNA Modification 114

RNA Synthesis

Minor RNA Phosphoramidites 119,124

RNA Phosphoramidites 117,118

RNA Supports 118,119

RNA Supports for TOM-RNA Synthesis 115,116

TOM-Protected Minor RNA Phosphoramidites 116,120,122

TOM-Protected RNA Phosphoramidites 115

S**s**

Pac-ds-CE Phosphoramidite 48

sTP 48

Saccharin 1-Methylimidazole 28**Sequence Modifiers 74****Serinol Backbone**

3'-6-Fluorescein Serinol CPG 90,97

3'-Alkyne-Modifier Serinol CPG 85,90

3'-Amino-Modifier Serinol CPG 75,90

3'-Protected BiotinLC Serinol CPG 90,94

3'-Protected Biotin Serinol CPG 90,94

6-Fluorescein Serinol Phosphoramidite 89,96

Alkyne-Modifier Serinol Phosphoramidite 85,90

Amino-Modifier Serinol Phosphoramidite 74,89

Protected BiotinLC Serinol Phosphoramidite 89,93

Protected Biotin Serinol Phosphoramidite 89,93

Silanized Bottles 37**SIMA**

SIMA (HEX)-dT Phosphoramidite 98

SIMA (HEX) Phosphoramidite 98

SMI 28**Spacer Modifiers**

3'-Spacer C3 CPG 80

dSpacer CE Phosphoramidite 80

PC Spacer Phosphoramidite 80,82

rSpacer CE Phosphoramidite 121

Spacer C12 CE Phosphoramidite 80

Spacer Phosphoramidite 18 80

Spacer Phosphoramidite 9 80

Spacer Phosphoramidite C3 80

Spermine Phosphoramidite 43**Spin Labels**

TEMPO Azide 88

TEMPO-TEG Azide 88

Stearyl Labelling 108

5'- Stearyl Phosphoramidite 108

Sterling

Introduction 5

Structural Studies 52**Structure/Activity Relationship 52****Sulfurizing Reagent 37****Sulfurizing Reagent II 37****Synthesis Columns**

ABI

Empty 8

Expedite

Empty 12

MerMade

Empty 15

T**T**

2-Thio-dT-CE Phosphoramidite 53

2',3'-ddT 51

3'-Amino-dT CPG 60

3'-Amino-Modifier C6 dT CPG 77

3'-dT-CE Phosphoramidite 61

3'-dT-CPG 50,62

3'-Fluorescein-dT CPG 97

4-Thio-dT-CE Phosphoramidite 53

5'-Amino-dT-CE Phosphoramidite 50

5'-Dabsyl-dT 112

5'-I-dT-CE Phosphoramidite 85

5'-OMe-dT-CE Phosphoramidite 50

5,6-Dihydro-dT-CE Phosphoramidite 57

Amino-Modifier C2 dT 73

Amino-Modifier C6 dT 73

C8-Alkyne-dT-CE Phosphoramidite 83

C8-TIPS-Alkyne-dT-CE Phosphoramidite 84

C8-TMS-Alkyne-dT-CE Phosphoramidite 84

CleanAmp™-dT-CE Phosphoramidite 49

DBCO-dT-CE Phosphoramidite 86

dT-CE Phosphoramidite 6,10,13,14,16,18

dT-H-Phosphonate 35

dT-Me Phosphoramidite 34

dT-Me Phosphonamide 32

dT-PACE Phosphoramidite 33

dT-Thiophosphoramidite 36

EDTA-C2-dT-CE Phosphoramidite 110

Ferrocene-dT-CE Phosphoramidite 110

Fluorescein-dT 96

N3-Cyanoethyl-dT 67

NPOM-Caged-dT 66

O4-Me-dT-CE Phosphoramidite 43,62

S-Bz-Thiol-Modifier C6-dT 74

TAMRA-dT 99

Thymidine Glycol CE Phosphoramidite 58

TAMRA Labelling

3'-TAMRA CPG 99

3'-TAMRA PS 99

3'-TAMRA PS 9

TAMRA-dT 99

TAMRA NHS Ester 99

tC

tC-CE Phosphoramidite 66

tCnitro

tCnitro-CE Phosphoramidite 66

tCo

tC°-CE Phosphoramidite 66

TEMPO

TEMPO Azide 88

TEMPO-TEG Azide 88

Termination, 3'

- 2',3'-ddC-CPG 51
- 2',3'-ddA 51
- 2',3'-ddC 51
- 2',3'-ddG 51
- 2',3'-ddT 51
- 3'-3' linkage 31,50
- 3'-dA-CPG 50
- 3'-dC-CPG 50
- 3'-dG-CPG 50
- 3'-dT-CPG 50
- 3'-Spacer C3 CPG 80

Termination, 5'

- 5'-OMe-dT-CE Phosphoramidite 50

Terminus Modifiers 70**TET 95**

- 6-TET Azide 87,88

Thio-dT

- 2-Thio-dT-CE Phosphoramidite 53
- 4-Thio-dT-CE Phosphoramidite 53

Thio-dU

- 4-Thio-dU-CE Phosphoramidite 53

Thio-G

- 6-Thio-dG-CE Phosphoramidite 53
- 6-Thio-G-CE Phosphoramidite 122

Thiol-Modifiers

- 3'-DTPA CPG 76
- 3'-Thiol-Modifier 6 S-S CPG 76
- 5'-Thiol-Modifier C6 72
- Dithiol Phosphoramidite 72
- DTPA 72
- S-Bz-Thiol-Modifier C6-dT 74
- Thiol-Modifier C6 S-S 72

Thiophosphoramidites 36**Thio-U**

- 4-Thio-U-TOM-CE Phosphoramidite 120

Thiotriphosphate Nucleotides

- 2'-deoxyAdenosine 134
- 2'-deoxyCytidine 135
- 2'-deoxyGuanosine 135
- 2'-deoxyUridine 135
- 2'-Fluoro-Adenosine 134
- 2'-OMe-Adenosine 134
- 2,6-Diaminopurine riboside 134
- 2-Aminopurine riboside 134
- 5-Methyl-Uridine 135
- 7-Deaza-Adenosine 133
- Adenosine 133
- Cytidine 135
- Guanosine 135
- Inosine 135
- N6-Me-Adenosine 133
- Purine riboside 134
- Thymidine 135
- Uridine 135

Thymidine Glycol

- Thymidine Glycol CE Phosphoramidite 58

Thymine Dimer

- Cis-syn Thymine Dimer Phosphoramidite 59

Tocopherol

- a-Tocopherol-TEG Phosphoramidite 108

TOM-Protecting-Group

- Ac-A-RNA-CPG 115
- Ac-C-RNA-CPG 116
- Ac-G-RNA-CPG 116
- A-TOM-CE Phosphoramidite 115,117
- C-TOM-CE Phosphoramidite 115,117
- G-TOM-CE Phosphoramidite 115,117
- U-RNA-CPG 116
- U-TOM-CE Phosphoramidite 115,117

Trebler Phosphoramidite

- Long Trebler 81
- Trebler 81

Trimer Phosphoramidites 38**Trimethoxystilbene**

- 5'-Trimethoxystilbene Cap Phosphoramidite 44

Triphosphate Nucleotides

- Biotin PaTP 48
- Pyrrolo-dCTP 64
- sTP 48

Triplex 52**TWIST™ Columns**

- Applied Biosystems 40nm, 0.2µm, 1µm, 10µm
- Empty 8
- Expedite 15µm
- Empty 12

U**U**

- 2'-F-U-ANA CE Phosphoramidite 131
- 2'-OMe-5-Br-U-CE Phosphoramidite 129
- 2'-OMe-5-F-U-CE Phosphoramidite 129
- 2'-OMe-5-Me-U-CE Phosphoramidite 128
- 2'-OMe-TMP-5-F-U-CE Phosphoramidite 129
- 2'-OMe-U-CE Phosphoramidite 125
- 2'-OMe-U-RNA 127
- 3'-Uaq Cap CPG 44,60
- 4-Thio-dU-CE Phosphoramidite 53
- 5,6-Dihydro-dU-CE Phosphoramidite 57
- 5-Br-dU-CE Phosphoramidite 55
- 5-Br-dU-CPG 55
- 5-Ethynyl-dU-CE Phosphoramidite 84,124
- 5-F-dU-CE Phosphoramidite 55
- 5-Hydroxymethyl-dU-CE Phosphoramidite 57
- 5-I-dU-CE Phosphoramidite 55
- 5-I-U-CE Phosphoramidite 122
- 5-OH-dU-CE Phosphoramidite 57
- Amino-Modifier C6-U Phosphoramidite 121
- Br-U-CE Phosphoramidite 122
- dU-CE Phosphoramidite 46
- dU-CPG 1000 46
- dU-CPG 500 46
- 04-Triazolyl-dU-CE Phosphoramidite 63
- pdU-CE Phosphoramidite 42
- Perylene-dU-CE Phosphoramidite 65

Pyrene-dU-CE Phosphoramidite 65,66
 TMP-F-dU-CE Phosphoramidite 63
 U-CE Phosphoramidite 117,118
 U-RNA-CPG 114,116,119
 U-TOM-CE Phosphoramidite 115,117

UltraMILD Deprotection

2'-OMe-Ac-C-CE Phosphoramidite 126
 2'-OMe-iPr-Pac-G-CE Phosphoramidite 126
 2'-OMe-Pac-A-CE Phosphoramidite 126
 Ac-C-CE Phosphoramidite 118
 Ac-dC-CE Phosphoramidite 20,68
 Cap Mix A 21,34,69,119,126
 iPr-Pac-dG-CE Phosphoramidite 20,68
 iPr-Pac-G-CE Phosphoramidite 118
 Pac-A-CE Phosphoramidite 118
 Pac-dA-CE Phosphoramidite 20,68
 Potassium Carbonate in Methanol 21,34,69,119,126

UniCap Phosphoramidite 29

Universal Support

Glen UnySupport FC 23
 Glen UnySupport™ 22,25
 Original Universal Support 22
 UnyLinker™ 22

Universal Support III

Universal Support III CPG 24
 Universal Support III HybridCPG™ 25
 Universal Support III PS 9,24

Unnatural Base Pairs

5-Me-isodC 48
 Biotin PaTP 48
 dDs 48
 dDss 48
 dPa 48
 ds 48
 isodG 48
 sTP 48

V

Vitamin E 108

X

X

2'-dX-CE Phosphoramidite 54
 7-deaza-dX-CE Phosphoramidite 52

Y

Yakima Yellow

Yakima Yellow™ CPG 103
 Yakima Yellow™ Phosphoramidite 102

Z

Zebularine

5-Me-2'-deoxyZebularine-CE Phosphoramidite 67
 Zebularine-CE Phosphoramidite 123

Zip Nucleic Acid 43

Spermine Phosphoramidite 43

ZNA 43

Spermine Phosphoramidite 43

ORDERING

Orders may be placed by mail, telephone, fax or email to:

Glen Research Corporation
22825 Davis Drive, Sterling, VA 20164, USA
(800) 327-GLEN (USA)
(703) 437-6191 (International)
(800) 934-2490 (Fax USA)
(703) 435-9774 (Fax)
orders@glenres.com (Internet email)
<http://www.glenres.com> (World Wide Web Home Page)

There is no minimum order requirement. Our office hours are 9:30 a.m. to 6:00 p.m. EST, Monday through Friday. During this time, Glen Research representatives will be available to take your order and answer any questions you may have regarding our product line or DNA synthesis in general. Messages may be left on voice mail at any other time. Orders placed by 4:00 p.m. EST are normally shipped the same day.

DISCOUNTS

Glen Research offers a 10% discount on catalog products to educational institutions. In addition, discounts based on volume of usage are available, so please request a specific quotation. We are also happy to quote on bulk quantities or custom packaging of our products and on custom synthesis of unique materials.

TERMS AND CONDITIONS OF SALE

Delivery will be F.O.B. Glen Research, Sterling, VA. Freight charges will be prepaid by Glen Research and will be included on the invoice sent to the Customer.

Glen Research will issue an invoice to the Customer upon shipment of Products which shall be due and payable in full within thirty (30) days of the date of the invoice. Amounts not paid when due may incur an interest charge of one and one-half (1-1/2) percent per month until fully paid. Customer shall be responsible for all taxes on Products sold to it, including state and local sales and use taxes. We reserve the right to require payment in advance before shipping any materials ordered.

Glen Research warrants that each product conforms to the specifications for such product. If the Customer notifies Glen Research within thirty (30) days of its receipt of a product that the product does not conform to the specifications, Glen Research will, at its option, replace the product or return the purchase price paid by Customer. No replacement or refund will be made if the Customer does not notify Glen Research of a non-conforming product within said thirty (30) day period.

Glen Research makes no other express or implied warranty with respect to the products. Glen Research disclaims all implied warranties, including the warranties of merchantability and fitness for a particular purpose. Glen Research's liability for breach of warranty is limited to refund of purchase price paid. In no event shall Glen Research be liable for lost profits, loss of goodwill or any incidental, special or consequential damages even if advised of the possibility of same.

It is the responsibility of the Buyer to determine the suitability of any of the listed products for any specific purpose. Responsibility for accidents arising from the handling and use of the listed products rests solely with the Buyer. All of the products should be handled only by trained personnel who are aware of the potential hazards and have ready access to suitable safety equipment. All products supplied are primarily for research and development purposes and Glen Research products are not intended for use in foods or drugs, or for any commercial purpose.

PATENTS

As a research-oriented company, we realize the desirability of patents to cover original research and it is our policy to avoid infringing any approved patents. Accordingly, it is possible that some of our products may have to be withdrawn or adjusted in price as patents are approved and issued.

©2015 Glen Research Corporation